

The Efroimsky formalism for weak gravitational waves adapted to high-frequency perturbations

O Svítek and J Podolský

Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles University
in Prague, V Holešovičkách 2, 180 00 Praha 8, Czech Republic

E-mail: ota@matfyz.cz and podolsky@mbox.troja.mff.cuni.cz

Received 11 November 2003

Published 5 July 2004

Online at stacks.iop.org/CQG/21/3579

doi:10.1088/0264-9381/21/14/017

Abstract

The Efroimsky perturbation scheme for consistent treatment of gravitational waves and their influence on the background is summarized and compared with the classical Isaacson high-frequency approach. We demonstrate that the Efroimsky method in its present form is not compatible with the Isaacson limit of high-frequency gravitational waves, and we propose its natural generalization to resolve this drawback.

PACS numbers: 04.30.-w, 04.25.-g

1. Introduction

Recently, Efroimsky introduced and developed a new formalism for the consistent treatment of weak gravitational waves [1, 2]. This interesting mathematical framework is remarkable, mainly due to the possibility of ascribing the stress–energy tensor even to low-frequency gravitational waves influencing the background, which is in contrast to the standard linearization approach where the background is kept fixed. This is achieved by introducing a natural low-frequency cut-off, employing three different metrics (the premetric, the complete physical metric and the average metric) and careful analysis of their mutual relations.

On the other hand, in a now classic paper [3] Isaacson (inspired by previous works [4, 5]) presented a perturbation method which can be used for studies of high-frequency gravitational waves. Such waves also influence the cosmological background in which they propagate. Isaacson's work stimulated further contributions in which his method was reformulated using various formalisms, and explicitly applied to particular spacetimes (see, e.g., [6–15]).

In our present work we first briefly summarize and compare the two above-mentioned perturbation schemes. In particular, it is shown that the Efroimsky method is not consistent if high-frequency gravitational waves are considered. Next (in section 3), we propose a possible modification of the Efroimsky formalism which may resolve this drawback.

2. The formalism

Efroimsky's approach [1, 2] is based on introducing three different smooth, non-degenerate, symmetric metrics on a differentiable manifold M , namely:

- (i) $\gamma_{\mu\nu}$ —the 'premetric', a vacuum metric corresponding to initial pure background without gravitational waves;
- (ii) $g_{\mu\nu}$ —the 'physical metric', a full vacuum metric which describes both the background and the waves;
- (iii) $q_{\mu\nu}$ —the 'average metric', a nonvacuum metric representing the background plus its perturbations with wavelength greater than L . In fact, it is the averaged full metric $g_{\mu\nu}$, where the cut-off value L depends on the observer's experimental abilities. Since no detector can measure gravitational waves of arbitrarily long wavelengths, the existence of such a low-frequency cut-off is a natural assumption.

One motivation for using these three distinct metrics is to resolve a (slight) discrepancy in the standard linearization approach which considers only the metrics $\gamma_{\mu\nu}$, $g_{\mu\nu}$, and decomposition $g_{\mu\nu} = \gamma_{\mu\nu} + h_{\mu\nu}$, where $h_{\mu\nu}$ is a small perturbation. The contravariant components obtained as an inverse of $g_{\mu\nu}$ are $g^{\mu\nu} = \gamma^{\mu\nu} - h^{\mu\nu} + O(h^2)$, but $\gamma_{\mu\nu}$ is commonly used for raising and lowering indices. It is thus not clear which semi-Riemannian manifold this equality relates to. Such inconsistency can be ignored in the lowest order because it leads to the correct linear approximation of the wave equation. To extend the weak-field formalism to higher-order terms, the distinction between the premetric γ and the average metric q is necessary as it exhibits the back-reaction of the waves on the background geometry. (Here and hereafter, indices of the metric tensors are sometimes suppressed for notational simplicity.)

The next step is to define the Ricci and Einstein tensors for an arbitrary metric g as

$$\begin{aligned} R_{\mu\nu}(g) &\equiv \left[\frac{1}{2} g^{\gamma\rho} (g_{\rho\nu,\mu} + g_{\rho\mu,\nu} - g_{\mu\nu,\rho}) \right]_{,\gamma} - \left[\frac{1}{2} g^{\gamma\rho} (g_{\rho\gamma,\mu} + g_{\rho\mu,\gamma} - g_{\mu\gamma,\rho}) \right]_{,\nu} \\ &\quad + \left[\frac{1}{2} g^{\gamma\delta} (g_{\rho\delta,\gamma} + g_{\rho\gamma,\delta} - g_{\gamma\delta,\rho}) \right] \left[\frac{1}{2} g^{\delta\rho} (g_{\rho\nu,\mu} + g_{\rho\mu,\nu} - g_{\mu\nu,\rho}) \right] \\ &\quad - \left[\frac{1}{2} g^{\gamma\rho} (g_{\rho\delta,\nu} + g_{\rho\nu,\delta} - g_{\nu\delta,\rho}) \right] \left[\frac{1}{2} g^{\delta\rho} (g_{\rho\gamma,\mu} + g_{\rho\mu,\gamma} - g_{\mu\gamma,\rho}) \right], \\ G_{\mu\nu}(g) &\equiv R_{\mu\nu}(g) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} R_{\alpha\beta}(g), \end{aligned} \quad (1)$$

where $g^{\rho\tau} = (g)_{\rho\tau}^{-1}$; the same expressions apply to γ and q . These equations remain a tensor even if we transfer to another semi-Riemann space (the reason is that covariant tensors are defined on a metric space rather than on some particular semi-Riemann one). From proposals (i)–(iii) it follows that $G_{\mu\nu}(\gamma) = 0 = G_{\mu\nu}(g)$, $G_{\mu\nu}(q) \neq 0$.

Now, the *differences* between the covariant components of the above metrics are introduced,

$$h_{\mu\nu} \equiv g_{\mu\nu} - q_{\mu\nu}, \quad \eta_{\mu\nu} \equiv q_{\mu\nu} - \gamma_{\mu\nu}. \quad (2)$$

It is necessary to specify the semi-Riemann space: for raising or lowering indices and for covariant differentiation, the averaged nonvacuum metric q will be used. Consequently, h and η are tensor fields on the semi-Riemann manifold (M, q) , i.e.

$$h^{\mu\nu} \equiv q^{\mu\alpha} q^{\nu\beta} h_{\alpha\beta}, \quad \eta^{\mu\nu} \equiv q^{\mu\alpha} q^{\nu\beta} \eta_{\alpha\beta}. \quad (3)$$

Treating $h_{\mu\nu}$ as a perturbation of the metric $q_{\mu\nu}$ the Ricci tensor (1) can be expanded in a power series

$$R_{\mu\nu}(g) = R_{\mu\nu}^{(0)}(q) + R_{\mu\nu}^{(1)}(q, h) + R_{\mu\nu}^{(2)}(q, h) + R_{\mu\nu}^{(3)}(q, h) + O(h^4), \quad (4)$$

where

$$\begin{aligned}
 R_{\mu\nu}^{(0)}(q) &\equiv R_{\mu\nu}(q), \\
 R_{\mu\nu}^{(1)}(q, h) &\equiv \frac{1}{2}q^{\rho\tau}(h_{\tau\mu;\nu\rho} + h_{\tau\nu;\mu\rho} - h_{\rho\tau;\mu\nu} - h_{\mu\nu;\rho\tau}), \\
 R_{\mu\nu}^{(2)}(q, h) &\equiv \frac{1}{2}\left[\frac{1}{2}h^{\rho\tau}{}_{;\mu}h_{\rho\tau;\nu} + h^{\rho\tau}(h_{\rho\tau;\mu\nu} + h_{\mu\nu;\rho\tau} - h_{\tau\mu;\nu\rho} - h_{\tau\nu;\mu\rho}) \right. \\
 &\quad \left. + h^\tau{}_{\nu;\rho}(h_{\tau\mu;\rho} - h_{\rho\mu;\tau}) - (h^{\rho\tau}{}_{;\rho} - \frac{1}{2}h^{\rho;\tau}) (h_{\tau\mu;\nu} + h_{\tau\nu;\mu} - h_{\mu\nu;\tau})\right].
 \end{aligned}
 \tag{5}$$

Analogously,

$$R_{\mu\nu}(\gamma) = R_{\mu\nu}^{(0)}(q) + R_{\mu\nu}^{(1)}(q, (-\eta)) + R_{\mu\nu}^{(2)}(q, (-\eta)) + O(\eta^3).
 \tag{6}$$

It is obvious that $R_{\mu\nu}^{(1)}(q, (-\eta)) = -R_{\mu\nu}^{(1)}(q, \eta)$ and $R_{\mu\nu}^{(2)}(q, (-\eta)) = R_{\mu\nu}^{(2)}(q, \eta)$. According to assumptions that both g and γ are vacuum metrics the following relation holds:

$$\begin{aligned}
 0 &= R_{\mu\nu}(g) - R_{\mu\nu}(\gamma) \\
 &= R_{\mu\nu}^{(1)}(q, h) + R_{\mu\nu}^{(2)}(q, h) + R_{\mu\nu}^{(1)}(q, \eta) + R_{\mu\nu}^{(3)}(q, h) + O(h^4) + O(\eta^2).
 \end{aligned}
 \tag{7}$$

At this point Efroimsky sets three assumptions:

Assumption 1. The perturbations h and η are small in the sense that the terms of the orders $O(h^4)$ and $O(\eta^2)$ are negligible.

Assumption 2. The perturbations η and h^2 are of the same order.

Assumption 3. The tensor field h consists of modes with short wavelengths which do not exceed the given maximal value L .

A physical interpretation of the perturbations given by (2) is thus the following: $h_{\mu\nu}$ characterizes measurable gravitational waves whereas $\eta_{\mu\nu}$ is a shift of the background geometry from vacuum premetric γ to nonvacuum effective average metric q due to the presence of gravitational waves. This enables us to interpret equation (7) as the wave equation for perturbations h on the background $q = \gamma + \eta$. To make this wave equation applicable, one has to express η in terms of h . Using the Brill–Hartle averaging procedure [5] over a spacetime volume of size L for (7) (Efroimsky considers only space averaging but when the measurement lasts much longer than the period of waves one can employ a spacetime average) we obtain

$$R_{\mu\nu}^{(1)}(q, \eta) = -\langle R_{\mu\nu}^{(2)}(q, h) \rangle_L.
 \tag{8}$$

The averaging brackets on the left-hand side are omitted because the term contains only the modes with wavelength greater than L . It is thus clear from (8) and (5) that assumption 2 is natural since the left-hand side is linear in η whereas the right-hand side is quadratic in h .

Let us finally recall the derivation of the stress–energy tensor of gravitational waves. By analogy with the Ricci tensor expansion (4) the Einstein tensor of the vacuum premetric γ is represented as a series

$$0 = G_{\mu\nu}(\gamma) = G_{\mu\nu}(q) + G_{\mu\nu}^{(1)}(q, (-\eta)) + O(\eta^2),
 \tag{9}$$

and the effective stress–energy tensor of gravitational waves is defined as

$$G_{\mu\nu}(q) = 8\pi T_{\mu\nu}^{(gw)} \equiv R_{\mu\nu}^{(1)}(q, \eta) - \frac{1}{2}q_{\mu\nu}q^{\alpha\beta}R_{\alpha\beta}^{(1)}(q, \eta).
 \tag{10}$$

From (8) it follows (considering the Brill–Hartle averaging) that this tensor fully agrees with that of Isaacson [3].

The main advantage of Efroimsky’s perturbation method above is the possibility of consistently treating all low-frequency gravitational waves, and of explicitly deriving an

effective stress–energy tensor (influencing the background) in this case. It can be extended to nonvacuum spacetimes with $T_{\mu\nu}$ of ideal fluid and/or with a possible cosmological constant Λ , see [1, 2]. However, there are some problems concerning high-frequency gravitational waves which will now be discussed.

3. Modification to include high-frequency waves

In this section we first explicitly demonstrate that one cannot consistently apply Efroimsky’s treatment on Isaacson’s high-frequency waves [3] because assumption 2 is not fulfilled in such a case. Then we will present a possible solution to this problem.

Let us start with the observation that it is the nonvacuum background curved by the presence of gravitational waves—not the vacuum premetric γ —which is the basis of Isaacson’s nonlinear approach. Therefore, the nonvacuum average metric q is considered as the background on which high-frequency gravitational waves h propagate.

We wish to use the Efroimsky formalism in the high-frequency regime such that the tensor field h contains high-frequency modes. We assume that they have short wavelengths λ , and a small amplitude $h = O(\varepsilon)$, where $\varepsilon = \lambda/S \ll 1$ is a small parameter because $\lambda \ll S$, S denoting a typical scale on which the background changes substantially.

Let us emphasize that we follow here the same definition of the symbol $O(\varepsilon^n)$ as in [3], namely $f = O(\varepsilon^n)$ if there exists a constant $C > 0$ such that $|f| < C\varepsilon^n$ as $\varepsilon \rightarrow 0$. The quantity f need not necessarily be proportional to ε^n , it *can be even smaller* than $C\varepsilon^n$ for $\varepsilon \rightarrow 0$. Therefore, the assumption $h = O(\varepsilon)$ does *not* automatically imply that $h \sim \varepsilon$. The spectrum of possible high-frequency waves is thus not *a priori* restricted, it is only required that their amplitudes fall to zero at least linearly with ε , i.e. $|h(\varepsilon)| < C\varepsilon$.

Since we can consider $S = O(1)$ it follows that $O(\varepsilon) = O(\lambda)$ and $\partial h \sim h/\lambda = O(1)$. In accordance with Isaacson’s approach (note that the decomposition now reads $g = q + h$, instead of the notation $g = \gamma + h$ used in [3]) we obtain the following orders of magnitude for the derivatives of the background q and the perturbation h :

$$\begin{aligned} q_{\mu\nu} &= O(1), & h_{\mu\nu} &= O(\varepsilon), \\ q_{\mu\nu,\alpha} &= O(1), & h_{\mu\nu,\alpha} &= O(1), \\ q_{\mu\nu,\alpha\beta} &= O(1), & h_{\mu\nu,\alpha\beta} &= O(\varepsilon^{-1}). \end{aligned} \quad (11)$$

This results in the orders of magnitude of the terms in the Ricci tensor expansion (4), (5) as $R_{\mu\nu}^{(0)} = O(1)$, $R_{\mu\nu}^{(1)} = O(\varepsilon^{-1})$, $R_{\mu\nu}^{(2)} = O(1)$, $R_{\mu\nu}^{(3)} = O(\varepsilon)$. (12)

To apply the Efroimsky approach in this case we must consider the decomposition $q = \gamma + \eta$, where γ is the vacuum premetric and η represents (in this case) a substantial shift of the background geometry due to the presence of high-frequency gravitational waves h . We also introduce the scale L , such that $\lambda \ll L \ll S$. This enables us simultaneously to consider an averaging procedure in accordance with the Isaacson approach, and also to introduce a meaningful cut-off scale L even if the wavelengths of high-frequency waves are not assumed to reach this value.

Of course, the geometry shift η does not contain high-frequency perturbations. Considering the wave equation (7) and using the Brill–Hartle averaging over a spacetime volume L to obtain equation (8) we get into a conflict with assumption 2 which prescribes $O(\eta) = O(h^2)$. Indeed, if $h = O(\varepsilon)$ there should be $\eta = O(\varepsilon^2)$. But the right-hand side of (8) is now of the order of $O(1)$, see (12), and the same magnitude should also have the left-hand side. Since η does not contain high-frequency waves, it is essential that $\eta = O(1)$. This is obviously in contradiction with both assumptions 1 and 2. In fact, it disables any consistent perturbation expansions in the powers of η .

Let us now suggest a modification of the Efroimsky formalism which will incorporate also the above case of a ‘substantial’ change of the background geometry due to the presence of high-frequency waves. Instead of the perturbation expansion (6) we consider a formal decomposition of the Ricci tensor of the premetric $\gamma = q - \eta$, namely

$$0 = R_{\mu\nu}(\gamma) = R_{\mu\nu}(q) + \Delta R_{\mu\nu}(q, (-\eta)), \tag{13}$$

by which the expression $\Delta R_{\mu\nu}$ is *defined*. Both terms on the right-hand side of (13) are of the same order $O(1)$. Moreover, the quantity $\Delta R_{\mu\nu}$ is conserved with respect to the background geometry q which is easily seen from equation (13) and the relation $(R_{\mu\nu}(q))^{; \nu} = 0$ (the differentiation relates to the background metric q).

The question concerning the gauge invariance of $\Delta R_{\mu\nu}$ with respect to generalized gauge transformations has recently been analysed in detail by Anderson [16] in connection with possible definitions of the wave equation and stress–energy tensor for gravitational waves. Let us consider an arbitrary coordinate transformation of the type

$$\bar{x}^\mu = x^\mu + \xi^\mu, \tag{14}$$

which does not change the functional form of the background geometry q , i.e. $\bar{q}(\bar{x}) = q(\bar{x})$ so that $\gamma(x) \rightarrow \bar{\gamma}(\bar{x}) = q(\bar{x}) - \bar{\eta}(\bar{x})$. Now, to prove the invariance of $\Delta R_{\mu\nu}$ we adopt (slightly modified) Anderson’s argumentation. Performing the above coordinate transformation (14) of the Ricci tensor decomposition (13) we obtain

$$\bar{R}_{\mu\nu}(q(\bar{x})) + \bar{\Delta R}_{\mu\nu}(q(\bar{x}), (-\bar{\eta}(\bar{x}))) = \bar{R}_{\mu\nu}(\bar{\gamma}(\bar{x})) = 0. \tag{15}$$

Here $\bar{R}_{\mu\nu}$ and $\bar{\Delta R}_{\mu\nu}$ are the same as $R_{\mu\nu}$ and $\Delta R_{\mu\nu}$, respectively, because definition (1) is maintained in any coordinate. Evaluating relation (15) at $\bar{x} = x$ we thus get $R_{\mu\nu}(q(x)) = -\Delta R_{\mu\nu}(q(x), (-\bar{\eta}(x)))$, and using (13) we obtain

$$\Delta R_{\mu\nu}(q(x), (-\eta(x))) = \Delta R_{\mu\nu}(q(x), (-\bar{\eta}(x))). \tag{16}$$

A generalized gauge transformation is defined in [16] as a transformation in which the quantity $\bar{\eta}(x)$ is substituted for $\eta(x)$ into the tensor expressions of interest. This incorporates, as a particular case, the well-known infinitesimal gauge transformation

$$\bar{\eta}_{\mu\nu}(x) = \eta_{\mu\nu}(x) + \xi_{\mu;\nu} + \xi_{\nu;\mu}, \tag{17}$$

where η, ξ and their derivatives are small. Obviously, equation (16) expresses a generalized gauge invariance of $\Delta R_{\mu\nu}$.

After introducing the above decomposition (13) and demonstrating its invariance we can now present modification and generalization of the Efroimsky formalism. Replacing the term $R_{\mu\nu}^{(1)}(q, \eta)$ by $-\Delta R_{\mu\nu}(q, (-\eta))$ in equations (7), (8), (10), and omitting the terms $O(\eta^2)$ we obtain relations

$$R_{\mu\nu}^{(1)}(q, h) + R_{\mu\nu}^{(2)}(q, h) - \Delta R_{\mu\nu}(q, (-\eta)) + R_{\mu\nu}^{(3)}(q, h) + O(h^4) = 0, \tag{18}$$

$$\Delta R_{\mu\nu}(q, (-\eta)) = \langle R_{\mu\nu}^{(2)}(q, h) \rangle_L, \tag{19}$$

$$G_{\mu\nu}(q) = 8\pi \tilde{T}_{\mu\nu}^{(gw)} \equiv -\Delta R_{\mu\nu}(q, (-\eta)) + \frac{1}{2} q_{\mu\nu} q^{\alpha\beta} \Delta R_{\alpha\beta}(q, (-\eta)). \tag{20}$$

In the case where the gravitational waves do not have high-frequency modes it is still possible to employ the expansion of $-\Delta R_{\mu\nu}(q, (-\eta))$ in powers of η and use its dominant term $R_{\mu\nu}^{(1)}(q, \eta)$ instead. Thus we recover Efroimsky’s previous results (cf (7), (8), (10)).

In general, however, expressing η in terms of h from equation (19) becomes an extremely difficult task because it is no longer a linear equation for η . To overcome this problem we

can use equation (19) and substitute for $\Delta R_{\mu\nu}$ into the remaining equations (18) and (20). We obtain the relations

$$R_{\mu\nu}^{(1)}(q, h) + R_{\mu\nu}^{(2)}(q, h) - \langle R_{\mu\nu}^{(2)}(q, h) \rangle_L + R_{\mu\nu}^{(3)}(q, h) + O(h^4) = 0, \quad (21)$$

$$-G_{\mu\nu}(q) = \langle R_{\mu\nu}^{(2)}(q, h) \rangle_L - \frac{1}{2}q_{\mu\nu}q^{\alpha\beta} \langle R_{\alpha\beta}^{(2)}(q, h) \rangle_L \equiv -8\pi T_{\mu\nu}^{BH}. \quad (22)$$

Equation (22) is obviously in perfect agreement with the Isaacson result [3] which represents the background response to the presence of high-frequency gravitational waves, using the Brill–Hartle averaging to introduce the effective stress–energy tensor $T_{\mu\nu}^{BH}$ for high-frequency gravitational waves. Equation (21) is the wave equation for perturbations h on the average metric q . In the highest order of high-frequency approximation this clearly reduces to $R_{\mu\nu}^{(1)} = 0$ which also fully reproduces Isaacson’s result. Additional terms in (21) can be used for study of nonlinear effects on the wave propagation.

Note finally another interesting consequence of equation (19) and the gauge invariance (16) of $\Delta R_{\mu\nu}$. This directly guarantees gauge invariance of the stress–energy tensor $T_{\mu\nu}^{BH}$ defined in (22) (in the highest order). Proof of this property was presented already in the classic work [3], using, however, a much more complicated method.

4. Concluding remarks

In our contribution we have compared the Efroimsky [1, 2] and the Isaacson [3] self-consistent perturbation schemes which describe propagation of weak gravitational waves on a cosmological background. In both these approaches the background is influenced by the waves, i.e. the nonlinear effects are taken into account. The classical Isaacson method applies to high-frequency waves. On the other hand, the Efroimsky formalism is applicable to low-frequency gravitational waves but does not admit the high-frequency limit. We have suggested a modification of the Efroimsky formalism by employing the gauge-invariant decomposition (13) of the Ricci tensor, introduced recently by Anderson [16]. The resulting generalized system of equations (18)–(20) fully recovers the Efroimsky results in the absence of high-frequency modes, in the high-frequency limit it reproduces Isaacson’s formulae.

Although we have considered here for simplicity only vacuum metrics $\gamma_{\mu\nu}$ and $g_{\mu\nu}$, possible generalization to nonvacuum spacetimes is straightforward. In fact, Efroimsky has already generalized his formalism to spacetimes with ideal-fluid-like matter and a cosmological term [1, 2]; in the case of the Isaacson high-frequency approach, this was done recently in [15].

Acknowledgments

The work was supported in part by the grants GAČR 202/02/0735 and GAUK 166/2003 of the Czech Republic and the Charles University in Prague.

References

- [1] Efroimsky M 1992 Gravity waves in vacuum and in media *Class. Quantum Grav.* **9** 2601
- [2] Efroimsky M 1994 Weak gravitation waves in vacuum and in media: taking nonlinearity into account *Phys. Rev. D* **49** 6512
- [3] Isaacson R A 1968 Gravitational radiation in the limit of high frequency: I, II *Phys. Rev.* **166** 1263
- [4] Wheeler J A 1962 *Geometrodynamics* (New York: Academic)

- [5] Brill D R and Hartle J B 1964 Method of the self-consistent field in general relativity and its application to the gravitational geon *Phys. Rev. B* **135** 271
- [6] Choquet-Bruhat Y 1968 Construction de solutions radiatives approchées des équations d'Einstein *Commun. Math. Phys.* **12** 16
- [7] MacCallum M A H and Taub A H 1973 The averaged Lagrangian and high-frequency gravitational waves *Commun. Math. Phys.* **30** 153
- [8] Araujo M E 1986 On the assumptions made in treating the gravitational wave problem by the high-frequency approximation *Gen. Rel. Grav.* **18** 219
- [9] Araujo M E 1989 Lagrangian methods and nonlinear high-frequency gravitational waves *Gen. Rel. Grav.* **21** 323
- [10] Elster T 1981 Propagation of high-frequency gravitational waves in vacuum: nonlinear effects *Gen. Rel. Grav.* **13** 731
- [11] Burnett G A 1989 The high-frequency limit in general relativity *J. Math. Phys.* **30** 90
- [12] Taub A H 1976 High frequency gravitational radiation in Kerr–Schild space-times *Commun. Math. Phys.* **47** 185
- [13] Taub A H 1980 High-frequency gravitational waves, two-timing, and averaged Lagrangians *General Relativity and Gravitation* vol 1 ed A Held (New York: Plenum) p 539
- [14] Hogan P A and Futamase T 1993 Some high-frequency spherical gravity waves *J. Math. Phys.* **34** 154
- [15] Podolský J and Svítek O 2004 Some high-frequency gravitational waves related to exact radiative spacetimes *Gen. Rel. Grav.* **36** 387
- [16] Anderson P R 1997 Gauge-invariant effective stress–energy tensors for gravitational waves *Phys. Rev. D* **55** 3440