

Algebraic structure of Robinson–Trautman and Kundt geometries in arbitrary dimension

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 Class. Quantum Grav. 32 015001

(<http://iopscience.iop.org/0264-9381/32/1/015001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 195.113.23.22

This content was downloaded on 04/12/2014 at 12:37

Please note that [terms and conditions apply](#).

Algebraic structure of Robinson–Trautman and Kundt geometries in arbitrary dimension

J Podolský and R Švarc

Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles University in Prague, V Holešovičkách 2, 180 00 Praha 8, Czech Republic

E-mail: podolsky@mbox.troja.mff.cuni.cz and robert.svarc@mff.cuni.cz

Received 19 June 2014, revised 16 September 2014

Accepted for publication 17 September 2014

Published 3 December 2014



CrossMark

Abstract

We investigate the Weyl tensor algebraic structure of a fully general family of D -dimensional geometries that admit a non-twisting and shear-free null vector field \mathbf{k} . From the coordinate components of the curvature tensor we explicitly derive all Weyl scalars of various boost weights. This enables us to give a complete algebraic classification of the metrics in the case when the optically privileged null direction \mathbf{k} is a (multiple) Weyl aligned null direction (WAND). No field equations are applied, so the results are valid not only in Einstein's gravity, including its extension to higher dimensions, but also in any metric gravitation theory that admits non-twisting and shear-free spacetimes. We prove that all such geometries are of type I(b), or more special, and we derive surprisingly simple necessary and sufficient conditions under which \mathbf{k} is a double, triple or quadruple WAND. All possible algebraically special types, including the refinement to subtypes, are thus identified, namely II(a), II(b), II(c), II(d), III(a), III(b), N, O, II_i, III_i, D, D(a), D(b), D(c), D(d), and their combinations. Some conditions are identically satisfied in four dimensions. We discuss both important subclasses, namely the Kundt family of geometries with the vanishing expansion ($\Theta = 0$) and the Robinson–Trautman family ($\Theta \neq 0$, and in particular $\Theta = 1/r$). Finally, we apply Einstein's field equations and obtain a classification of all Robinson–Trautman vacuum spacetimes. This reveals fundamental algebraic differences in the $D > 4$ and $D = 4$ cases, namely that in higher dimensions there only exist such spacetimes of types $D(a) \equiv D(abd)$, $D(c) \equiv D(bcd)$ and O.

Keywords: algebraic classification, Kundt, Robinson–Trautman

PACS numbers: 04.20.Jb, 04.50.-h, 04.30.-w

1. Introduction

Exact spacetimes play a crucial role in understanding Einstein's general relativity and other metric theories of gravity. They enable us to investigate many mathematical and physical aspects of fundamental models in cosmology, black hole physics and theory of gravitational waves.

Among the most important of such classes of exact solutions there are Robinson–Trautman [1, 2] and Kundt [3, 4] geometries. They were discovered almost simultaneously about half a century ago, shortly after the advent of new concepts and techniques in general relativity, in particular geometrical optics of null congruences and algebraic classification of the Weyl tensor. Since then, an enormous progress has been made in investigation of their various properties.

From the geometrical point of view, both these classes belong to a large family of geometries admitting a non-twisting shear-free congruence of geodesics, generated by a null vector field \mathbf{k} . The Kundt class is defined by having vanishing expansion while the other case with non-vanishing expansion defines the Robinson–Trautman class. The former includes the famous pp-waves (with a covariantly constant \mathbf{k}), more general non-expanding gravitational waves (including gyratons, non-vanishing cosmological constant Λ , impulsive limits), VSI and CSI spacetimes (for which all scalar invariants of curvature vanish and are constant, respectively), or the direct-product spacetimes (Bertotti–Robinson, Nariai, Plebański–Hacyan). In the Robinson–Trautman class there are, e.g., some well-known black holes (Schwarzschild, Reissner–Nordström, Schwarzschild–de Sitter, Vaidya), expanding spherical gravitational waves (including Λ), the C-metric (representing the field of accelerated black holes) or Kinnersley's and Bonnor's 'photon rockets'. Details and a number of references can be found in the monographs [5, 6] (chapters 28, 31 and 19, 18, respectively).

In view of the growing interest to generalize Einstein's theory and to extend it to higher dimensions, it is a natural task to find and analyze specific properties of such spacetimes. Assuming the validity of Einstein's field equations (for vacuum with Λ , aligned electromagnetic field, pure radiation, gyratonic matter), the explicit Robinson–Trautman class in any dimension was studied in [7–9]. The complementary Kundt class was also investigated, e.g., in [10, 11]. The results were summarized in the recent review [12] on algebraic properties of higher dimensional spacetimes.

In this paper we consider the fully general class of non-twisting and shear-free geometries in an arbitrary dimension $D \geq 4$, without assuming any field equations. Starting from the canonical form of the metric we derive all components of the Riemann, Ricci and Weyl tensors, and the Weyl tensor is projected onto a suitable null frame. It enables us to give a complete and explicit classification of the whole class into the algebraic types and subtypes based on the WAND multiplicity of the optically privileged null vector field \mathbf{k} . Our new results thus considerably generalize the study of algebraic structure of the non-expanding Kundt family of geometries [13] and exemplify general conclusions of previous works [14–17].

We introduce the general metric in section 2. In section 3 the null frame components of the Weyl tensor are employed for the algebraic classification. The necessary and sufficient conditions for all principal and secondary alignment (sub)types are discussed in sections 4 and 5. Results for the Kundt class are summarized in section 6 while those for the Robinson–Trautman class are contained in section 7. In section 8 we discuss a special case, namely the Robinson–Trautman vacuum spacetimes in D -dim Einstein's theory. Coordinate components of the Riemann, Ricci, and Weyl tensors for the generic non-twisting shear-free geometry are given in appendix A.

2. Robinson–Trautman and Kundt geometries

The metric of the most general non-twisting D -dimensional geometry can be written in the form [7]

$$ds^2 = g_{pq}(r, u, x)dx^p dx^q + 2g_{up}(r, u, x)du dx^p - 2du dr + g_{uu}(r, u, x)du^2, \quad (1)$$

if natural coordinates are used. A non-twisting character of the spacetime implies the existence of a foliation by null hypersurfaces $u = \text{const.}$, i.e., a family of maximal integral submanifolds labeled by the coordinate u . By the Frobenius theorem, this is equivalent to the existence of a non-twisting null vector field \mathbf{k} that is everywhere tangent (and normal) to $u = \text{const.}$ Since this field \mathbf{k} generates a congruence of null geodesics in the whole spacetime, it is most natural to take their affine parameter r as the second coordinate, so that $\mathbf{k} = \partial_r$. At any fixed u and r we are thus left with a $(D - 2)$ -dimensional Riemannian manifold covered by the spatial coordinates x^p . We will use the indices m, n, p, q (ranging from 2 to $D - 1$) to label these spatial coordinates on the transverse space, and a shorthand x for their complete set. The non-vanishing contravariant metric components are

$$g^{pq}, \quad g^{ru} = -1, \quad g^{rp} = g^{pq}g_{uq}, \quad g^{rr} = -g_{uu} + g^{pq}g_{up}g_{uq}, \quad (2)$$

where g^{pq} is an inverse matrix to g_{pq} . This implies

$$g_{up} = g_{pq}g^{rq}, \quad g_{uu} = -g^{rr} + g_{pq}g^{rp}g^{rq}. \quad (3)$$

The covariant derivative of the geometrically privileged null vector field $\mathbf{k} = \partial_r$ with respect to the metric (1) is $k_{a;b} = \Gamma_{ab}^u = \frac{1}{2}g_{ab,r}$ so that $k_{r;b} = 0 = k_{a;r}$. Consequently, the *optical matrix* [12] defined as $\rho_{ij} \equiv k_{a;b} m_i^a m_j^b$, where m_i^a are components of $(D - 2)$ unit vectors \mathbf{m}_i such that $\mathbf{m}_i \cdot \mathbf{k} = 0 \Rightarrow m_i^u = 0$, forming an orthonormal basis in the transverse Riemannian space, is simply given by $\rho_{ij} = k_{p;q} m_i^p m_j^q = \frac{1}{2}g_{pq,r} m_i^p m_j^q$. This can be decomposed as $\rho_{ij} = A_{ij} + \sigma_{ij} + \Theta\delta_{ij}$, where $A_{ij} \equiv \rho_{[ij]}$ is the antisymmetric *twist matrix*, σ_{ij} is the symmetric traceless *shear matrix*, and the scalar $\Theta \equiv \frac{1}{D-2} \delta^{ij} \rho_{ij}$ determines the *expansion* of the privileged vector field \mathbf{k} .

It can be observed that $A_{ij} = \frac{1}{2}g_{pq,r} m_i^p m_j^q = 0$ which confirms that the metric (1) is *non-twisting*. Now, imposing the additional condition that the metric is *shear-free*, $\sigma_{ij} = 0$, we obtain the relation

$$\rho_{ij} = \Theta\delta_{ij} = \frac{1}{2}g_{pq,r} m_i^p m_j^q. \quad (4)$$

Using the orthonormality relation $\delta_{ij} = g_{pq} m_i^p m_j^q$ we thus immediately infer

$$g_{pq,r} = 2\Theta g_{pq}, \quad (5)$$

implying $g_{pq,rr} = 2(\Theta_{,r} + 2\Theta^2)g_{pq}$. The expression (5) can be integrated as

$$g_{pq} = R^2(r, u, x)h_{pq}(u, x), \quad \text{where} \quad \frac{R_{,r}}{R} = \Theta. \quad (6)$$

Since either $\Theta = 0$ or $\Theta \neq 0$, there are thus two distinct classes of non-twisting shear-free geometries. The *Kundt class* [3–6, 10, 12, 13, 18] is defined by having the vanishing expansion, $\Theta = 0$, in which case the spatial metric $g_{pq}(u, x) = h_{pq}(u, x)$ is independent of the affine parameter r (and R in (6) effectively reduces to $R = 1$). The other case $\Theta \neq 0$ gives the expanding *Robinson–Trautman class* [1, 2, 5–8, 12], for which R is a non-trivial function of r determined by $R = \exp\left(\int \Theta(r, u, x) dr\right)$.

3. Frame components of the Weyl tensor and its classification

The natural null frame for the general metric (1) is given by

$$\mathbf{k} = \mathbf{k} = \partial_r, \quad \mathbf{l} = \frac{1}{2}g_{uu}\partial_r + \partial_u, \quad \mathbf{m}_i = m_i^p \left(g_{up}\partial_r + \partial_p \right), \quad (7)$$

satisfying $\mathbf{k} \cdot \mathbf{l} = -1$, $\mathbf{m}_i \cdot \mathbf{m}_j = \delta_{ij}$. All the Weyl tensor components with respect to such a frame $(\mathbf{k}, \mathbf{l}, \mathbf{m}_i)$, sorted by the boost weight, can be denoted as [13, 19]

$$\begin{aligned} \Psi_{0ij} &= C_{abcd} k^a m_i^b k^c m_j^d, \\ \Psi_{1jk} &= C_{abcd} k^a m_i^b m_j^c m_k^d, \quad \Psi_{1T^i} = C_{abcd} k^a l^b k^c m_i^d, \\ \Psi_{2^{ijkl}} &= C_{abcd} m_i^a m_j^b m_k^c m_l^d, \quad \Psi_{2S} = C_{abcd} k^a l^b l^c k^d, \\ \Psi_{2^{ij}} &= C_{abcd} k^a l^b m_i^c m_j^d, \quad \Psi_{2T^{ij}} = C_{abcd} k^a m_i^b l^c m_j^d, \\ \Psi_{3^{ijk}} &= C_{abcd} l^a m_i^b m_j^c m_k^d, \quad \Psi_{3T^i} = C_{abcd} l^a k^b l^c m_i^d, \\ \Psi_{4^{ij}} &= C_{abcd} l^a m_i^b l^c m_j^d. \end{aligned} \quad (8)$$

The scalars in the right column could, in fact, be obtained from those in the left column by contractions, namely $\Psi_{1T^i} = \Psi_{1^{k_i}}$, $\Psi_{2S} = \Psi_{2T^{kk}}$, $\Psi_{2T^{(ij)}} = \frac{1}{2}\Psi_{2^{ijk}}$, $\Psi_{2T^{[ij]}} = \frac{1}{2}\Psi_{2^{ij}}$, $\Psi_{3T^i} = \Psi_{3^{k_i}}$. Relations of these Newman–Penrose-like quantities to other equivalent notations employed in [12] and elsewhere can be found in [13].

For the invariant (sub)classification of the Weyl tensor algebraic structure it is also necessary to introduce the following irreducible components of these scalars (see [12]):

$$\tilde{\Psi}_{1^{ijk}} \equiv \Psi_{1^{ijk}} - \frac{1}{D-3} \left(\delta_{ij}\Psi_{1T^k} - \delta_{ik}\Psi_{1T^j} \right), \quad (9)$$

$$\tilde{\Psi}_{2T^{(ij)}} \equiv \Psi_{2T^{(ij)}} - \frac{1}{D-2} \delta_{ij}\Psi_{2S}, \quad (10)$$

$$\begin{aligned} \tilde{\Psi}_{2^{ijkl}} &\equiv \Psi_{2^{ijkl}} - \frac{2}{D-4} \left(\delta_{ik}\tilde{\Psi}_{2T^{(jl)}} + \delta_{jl}\tilde{\Psi}_{2T^{(ik)}} - \delta_{il}\tilde{\Psi}_{2T^{(jk)}} - \delta_{jk}\tilde{\Psi}_{2T^{(il)}} \right) \\ &\quad - \frac{2}{(D-2)(D-3)} \left(\delta_{ik}\delta_{jl} - \delta_{il}\delta_{jk} \right) \Psi_{2S}, \end{aligned} \quad (11)$$

$$\tilde{\Psi}_{3^{ijk}} \equiv \Psi_{3^{ijk}} - \frac{1}{D-3} \left(\delta_{ij}\Psi_{3T^k} - \delta_{ik}\Psi_{3T^j} \right). \quad (12)$$

The main step now is to project the coordinate components (A.34)–(A.43) of the Weyl tensor of a generic non-twisting shear-free geometry (see appendix A) onto the null frame (7). A long calculation with non-trivial cancelations of various terms reveals that the corresponding Weyl scalars take the following explicit and surprisingly simple form

$$\Psi_{0ij} = 0, \quad (13)$$

$$\Psi_{1T^i} = m_i^p \frac{D-3}{D-2} \left[\left(-\frac{1}{2}g_{up,r} + \Theta g_{up} \right)_{,r} + \Theta_{,p} \right], \quad (14)$$

$$\tilde{\Psi}_{1^{ijk}} = 0, \quad (15)$$

$$\Psi_{2S} = \frac{D-3}{D-1} P, \quad (16)$$

$$\tilde{\Psi}_{2T^{(ij)}} = m_i^p m_j^q \frac{1}{D-2} \left(Q_{pq} - \frac{1}{D-2} g_{pq} Q \right), \quad (17)$$

$$\tilde{\Psi}_{2^{ijkl}} = m_i^m m_j^p m_k^n m_l^q {}^S C_{mpnq}, \quad (18)$$

$$\Psi_{2^{ij}} = m_i^p m_j^q F_{pq}, \quad (19)$$

$$\Psi_{3T^i} = m_i^p \frac{D-3}{D-2} V_p, \quad (20)$$

$$\tilde{\Psi}_{3^{ijk}} = m_i^p m_j^m m_k^q \left(X_{pmq} - \frac{2}{D-3} g_{p[m} X_{q]1} \right), \quad (21)$$

$$\Psi_{4^{ij}} = m_i^p m_j^q \left(W_{pq} - \frac{1}{D-2} g_{pq} W \right), \quad (22)$$

where

$$\begin{aligned} P = & \left(\frac{1}{2} g_{uu,r} - \Theta g_{uu} \right)_r + \frac{1}{(D-2)(D-3)} {}^S R - \frac{1}{4} \frac{D-4}{D-2} g^{mn} g_{um,r} g_{un,r} \\ & + \frac{1}{D-2} \left(g^{rn} g_{un,rr} + g^{mn} g_{um,r||n} \right) - \frac{2}{D-2} g^{rn} g_{un} \Theta_{,r} - 2\Theta_{,u} - \frac{4}{D-2} g^{rn} \Theta_{,n} \\ & - \Theta^2 \frac{D-4}{D-2} g^{rn} g_{un} + \Theta \left(\frac{D-6}{D-2} g^{rn} g_{un,r} - \frac{2}{D-2} g^{mn} g_{um||n} \right), \end{aligned} \quad (23)$$

$$\begin{aligned} Q_{pq} = & {}^S R_{pq} + (D-4) \left[\frac{1}{2} \left(f_{pq} + g_{u(p} g_{q)u,rr} \right) - \left(\Theta_{,r} - \Theta^2 \right) g_{up} g_{uq} \right. \\ & \left. - 2g_{u(p} \Theta_{,q)} - \Theta \left(g_{u(p||q)} + 2g_{u(p} g_{q)u,r} \right) \right], \end{aligned} \quad (24)$$

$$F_{pq} = g_{u[p,q],r} - g_{u[p} g_{q]u,rr} + 2\Theta \left(g_{u[p} g_{q]u,r} - g_{u[p,q]} \right), \quad (25)$$

$$\begin{aligned} V_p = & \frac{1}{2} \left[\frac{1}{2} g_{uu} g_{up,rr} - g_{uu,rp} + g_{up,ru} - \frac{1}{2} g^{rn} g_{un,r} g_{up,r} \right. \\ & \left. + g^{mn} g_{um,r} E_{np} - g_{up} \left(g_{uu,rr} - \frac{1}{2} g^{mn} g_{um,r} g_{un,r} \right) \right] \\ & + \frac{1}{D-3} \left[\frac{1}{2} g^{rn} g_{un} g_{up,rr} + g^{mn} e_{m[n} g_{p]u,r} - g^{rn} g_{u[n,p],r} + \frac{1}{2} g^{rn} \left(g_{u[p,r||n]} + f_{pn} \right) \right. \\ & \left. - g^{mn} \left(g_{m[p,u||n]} + g_{u[m,p]||n} \right) - \frac{1}{2} g_{up} \left(g^{rn} g_{un,rr} + g^{mn} f_{mn} \right) \right] \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2}g_{up}g_{uu}\Theta_{,r} + g_{up}\Theta_{,u} + \frac{1}{2}g_{uu}\Theta_{,p} \\
& - \Theta \left[\frac{1}{2}g_{uu}g_{up,r} - g_{uu,p} + g_{up,u} - g^m g_{u[n}g_{p]u,r} + g^m E_{np} - g_{up}g_{uu,r} \right. \\
& \left. + \frac{1}{D-3} \left(3g^m g_{u[n}g_{p]u,r} - 3g^m g_{u[n,p]} - \frac{1}{2}g_{up}g^{mn}g_{mn,u} + \frac{1}{2}g^m g_{np,u} \right) \right], \quad (26)
\end{aligned}$$

$$\begin{aligned}
X_{pmq} & = g_{p[m,u]q} + g_{u[q,m]p} + g_{up}g_{u[m}g_{q]u,rr} + e_{p[m}g_{q]u,r} \\
& - g_{u[q}g_{m]u,r}p - g_{up}g_{u[m,r]q} - \frac{1}{2}g_{u[q}g_{m]u,r}g_{up,r} \\
& + \Theta \left(3g_{u[q}g_{m]u,r}g_{up} + g_{u[q}g_{m]p,u} + g_{u[q}g_{m]u}p - g_{up}g_{[m}g_{q]u} - 2g_{u[q,m]}g_{up} \right), \quad (27)
\end{aligned}$$

$$\begin{aligned}
W_{pq} & = -\frac{1}{2}g_{uu}g_{p}q - \frac{1}{2}g_{pq,uu} + g_{u(p,u)q} - \frac{1}{2}g_{uu,r}e_{pq} + \frac{1}{2}g_{uu,(p}g_{q)u,r} - g_{uu,r(p}g_{q)u} \\
& + \frac{1}{2}g_{uu}g_{u(p,r)q} + \frac{1}{2}g_{uu}g_{u(q}g_{p)u,rr} - \frac{1}{2}g_{uu,rr}g_{up}g_{uq} + g_{u(q}g_{p)u,ru} \\
& + \frac{1}{4}g^{mn} \left(g_{um}g_{un}g_{up,r}g_{uq,r} + g_{um,r}g_{un,r}g_{up}g_{uq} \right) - \frac{1}{2}g^{mn}g_{um}g_{un,r}g_{u(q}g_{p)u,r} \\
& + g^{mn} \left(E_{mp}E_{nq} + g_{um,r}E_{n(p}g_{q)u} - g_{um}E_{n(p}g_{q)u,r} \right) \\
& + \Theta \left(g_{up}g_{uq}g_{uu,r} + g_{uu,(p}g_{q)u} - g_{uu}g_{u(p}g_{q)u,r} - 2g_{u(p}g_{q)u,u} - \frac{1}{2}g_{uu}g_{pq,u} \right), \quad (28)
\end{aligned}$$

their contractions are defined as $Q \equiv g^{pq}Q_{pq}$, $W \equiv g^{pq}W_{pq}$ and $X_q \equiv g^{pm}X_{pmq}$, and the auxiliary quantities are defined in (A.44)–(A.53).

As a generalization of the classical Petrov classification of four-dimensional spacetimes in Einstein's theory, classification scheme of the algebraic structure of the Weyl tensor is based on whether the scalars (8) of various boost weights vanish or not in a *suitable* null frame [20], see [12] for a recent comprehensive review.

Specifically, it is possible to introduce the *principal alignment types and subtypes* of the Weyl tensor in any dimension D based on the existence of the (multiple) *Weyl aligned null direction* (WAND) k , as summarized in table 1. Apart from a fully generic type G with all Weyl tensor components non-vanishing, there is type I with subtypes I(a) and I(b), type II with four possible subtypes II(a), II(b), II(c) and II(d), type III with two subtypes III(a) and III(b), type N and type O corresponding to the case when the Weyl tensor vanishes completely.

Subsequently, it is possible to introduce *secondary alignment types* of the Weyl tensor defined by the property that there exists an *additional* WAND l , namely I_i , II_i and III_i . These are the types I, II, and III, respectively, for which not only $\Psi_{0\dot{j}} = 0$ but also $\Psi_{4\dot{j}} = 0$. There is also the 'degenerate' case of type D equivalent to II_i for which only the zero-boost weight Weyl scalars $\Psi_{2..}$ are non-vanishing (see table 2).

Of course, various combinations of these possibilities can occur. For example, there may be a spacetime with the algebraic structure II(ab) which means that it is *both* of subtype II(a) and II(b). Clearly, $II = I(ab)$, $III = II(abcd)$, $N = III(ab)$. In addition, there may be, for example, a $II(c)_i$ spacetime which means that it is simultaneously of subtype II(c) and II_i . Or there can

Table 1. The principal alignment types and subtypes of the Weyl tensor defined by the existence of a (multiple) WAND k .

Type	Vanishing components									
G	No null frame exists in which all components $\Psi_{0^{\dot{a}b}}$ vanish									
I	$\Psi_{0^{\dot{a}b}}$									
I(a)	$\Psi_{0^{\dot{a}b}}$	Ψ_{1T^i}								
I(b)	$\Psi_{0^{\dot{a}b}}$	$\tilde{\Psi}_{\dot{p}^{\dot{q}k}}$								
II	$\Psi_{0^{\dot{a}b}}$	Ψ_{1T^i}	$\tilde{\Psi}_{\dot{p}^{\dot{q}k}}$							
II(a)	$\Psi_{0^{\dot{a}b}}$	Ψ_{1T^i}	$\tilde{\Psi}_{\dot{p}^{\dot{q}k}}$	Ψ_{2S}						
II(b)	$\Psi_{0^{\dot{a}b}}$	Ψ_{1T^i}	$\tilde{\Psi}_{\dot{p}^{\dot{q}k}}$	$\tilde{\Psi}_{2T^{(ij)}}$						
II(c)	$\Psi_{0^{\dot{a}b}}$	Ψ_{1T^i}	$\tilde{\Psi}_{\dot{p}^{\dot{q}k}}$	$\tilde{\Psi}_{2^{\dot{p}k\dot{l}}}$						
II(d)	$\Psi_{0^{\dot{a}b}}$	Ψ_{1T^i}	$\tilde{\Psi}_{\dot{p}^{\dot{q}k}}$	$\Psi_{2^{\dot{p}j}}$						
III	$\Psi_{0^{\dot{a}b}}$	Ψ_{1T^i}	$\tilde{\Psi}_{\dot{p}^{\dot{q}k}}$	Ψ_{2S}	$\tilde{\Psi}_{2T^{(ij)}}$	$\tilde{\Psi}_{2^{\dot{p}k\dot{l}}}$	$\Psi_{2^{\dot{p}j}}$			
III(a)	$\Psi_{0^{\dot{a}b}}$	Ψ_{1T^i}	$\tilde{\Psi}_{\dot{p}^{\dot{q}k}}$	Ψ_{2S}	$\tilde{\Psi}_{2T^{(ij)}}$	$\tilde{\Psi}_{2^{\dot{p}k\dot{l}}}$	$\Psi_{2^{\dot{p}j}}$	Ψ_{3T^i}		
III(b)	$\Psi_{0^{\dot{a}b}}$	Ψ_{1T^i}	$\tilde{\Psi}_{\dot{p}^{\dot{q}k}}$	Ψ_{2S}	$\tilde{\Psi}_{2T^{(ij)}}$	$\tilde{\Psi}_{2^{\dot{p}k\dot{l}}}$	$\Psi_{2^{\dot{p}j}}$	$\tilde{\Psi}_{3^{\dot{p}k}}$		
N	$\Psi_{0^{\dot{a}b}}$	Ψ_{1T^i}	$\tilde{\Psi}_{\dot{p}^{\dot{q}k}}$	Ψ_{2S}	$\tilde{\Psi}_{2T^{(ij)}}$	$\tilde{\Psi}_{2^{\dot{p}k\dot{l}}}$	$\Psi_{2^{\dot{p}j}}$	Ψ_{3T^i}	$\tilde{\Psi}_{3^{\dot{p}k}}$	
O	$\Psi_{0^{\dot{a}b}}$	Ψ_{1T^i}	$\tilde{\Psi}_{\dot{p}^{\dot{q}k}}$	Ψ_{2S}	$\tilde{\Psi}_{2T^{(ij)}}$	$\tilde{\Psi}_{2^{\dot{p}k\dot{l}}}$	$\Psi_{2^{\dot{p}j}}$	Ψ_{3T^i}	$\tilde{\Psi}_{3^{\dot{p}k}}$	$\Psi_{4^{\dot{p}j}}$

Table 2. The secondary alignment types of the Weyl tensor defined by the existence of another WAND l (which is a double WAND in the case of type D).

Type	Principal type	Additional vanishing components
I_i	I	$\Psi_{4^{\dot{a}b}}$
II_i	II	$\Psi_{4^{\dot{a}b}}$
III_i	III	$\Psi_{4^{\dot{a}b}}$
$D \equiv II_{ii}$	II	Ψ_{3T^i} $\tilde{\Psi}_{3^{\dot{p}k}}$ $\Psi_{4^{\dot{p}j}}$

be a D(bcd) spacetime defined by the property that there exists a double WAND k and a double WAND l such that the only non-vanishing Weyl scalar is Ψ_{2S} .

In our present contribution, we are going to apply this algebraic classification scheme to the fully general family of non-twisting and shear-free geometries. This contains both the non-expanding Kundt class ($\Theta = 0$) and the expanding Robinson–Trautman class ($\Theta \neq 0$). In particular, we will completely characterize all possible principal alignment types of the Weyl tensor with respect to the optically privileged null vector field $k = \partial_r$.

Using the fact that $\Psi_{0^{\dot{a}b}} = 0$, see (13), and table 1, it immediately follows that a *generic Kundt or Robinson–Trautman geometry is of algebraic type I* (or more special), so that the optically privileged (non-twisting and shear-free) null vector field $k = \partial_r$ is a WAND. Moreover, since the relation (15) reads $\tilde{\Psi}_{\dot{p}^{\dot{q}k}} = 0$, *any Kundt or Robinson–Trautman geometry is, in fact, of algebraic subtype I(b)*, or more special.

We will now discuss all possibilities when $k = k$ is a *multiple WAND*. In other words, the spacetime geometry is (at least) of algebraic type II with respect to ∂_r .

4. Multiple WAND \mathbf{k} and algebraically special (sub)types

When the vector field $\mathbf{k} = \partial_r$ is (at least) a *double* WAND then the spacetime is *algebraically special* (of type II, or more special)¹. It follows from table 1 that such a situation occurs if, and only if,

$$\Psi_{1T^i} = 0, \quad (29)$$

where Ψ_{1T^i} is given by (14). Since the spatial vectors \mathbf{m}_i are linearly independent, this condition is equivalent to $(g_{up,r} - 2\Theta g_{up})_{,r} = 2\Theta_{,p}$ which can be rewritten as

$$g_{up,rr} = 2g_{up}\Theta_{,r} + 2\Theta g_{up,r} + 2\Theta_{,p}, \quad (30)$$

or integrated to

$$g_{up,r} = 2\Theta g_{up} + f_p, \quad (31)$$

where $f_p(r, u, x) \equiv 2 \int \Theta_{,p} dr + \varphi_p(u, x)$, that is

$$f_{p,r} = 2\Theta_{,p}. \quad (32)$$

Applying the condition $\Psi_{1T^i} = 0$ which implies (31) and (30), the functions (23)–(28) determining the remaining Weyl scalars (16)–(22) simplify considerably to

$$\begin{aligned} P &= \left(\frac{1}{2} g_{uu,r} - \Theta g_{uu} \right)_{,r} + \frac{1}{(D-2)(D-3)} {}^S R \\ &\quad + \frac{1}{D-2} g^{mn} f_{m||n} - \frac{1}{4} \frac{D-4}{D-2} g^{mn} f_m f_n - 2\Theta_{,u}, \end{aligned} \quad (33)$$

$$Q_{pq} = {}^S R_{pq} + \frac{1}{2}(D-4) \left(f_{(p||q)} + \frac{1}{2} f_p f_q \right), \quad (34)$$

$$F_{pq} = f_{[p,q]}, \quad (35)$$

$$\begin{aligned} V_p &= \frac{1}{2} (f_{p,u} - g_{uu,rp} - g_{up} g_{uu,rr} + g^{mn} f_m E_{np}) + \frac{1}{2} \frac{D-4}{D-3} g^{mn} g_{u[p} f_m] f_n \\ &\quad - \frac{1}{D-3} g^{mn} \left[g_{m[p,u]||n} + g_{u[m,p]||n} + e_{m[p} f_n] + \frac{1}{2} g_{up} f_m] f_n \right. \\ &\quad \left. - \frac{1}{2} g_{um} (f_{(p||p)} - 3f_{[n,p]}) \right] \\ &\quad + g_{up} g_{uu} \Theta_{,r} + 2g_{up} \Theta_{,u} + g_{uu} \Theta_{,p} + \Theta (g_{up} g_{uu,r} + g_{uu,p}), \end{aligned} \quad (36)$$

$$X_{pmq} = g_{p[m,u]||q} + g_{u[q,m]||p} + e_{p[m} f_q] - g_{u[q} f_m]||p - g_{up} f_{[m,q]} - \frac{1}{2} g_{u[q} f_m] f_p, \quad (37)$$

¹ In principle, there could exist ‘peculiar’ algebraically special spacetimes for which \mathbf{k} is *not* a *double* WAND and there is *another* double WAND vector field.

Table 3. Principal alignment types and subtypes of the algebraically special Weyl tensor with respect to the multiple WAND $\mathbf{k} = \partial_r$.

Type	Necessary and sufficient condition	Equation
II(a)	$P = 0$	(40)
II(b)	$Q_{pq} - \frac{1}{D-2} g_{pq} Q = 0$	(41)
II(c)	${}^S C_{mpnq} = 0$	(42)
II(d)	$F_{pq} = 0$	(43)
III	II(abcd)	—
III(a)	$V_p = 0$	(47)
III(b)	$X_{pmq} - \frac{2}{D-3} g_{p[m} X_{q]} = 0$	(48)
N	III(ab)	—
O	$W_{pq} - \frac{1}{D-2} g_{pq} W = 0$	(50)

$$\begin{aligned}
W_{pq} = & -\frac{1}{2} g_{uu} \parallel_{p||q} - \frac{1}{2} g_{pq,uu} + g_{u(p,u||q)} - \frac{1}{2} g_{uu,r} e_{pq} - g_{uu,r(p} g_{q)u} - \frac{1}{2} g_{up} g_{uq} g_{uu,rr} \\
& + \frac{1}{2} g_{uu,(p} f_{q)} + \frac{1}{2} g_{uu} f_{(p||q)} + g_{u(p} f_{q),u} + \frac{1}{4} g^{mn} (g_{um} g_{un} f_{p} f_{q} + f_m f_n g_{up} g_{uq}) \\
& - \frac{1}{2} g^{mn} g_{um} f_n g_{u(p} f_{q)} + g^{mn} (E_{mp} E_{nq} + f_m E_n (p} g_{q)u} - g_{um} E_n (p} f_{q)}) \\
& + g_{uu} g_{up} g_{uq} \Theta_{,r} + 2g_{up} g_{uq} \Theta_{,u} + 2g_{uu} g_{u(p} \Theta_{,q)} \\
& + \Theta \left(2g_{uu,(p} g_{q)u} + g_{uu} g_{u(p||q)} + g_{up} g_{uq} g_{uu,r} - \frac{1}{2} g_{uu} g_{pq,u} \right). \tag{38}
\end{aligned}$$

Recall that

$$e_{pq} = g_{u(p||q)} - \frac{1}{2} g_{pq,u}, \quad E_{pq} = g_{u[p,q]} + \frac{1}{2} g_{pq,u}, \tag{39}$$

$f_{pq} = f_{(p||q)} + \frac{1}{2} f_{,p} f_{q} + 2g_{u(p} \Theta_{,q)} + 2\Theta^2 g_{up} g_{uq} + 2\Theta (g_{u(p||q)} + g_{u(p} f_{q)})$, and \parallel denotes the covariant derivative with respect to the spatial metric g_{pq} , while the corresponding Ricci tensor and Ricci scalar are ${}^S R_{pq}$ and ${}^S R$, respectively, see appendix A.

Using table 1 we can thus explicitly present the necessary and sufficient conditions determining all possible principal algebraic (sub)types of non-twisting shear-free geometries with a double, triple and quadruple WAND $\mathbf{k} = \partial_r$. These are summarized in table 3.

4.1. Type II subtypes with a double WAND \mathbf{k}

The Robinson–Trautman or Kundt spacetimes (1) with (5) satisfying the condition (31), i.e., $\Psi_{1T^i} = 0$ implying that they are (at least) of type II with respect to the null direction $\mathbf{k} = \partial_r$, admit the following particular algebraic subtypes of the Weyl tensor:

- Subtype II(a) $\Leftrightarrow \Psi_{2S} = 0 \Leftrightarrow P = 0 \Leftrightarrow$ the metric function g_{uu} satisfies the relation:

$$\left(\frac{1}{2}g_{uu,r} - \Theta g_{uu}\right)_{,r} = -\frac{1}{(D-2)(D-3)} sR - \frac{1}{D-2} g^{mn} f_{m||n} + \frac{1}{4} \frac{D-4}{D-2} g^{mn} f_m f_n + 2\Theta_{,u}. \quad (40)$$

This determines the specific dependence of $g_{uu}(r, u, x)$ on the coordinate r which is the affine parameter along the null congruence generated by \mathbf{k} .

- Subtype II(b) $\Leftrightarrow \check{\Psi}_{2T^{(ii)}} = 0 \Leftrightarrow Q_{pq} = \frac{1}{D-2} g_{pq} Q$:

$$sR_{pq} - \frac{g_{pq}}{D-2} sR = -\frac{1}{2}(D-4) \left[\left(f_{p||q} + \frac{1}{2} f_p f_q \right) - \frac{g_{pq}}{D-2} g^{mn} \left(f_{m||n} + \frac{1}{2} f_m f_n \right) \right]. \quad (41)$$

This is identically satisfied when $D = 4$ since for any two-dimensional Riemannian space there is $sR_{pq} = \frac{1}{2} g_{pq} sR$.

- Subtype II(c) $\Leftrightarrow \check{\Psi}_{2ijkl} = 0$:

$$sC_{mpnq} = 0. \quad (42)$$

This is always satisfied when $D = 4$ and $D = 5$ since the Weyl tensor vanishes identically in dimensions 2 and 3.

- Subtype II(d) $\Leftrightarrow \Psi_{2ij} = 0 \Leftrightarrow F_{pq} = 0$:

$$f_{[p,q]} = 0. \quad (43)$$

Introducing a 1-form $\phi \equiv f_p dx^p$ in the transverse $(D-2)$ -dim Riemannian space, this condition is equivalent to the condition that ϕ is closed ($d\phi = 0$). By the Poincaré lemma, on any contractible domain there exists a potential function \mathcal{F} such that $\phi = d\mathcal{F}$, that is $f_p = \mathcal{F}_{,p}$. In a general case, such \mathcal{F} exists only *locally*.

These four distinct subtypes of type II can be arbitrarily combined. Clearly, in the $D = 4$ case the algebraically special non-twisting shear-free geometries are always of subtype II(bc).

4.2. Type III subtypes with a triple WAND \mathbf{k}

The Robinson–Trautman or Kundt spacetime is of algebraic type III with respect to the triple WAND $\mathbf{k} = \partial_r$ if *all four* independent conditions (40)–(43) are satisfied *simultaneously*. In such a case the zero-boost-weight Weyl tensor components $\Psi_{2..}$ vanish and we obtain geometries of type II(abcd) \equiv III, with the remaining Weyl scalars (20)–(22) determined by the structural functions (36)–(38) now simplified to

$$V_p = \frac{1}{2} \left(f_{p,u} - g_{uu,rp} + g^{mn} f_m E_{np} - \frac{1}{2} g^{mn} g_{um} f_n f_p \right) + X_p + \frac{1}{D-2} g_{up} \left[\frac{1}{D-3} sR + g^{mn} \left(f_{m||n} + \frac{1}{2} f_m f_n \right) \right] + g_{uu} \Theta_{,p} + \Theta g_{uu,p}, \quad (44)$$

$$X_{pmq} = g_{p[m,u]||q]} + g_{u[q,m]||p]} + e_{p[m} f_{q]} - g_{u[q} f_{m]||p]} - \frac{1}{2} g_{u[q} f_m] f_p, \quad (45)$$

$$\begin{aligned}
W_{pq} = & -\frac{1}{2}g_{uu}\|p\|q - \frac{1}{2}g_{pq,uu} + g_{u(p,u)\|q} - \left(\frac{1}{2}g_{uu,r} - \Theta g_{uu}\right)e_{pq} - g_{uu,r(p}g_{q)u} \\
& + \frac{1}{2}g_{uu,(p}f_{q)} + \frac{1}{2}g_{uu}f_{(p}\|q)} + g_{u(p}f_{q),u} + \frac{1}{4}g^{mn}g_{um}g_{un}f_p f_q \\
& - \frac{1}{2}g^{mn}g_{um}f_n g_{u(p}f_{q)} + g^{mn}\left(E_{mp}E_{nq} + f_m E_n(p}g_{q)u} - g_{um}E_n(p}f_{q)}\right) \\
& + \frac{1}{D-2}g_{up}g_{uq}\left[\frac{1}{D-3}S_R + g^{mn}\left(f_{m\|n} + \frac{1}{2}f_m f_n\right)\right] \\
& + 2g_{uu}g_{u(p}\Theta_{,q)} + 2\Theta g_{uu,(p}g_{q)u}.
\end{aligned} \tag{46}$$

Consequently, in view of table 1, the explicit conditions for the subtypes III(a) and III(b) with the triple WAND \mathbf{k} are:

- *Subtype III(a)* $\Leftrightarrow \Psi_{3T^i} = 0 \Leftrightarrow V_p = 0 \Leftrightarrow$ the function $g_{uu,rp}$ is explicitly given as:

$$\begin{aligned}
g_{uu,rp} = & f_{p,u} + g^{mn}f_m E_{np} - \frac{1}{2}g^{mn}g_{um}f_n f_p + \frac{2}{D-3}X_p \\
& + \frac{2}{D-2}g_{up}\left[\frac{1}{D-3}S_R + g^{mn}\left(f_{m\|n} + \frac{1}{2}f_m f_n\right)\right] \\
& + 2g_{uu}\Theta_{,p} + 2\Theta g_{uu,p}.
\end{aligned} \tag{47}$$

This is a specific restriction on the spatial derivatives of the function $g_{uu,r}$.

- *Subtype III(b)* $\Leftrightarrow \tilde{\Psi}_{3\dot{y}^k} = 0$:

$$X_{pmq} = \frac{2}{D-3}g_{p[m}X_{q]}, \quad \text{where } X_q = g^{pm}X_{pmq} \tag{48}$$

and X_{pmq} is given by expression (45). Using the fact that any two-dimensional metric g_{pq} is conformally flat, $g_{pq} = \Omega \delta_{pq}$, it can be easily checked that the condition (48) is identically satisfied in $D = 4$.

4.3. Type N with a quadruple WAND \mathbf{k}

When *both* conditions (47) and (48) are satisfied, the only remaining Weyl scalar is $\Psi_{\dot{y}^i} = m_i^p m_j^q \left(W_{pq} - \frac{1}{D-2}g_{pq}W\right)$, see (22). In such a case we obtain the Robinson–Trautman or Kundt spacetimes of algebraic type N with the quadruple WAND $\mathbf{k} = \partial_r$. Using $V_p = 0$, that is by substituting (47) into (46), the function W_{pq} for such type N geometries reduces to a simple expression

$$\begin{aligned}
W_{pq} = & -\frac{1}{2}g_{uu}\|p\|q - \frac{1}{2}g_{pq,uu} + g_{u(p,u)\|q} - \left(\frac{1}{2}g_{uu,r} - \Theta g_{uu}\right)e_{pq} \\
& + \frac{1}{2}g_{uu,(p}f_{q)} + \frac{1}{2}g_{uu}f_{(p}\|q)} + \frac{1}{4}g^{mn}g_{um}g_{un}f_p f_q
\end{aligned}$$

$$\begin{aligned}
& -\frac{1}{D-2} g_{up} g_{uq} \left[\frac{1}{D-3} sR + g^{mn} \left(f_{m|n} + \frac{1}{2} f_m f_n \right) \right] \\
& -\frac{2}{D-3} X_{(p} g_{q)u} + g^{mn} \left(E_{mp} E_{nq} - g_{um} E_{n(p} f_{q)} \right), \tag{49}
\end{aligned}$$

determined by the metric functions (1) and their first and second derivatives. The set of functions W_{pq} directly encodes the amplitudes $\Psi_{\dot{a}\dot{b}}$ of the corresponding gravitational waves, forming a symmetric traceless matrix of dimension $(D-2) \times (D-2)$.

4.4. Type O geometries

The Weyl tensor vanishes completely if, and only if, *all* the above conditions are satisfied and, *in addition*, $\Psi_{\dot{a}\dot{b}} = 0$. This clearly occurs when

$$W_{pq} = \frac{1}{D-2} g_{pq} W, \tag{50}$$

with $W = g^{pq} W_{pq}$, which is a restriction on the functions W_{pq} given by (49).

5. Secondary alignment types of the Weyl tensor

In the non-twisting and shear-free geometries there may also exist an *additional* WAND l distinct from the (multiple) WAND $k = \mathbf{k} = \partial_r$. In such cases the spacetimes are of type I_i , II_i , III_i or D, see table 2.

In general, when $\Psi_{\dot{a}\dot{b}} = m_i^p m_j^q \left(W_{pq} - \frac{1}{D-2} g_{pq} W \right) = 0$ with W_{pq} given by (28), the geometry is of type I_i (the subtype $I(b)_i$, in fact). In such a case $k = \mathbf{k} = \partial_r$ and $l = \frac{1}{2} g_{uu} \partial_r + \partial_u$ are two distinct WANDs, see (7).²

When the geometry is of type II with the double WAND k and $\Psi_{\dot{a}\dot{b}} = 0$, where W_{pq} is given by (38), the geometry is of type II_i with another WAND l . If the additional conditions (40)–(43) are satisfied, we obtain the subtypes $II(a)_i$, $II(b)_i$, $II(c)_i$ and $II(d)_i$, respectively (or their combinations).

The type III_i geometry is equivalent to $II(abcd)_i$, in which case there is the triple WAND k and an additional WAND l . The subtypes $III(a)_i$ and $III(b)_i$ occur if conditions (47) and (48) are also satisfied. For such spacetimes, the only non-vanishing Weyl scalars are $\tilde{\Psi}_{3\dot{a}\dot{b}\dot{c}}$ and Ψ_{3T^i} , respectively.

Finally, there is also the ‘degenerate’ case $D \equiv II_i$ which admits the *double* WAND $k = \partial_r$ and the *double* WAND $l = \frac{1}{2} g_{uu} \partial_r + \partial_u$. Its only non-vanishing Weyl scalars are $\Psi_{2\dot{a}\dot{b}}$, i.e., those of zero-boost-weight. In view of (13)–(22) and (33)–(38), this occurs if (and only if) $V_p = 0$, $X_{pmq} = \frac{2}{D-3} g_{p[m} X_{q]}$, $W_{pq} = \frac{1}{D-2} g_{pq} W$, in which the functions are determined by (36)–(38). Of course, with the additional constraints (40)–(43) we obtain the subtypes D(a), D(b), D(c) and D(d), respectively, and their various combinations. The simplest type D geometry thus seems to be of the subtype D(bcd) for which the only non-vanishing Weyl scalar is $\Psi_{2S} = \frac{D-3}{D-1} P$ with P given by expression (33). This involves, for example, generalizations of Schwarzschild black hole spacetimes, see section 8.1.

² There may exist other WANDs distinct from $l = \frac{1}{2} g_{uu} \partial_r + \partial_u$, see e.g. the case $D > 4$ in section 8.1. However, their systematic investigation is not the topic of the present work.

6. Kundt geometries

We will now discuss the two distinct important subclasses of non-twisting shear-free geometries, namely the non-expanding Kundt and the expanding Robinson–Trautman metrics (in the next section).

The Kundt family is defined by having a *vanishing expansion*, $\Theta = 0$. It implies that the spatial metric g_{pq} in (1) is r -independent,

$$g_{pq} \equiv h_{pq}(u, x), \quad (51)$$

see end of section 2. This significantly simplifies the Riemann tensor (A.17)–(A.26), the Ricci tensor (A.27)–(A.32) and the Weyl tensor (A.34)–(A.43) listed explicitly in appendix A. In fact, it is a complete generalization of the analogous results presented previously in [13] since *no field equations* and *no constraints on algebraically special types* have been employed. The curvature tensor components given in appendix A thus characterize the most general Kundt geometry, which is of algebraic type I(b), whereas the results in appendix B of [13] are only valid for Kundt spacetimes of type II (or more special).

For such a generic Kundt geometry the Weyl tensor frame components are given by (13)–(22) with (23)–(28), simplified by setting $\Theta = 0$. These represent an explicit form of the expressions (7)–(16) written in [13].

In the case of a vanishing expansion we may fully express the conditions discussed in section 4 for algebraically special Kundt geometries with respect to the WAND $\mathbf{k} = \partial_r$. Specifically, we integrate equations (31) and (40), obtaining an explicit r -dependence of the metric functions g_{up} and g_{uv} , respectively. After substituting them into the remaining conditions and separating in r , we obtain:

- The type I Kundt geometry is of subtype I(a) = I(ab) \equiv II $\Leftrightarrow \Psi_{1T^i} = 0 \Leftrightarrow$

$$g_{up} = e_p(u, x) + f_p(u, x) r, \quad (52)$$

where e_p and f_p are arbitrary functions of the coordinates u and x .

- The type II \equiv I(ab) Kundt geometry is of subtype II(a) $\Leftrightarrow \Psi_{2S} = 0 \Leftrightarrow$

$$g_{uu} = a(u, x) r^2 + b(u, x) r + c(u, x), \quad (53)$$

where $a = \frac{1}{4}f^p f_p - \frac{1}{D-2} \left(\frac{1}{D-3} \mathcal{S}R + g^{pq} f_{pq} \right)$ with $f^p \equiv g^{pq} f_q$, $f_{pq} \equiv f_{(p||q)} + \frac{1}{2}f_p f_q$, b and c are arbitrary functions of the coordinates u and x .

- The type II Kundt geometry is of subtype II(b) $\Leftrightarrow \tilde{\Psi}_{2T^{(ij)}} = 0 \Leftrightarrow$

$$\mathcal{S}R_{pq} - \frac{g_{pq}}{D-2} \mathcal{S}R = -\frac{1}{2}(D-4) \left(f_{pq} - \frac{g_{pq}}{D-2} g^{mn} f_{mn} \right). \quad (54)$$

- The type II Kundt geometry is of subtype II(c) $\Leftrightarrow \tilde{\Psi}_{2^{ijkl}} = 0 \Leftrightarrow$

$$\mathcal{S}C_{mpnq} = 0. \quad (55)$$

- The type II Kundt geometry is of subtype II(d) $\Leftrightarrow \Psi_{2ij} = 0 \Leftrightarrow$

$$F_{pq} \equiv f_{[p,q]} = 0. \quad (56)$$

- The type III \equiv II(abcd) Kundt geometry is of subtype III(a) $\Leftrightarrow \Psi_{3T^i} = 0 \Leftrightarrow$

$$a_{,q} + f_q a = 0, \quad b_{,q} - f_{q,u} = T_q, \quad (57)$$

where

$$T_q \equiv -2 e_q \left(a - \frac{1}{4} f^p f_p \right) - \frac{1}{2} e^p f_p f_q + f^p E_{pq} + \frac{2}{D-3} X_q, \quad (58)$$

with $e^p \equiv g^{pq} e_q$, $e_{pq}^k \equiv e_{(p||q)} - \frac{1}{2} g_{pq,u}$, $E_{pq} = e_{[p,q]} + \frac{1}{2} g_{pq,u}$, $X_q \equiv g^{pm} X_{pmq}$ and

$$X_{pmq} = g_{p[m,u||q]} + e_{[q,m]||p} + e_{p[m]f_q}^k - e_{[q]f_m]p}. \quad (59)$$

- The type III Kundt geometry is of subtype III(b) $\Leftrightarrow \check{\Psi}_{3^{jk}} = 0 \Leftrightarrow$

$$X_{pmq} = \frac{1}{D-3} \left(g_{pm} X_q - g_{pq} X_m \right). \quad (60)$$

- The type N \equiv III(ab) Kundt geometry is completely described by the symmetric traceless matrix $\Psi_{\dot{a}\dot{b}}$, which is determined by

$$\begin{aligned} W_{pq} = & -\frac{1}{2} c_{||p||q} + \frac{1}{2} c_{(p} f_q) + \frac{1}{2} c f_{(p||q)} - \frac{1}{2} b e_{pq}^k + \left(a - \frac{1}{4} f^n f_n \right) e_p e_q + e_{(p,u||q)} \\ & - \frac{1}{2} g_{pq,uu} + \frac{1}{4} e^n e_n f_p f_q - e^n E_{n(p} f_q) + g^{mn} E_{mp} E_{nq} - \frac{2}{D-3} X_{(p} e_q) \\ & - \frac{1}{2} \left(2a e_{pq}^k + T_{(p||q)} + T_{(p} f_q) - f_{(p,u||q)} - f_{(p} f_q)_{,u} \right) r. \end{aligned} \quad (61)$$

- The type N becomes type O $\Leftrightarrow \Psi_{\dot{a}\dot{b}} = 0 \Leftrightarrow W_{pq} = \frac{1}{D-2} g_{pq} W$ with $W \equiv g^{pq} W_{pq}$.

These are the same conditions as those presented in [13]. For Kundt geometries of type II (whose metric functions g_{up} satisfy the condition (52)) the shorthands (35), (39), (37) used in this paper become

$$F_{pq} = f_{[p||q]}, \quad (62)$$

$$f_{pq} = f_{(p||q)} + \frac{1}{2} f_p f_q, \quad (63)$$

$$E_{pq} = e_{[p||q]} + \frac{1}{2} g_{pq,u} + r f_{[p||q]}, \quad (64)$$

$$e_{pq} = e_{(p||q)} - \frac{1}{2} g_{pq,u} + r f_{(p||q)}, \quad (65)$$

$$\begin{aligned} X_{pmq} = & e_{[q||m]||p} + F_{qm} e_p + F_{p[m} e_q] + e_{p[m} f_q] - f_{p[m} e_q] + g_{p[m,u||q]} \\ & + r \left(f_{[q||m]||p} + F_{qm} f_p + F_{p[m} f_q] \right). \end{aligned} \quad (66)$$

The identification is

$$F_{pq} \equiv F_{pq}^k, \quad (67)$$

$$f_{pq} \equiv f_{pq}^{\kappa}, \quad (68)$$

$$E_{pq} \equiv E_{pq}^{\kappa} + r F_{pq}^{\kappa}, \quad (69)$$

$$e_{pq} \equiv e_{pq}^{\kappa} + r f_{(p||q)}, \quad (70)$$

$$X_{pmq} \equiv X_{pmq}^{\kappa} + r Y_{pmq}^{\kappa}, \quad (71)$$

where the superscript κ denotes the quantities defined and employed in [13].

7. Particular Robinson–Trautman geometries

The algebraic structure of generic Robinson–Trautman geometries with an arbitrary $\Theta \neq 0$ has been described in sections 3 and 4. Let us now investigate in detail a large particular subclass such that the non-twisting shear-free congruence generated by the null vector field $\mathbf{k} = \partial_r$ has an expansion of the form

$$\Theta = \frac{1}{r}. \quad (72)$$

This is an important subcase since $\Theta_{,r} + \Theta^2 = 0$ and thus, in view of (A.17) and (A.27), there is $R_{rprq} = 0 = R_{rr}$. Consequently, $R_{abcd} k^a m_i^b k^c m_j^d = 0 = R_{ab} k^a k^b$ which means that such Robinson–Trautman geometries are of Riemann type I and also of Ricci type I.

For the case (72) we can explicitly integrate all the conditions with respect to r and determine the algebraic types and subtypes. First, the spatial metric g_{pq} becomes

$$g_{pq} = r^2 h_{pq}(u, x), \quad (73)$$

see (6) for $R = r$, which is obtained by solving $R_{,r} = \Theta R$. Such geometries are, in general, of the Weyl (sub)type I(b).

The metrics are of the Weyl type II (or more special) with the double WAND \mathbf{k} if, and only if, the condition $\Psi_{1T^i} = 0$ is satisfied. For (72) we have $f_{p,r} = 0$ due to (32), i.e., the functions f_p are *independent of r* . By integrating (31) we then obtain

$$g_{up} = e_p(u, x) r^2 - f_p(u, x) r, \quad (74)$$

where e_p and f_p are arbitrary functions of the coordinates u and x .

With the conditions (72)–(74), the functions (33)–(38) determining the Weyl scalars reduce to

$$P = \left(\frac{1}{2} g_{uu,r} - g_{uu} r^{-1} \right)_{,r} - \alpha r^{-2}, \quad (75)$$

$$Q_{pq} = R_{pq} + \frac{1}{2} (D - 4) \left(f_{(p||q)} + \frac{1}{2} f_p f_q \right), \quad (76)$$

$$F_{pq} = f_{[p,q]}, \quad (77)$$

$$\begin{aligned}
V_p = & -\frac{1}{2}g_{uu,rp} + g_{uu,p} r^{-1} - (e_p r^2 - f_p r) \left(\frac{1}{2}g_{uu,rr} - g_{uu,r} r^{-1} + g_{uu} r^{-2} \right) \\
& + \frac{1}{2}f_{p,u} + \frac{1}{2}f^n E_{np}^{\text{RT}} + \frac{1}{2} \frac{D-4}{D-3} f^n e_{[p} f_{n]} - \frac{1}{2} \frac{D-6}{D-3} f^n F_{np} r^{-1} \\
& - \frac{1}{D-3} h^{mn} \left[h_{m[p,u]n} + e_{[m,p]n} + e_m^{\text{RT}} f_{n]} \right] \\
& + \frac{1}{2} \left(e_p f_{m|n} - e_m f_{(n|p)} \right) + \frac{3}{2} e_m F_{np} - F_{mp|n} r^{-1} \Big], \tag{78}
\end{aligned}$$

$$X_{pmq} = X_{pmq}^{\text{RT}} r^2 + Y_{pmq}^{\text{RT}} r, \tag{79}$$

$$\begin{aligned}
W_{pq} = & -\frac{1}{2}g_{uu|p|q} + e_{pq}^{\text{RT}} \left(g_{uu} r - \frac{1}{2}g_{uu,r} r^2 \right) - \frac{1}{2}f_{(p|q)} \left(g_{uu} - g_{uu,r} r \right) \\
& - g_{uu,r(p} e_{q)} r^2 + 2g_{uu,(p} e_{q)} r - \frac{3}{2}g_{uu,(p} f_{q)} + g_{uu,r(p} f_{q)} r \\
& - \left(e_p e_q r^2 - 2e_{(p} f_{q)} r + f_p f_q \right) \left(\frac{1}{2}g_{uu,rr} r^2 - g_{uu,r} r + g_{uu} \right) \\
& + \left[e_{(p,u|q)} + e_{(p} f_{q),u} - \frac{1}{2}h_{pq,uu} + \frac{1}{4} \left(e^n e_n f_p f_q + f^n f_n e_p e_q \right) - \frac{1}{2}e^n f_n e_{(p} f_{q)} \right. \\
& \left. + h^{mn} E_{mp}^{\text{RT}} E_{nq}^{\text{RT}} + f^n E_{n(p}^{\text{RT}} e_{q)} - e^n E_{n(p}^{\text{RT}} f_{q)} \right] r^2 \\
& - \left[f_{(p,u|q)} + f_{(p} f_{q),u} - 2h^{mn} E_{m(p}^{\text{RT}} F_{q)n} + f^n F_{n(p} e_{q)} - e^n F_{n(p} f_{q)} \right] r \\
& + h^{mn} F_{mp} F_{nq}, \tag{80}
\end{aligned}$$

where $\mathcal{R} = {}^S R r^2$ and $\mathcal{R}_{pq} = {}^S R_{pq}$ are the Ricci scalar and the Ricci tensor with respect to the Riemannian metric $h_{pq}(u, x)$, respectively,

$$\alpha(u, x) \equiv \frac{1}{4}f^p f_p - \frac{1}{D-2} \left(\frac{1}{D-3} \mathcal{R} + f^p{}_{||p} + \frac{1}{2}f^p f_p \right), \tag{81}$$

$$X_{pmq}^{\text{RT}} \equiv h_{p[m,u]q} + e_{[q,m]||p} + e_{p[m}^{\text{RT}} f_{q]} - e_{[q} f_{m]||p} - e_p F_{mq} - \frac{1}{2}e_{[q} f_{m]} f_p, \tag{82}$$

$$Y_{pmq}^{\text{RT}} \equiv -F_{qm||p} + f_{[q} F_{m]p} + f_p F_{mq}, \tag{83}$$

and

$$e^p \equiv h^{pq} e_q, \quad f^p \equiv h^{pq} f_q, \tag{84}$$

$$e_{pq}^{\text{RT}} \equiv e_{(p|q)} - \frac{1}{2}h_{pq,u}, \quad E_{pq}^{\text{RT}} \equiv e_{[p|q]} + \frac{1}{2}h_{pq,u}, \tag{85}$$

$$X_q^{\text{RT}} \equiv h^{pm} X_{pmq}^{\text{RT}}, \quad Y_q^{\text{RT}} \equiv h^{pm} Y_{pmq}^{\text{RT}}. \tag{86}$$

Using the results of sections 4.1–4.4 we can thus explicitly express the conditions for the principal alignment (sub)types of the algebraically special Weyl tensor:

- The type II Robinson–Trautman geometry is of subtype II(a) $\Leftrightarrow \Psi_{2s} = 0 \Leftrightarrow$

$$g_{uu} = \alpha(u, x) + \beta(u, x) r + \gamma(u, x) r^2, \tag{87}$$

where α is given by (81) while β, γ are arbitrary functions of u and x .

- The type II Robinson–Trautman geometry is of subtype II(b) $\Leftrightarrow \tilde{\Psi}_{2T^{(ij)}} = 0 \Leftrightarrow$

$$\mathcal{R}_{pq} - \frac{h_{pq}}{D-2} \mathcal{R} = -\frac{1}{2}(D-4) \left[\left(f_{(p\parallel q)} + \frac{1}{2} f_p f_q \right) - \frac{h_{pq}}{D-2} \left(f^n_{\parallel n} + \frac{1}{2} f^n f_n \right) \right], \tag{88}$$

where \mathcal{R}_{pq} is the Ricci tensor with respect to the metric h_{pq} .

- The type II Robinson–Trautman geometry is of subtype II(c) $\Leftrightarrow \tilde{\Psi}_{2^{ijkl}} = 0 \Leftrightarrow$

$$C_{mpnq} = 0, \tag{89}$$

where $C_{mpnq} = {}^s C_{mpnq} r^{-2}$ is the Weyl tensor corresponding to the metric h_{pq} .

- The type II Robinson–Trautman geometry is of subtype II(d) $\Leftrightarrow \Psi_{2^{ij}} = 0 \Leftrightarrow$

$$F_{pq} \equiv f_{[p,q]} = 0. \tag{90}$$

- The type III Robinson–Trautman geometry is of subtype III(a) $\Leftrightarrow \Psi_{3T^i} = 0 \Leftrightarrow$

$$\alpha_{,q} + f_q \alpha = 0, \quad \beta_{,q} + f_{q,u} = \mathcal{T}_q, \tag{91}$$

where

$$\mathcal{T}_q \equiv 2 e_q \left(\alpha - \frac{1}{4} f^p f_p \right) + \frac{1}{2} e^p f_p f_q - f^p E_{pq}^{\text{RT}} - \frac{2}{D-3} X_q, \tag{92}$$

with $X_q = h^{pm} X_{pmq}^{\text{RT}} = g^{pm} X_{pmq}$,

$$X_{pmq}^{\text{RT}} = h_{p[m,u\parallel q]} + e_{[q,m]\parallel p} + e_{p[m}^{\text{RT}} f_{q]} - e_{[q} f_{m]\parallel p} - \frac{1}{2} e_{[q} f_{m]} f_p. \tag{93}$$

- The type III Robinson–Trautman geometry is of subtype III(b) $\Leftrightarrow \tilde{\Psi}_{3^{ijk}} = 0 \Leftrightarrow$

$$X_{pmq}^{\text{RT}} = \frac{1}{D-3} (h_{pm} X_q - h_{pq} X_m). \tag{94}$$

- The type N Robinson–Trautman geometry is described by the symmetric traceless matrix $\Psi_{4^{ij}}$, which is completely determined by

$$\begin{aligned} W_{pq} = & \left(-\frac{1}{2} \gamma_{\parallel p\parallel q} + \frac{1}{2} \gamma_{,(p} f_{q)} + \frac{1}{2} \gamma f_{(p\parallel q)} + \frac{1}{2} \beta e_{pq}^{\text{RT}} + \left(\alpha - \frac{1}{4} f^n f_n \right) e_p e_q + e_{(p,u\parallel q)} \right. \\ & \left. - \frac{1}{2} h_{pq,uu} + \frac{1}{4} e^n e_n f_p f_q - e^n E_{n(p}^{\text{RT}} f_{q)} + h^{mn} E_{mp}^{\text{RT}} E_{nq}^{\text{RT}} - \frac{2}{D-3} X_{(p} e_{q)} \right) r^2 \\ & + \frac{1}{2} \left(2\alpha e_{pq}^{\text{RT}} - \mathcal{T}_{(p\parallel q)} - \mathcal{T}_{(p} f_{q)} - f_{(p,u\parallel q)} - f_{(p} f_{q),u} \right) r. \end{aligned} \tag{95}$$

- The type N becomes type O $\Leftrightarrow \Psi_{4^{ij}} = 0 \Leftrightarrow W_{pq} = \frac{1}{D-2} h_{pq} h^{mn} W_{mn}$, where W_{pq} is given by (95).

This completes the classification of *principal* alignment (sub)types of the Weyl tensor of Robinson–Trautman geometries with expansion (72) and the multiple WAND \mathbf{k} .

The *secondary* alignment (sub)types with the additional WAND $l = \frac{1}{2}g_{uu}\partial_r + \partial_u$ are obtained when the conditions summarized in table 2 are satisfied, namely:

- The type II Robinson–Trautman geometry is of type $\text{II}_i \Leftrightarrow \Psi_{4\dot{\psi}} = 0 \Leftrightarrow$

$$W_{pq} = \frac{1}{D-2} h_{pq} h^{mn} W_{mn}, \quad (96)$$

where W_{pq} is given by (80).

- The type III Robinson–Trautman geometry is of type $\text{III}_i \Leftrightarrow \Psi_{4\dot{\psi}} = 0 \Leftrightarrow$ the condition (96) is satisfied, where W_{pq} is given by

$$\begin{aligned} W_{pq} = & \left[-\frac{1}{2}\gamma_{\parallel p\parallel q} + \frac{1}{2}\gamma_{(p}f_{q)} + \frac{1}{2}\gamma f_{(p\parallel q)} + \frac{1}{2}\beta e_{pq}^{\text{RT}} + \beta_{(p}e_{q)} - \left(\alpha - \frac{1}{4}f^n f_n\right) e_p e_q \right. \\ & + e_{(p,u\parallel q)} + e_{(p}f_{q),u} - \frac{1}{2}h_{pq,uu} + \frac{1}{4}e^n e_n f_p f_q - \frac{1}{2}e^n f_n e_{(p}f_{q)} \\ & \left. + h^{mn} E_{mp}^{\text{RT}} E_{nq}^{\text{RT}} + f^n E_{n(p}^{\text{RT}} e_{q)} - e^n E_{n(p}^{\text{RT}} f_{q)} \right] r^2 \\ & - \left(\frac{1}{2}\beta_{\parallel p\parallel q} + \frac{1}{2}\beta_{(p}f_{q)} - \alpha e_{pq}^{\text{RT}} - 2\alpha_{(p}e_{q)} - 2\alpha e_{(p}f_{q)} + f_{(p,u\parallel q)} + f_{(p}f_{q),u} \right) r \\ & - \frac{1}{2}\alpha_{\parallel p\parallel q} - \frac{1}{2}\alpha f_{(p\parallel q)} - \frac{3}{2}\alpha_{(p}f_{q)} - \alpha f_p f_q. \end{aligned} \quad (97)$$

This, in fact, is a general form of W_{pq} for the type III spacetimes.

- The type II Robinson–Trautman geometry is of type D with respect to the double WAND $l = \frac{1}{2}g_{uu}\partial_r + \partial_u \Leftrightarrow \Psi_{3T^i} = \tilde{\Psi}_{3^{jk}} = 0$ and $\Psi_{4\dot{\psi}} = 0 \Leftrightarrow$

$$\begin{aligned} X_{pmq}^{\text{RT}} &= \frac{1}{D-3} (h_{pm} X_q^{\text{RT}} - h_{pq} X_m^{\text{RT}}), \\ Y_{pmq}^{\text{RT}} &= \frac{1}{D-3} (h_{pm} Y_q^{\text{RT}} - h_{pq} Y_m^{\text{RT}}), \\ V_p &= 0, \quad W_{pq} = \frac{1}{D-2} h_{pq} h^{mn} W_{mn}, \end{aligned} \quad (98)$$

where the corresponding functions are given by (78), (80), (82), (83), (86). The particular subtypes D(a), D(b), D(c), D(d) and their various combinations occur if the additional conditions (87)–(90) are also valid.

8. Example: vacuum Robinson–Trautman spacetimes

Motivated by our previous studies [7, 9] of Robinson–Trautman spacetimes in general relativity (extended to any dimension $D \geq 4$), let us consider a metric of the form

$$ds^2 = r^2 h_{pq} (dx^p + e^p du)(dx^q + e^q du) - 2 du dr - g^{rr} du^2, \quad (99)$$

where

$$g^{rr} = \frac{\mathcal{R}}{(D-2)(D-3)} + \frac{b(u)}{r^{D-3}} - \frac{2}{D-2} \left(e^p \parallel_p - \frac{1}{2} h^{pq} h_{pq,u} \right) r - \frac{2\Lambda}{(D-1)(D-2)} r^2. \quad (100)$$

In fact, this is the most general Robinson–Trautman vacuum line element in Einstein’s theory (extended to an arbitrary dimension D), with a cosmological constant Λ and possibly a pure radiation field aligned with $\mathbf{k} = \partial_r$.

Employing the results of the previous section, it is straightforward to obtain explicit conditions under which this geometry becomes of a specific algebraic type. Since

$$f_p = 0 \quad (101)$$

and $g_{up} = e_p(u, x)r^2$, the condition (74) is satisfied, so that the spacetime is of Weyl type II (or more special) with respect to the multiple WAND \mathbf{k} . Moreover, $g_{uu} = r^2 e^n e_n - g^{rr}$, that is

$$g_{uu} = -a - b r^{3-D} - c r + \gamma r^2, \quad (102)$$

where

$$a = \frac{\mathcal{R}}{(D-2)(D-3)}, \quad (103)$$

$$c = -\frac{2}{D-2} \left(e^n \parallel_n - \frac{1}{2} h^{mn} h_{mn,u} \right), \quad (104)$$

$$\gamma = e^n e_n + \frac{2\Lambda}{(D-1)(D-2)}. \quad (105)$$

For $f_p = 0$ with g_{uu} of the form (102) we obtain from (75)–(80) that

$$P = -\frac{1}{2}(D-1)(D-2)b r^{1-D}, \quad (106)$$

$$Q_{pq} = \mathcal{R}_{pq}, \quad (107)$$

$$F_{pq} = 0, \quad (108)$$

$$V_p = \left(a e_p - \frac{1}{2} c_{,p} + \frac{1}{D-3} X_p \right) - a_{,p} r^{-1} + \frac{1}{2}(D-1)(D-2)b e_p r^{3-D}, \quad (109)$$

$$X_{pmq} = \left(h_{p[m,u]q} + e_{[q,m]p} \right) r^2, \quad (110)$$

$$\begin{aligned}
 W_{pq} = & \frac{1}{2} a_{\|p\|q} + \left(\frac{1}{2} c_{\|p\|q} - a e_{pq}^{\text{RT}} - 2a_{,(p} e_{q)} \right) r \\
 & + \left(-\frac{1}{2} (e^n e_n)_{\|p\|q} - \frac{1}{2} c e_{pq}^{\text{RT}} + a e_p e_q - c_{,(p} e_{q)} \right. \\
 & \left. + e_{(p,u\|q)} - \frac{1}{2} h_{pq,uu} + h^{mn} E_{mp}^{\text{RT}} E_{nq}^{\text{RT}} \right) r^2 \\
 & - \frac{1}{2} (D-1) b e_{pq}^{\text{RT}} r^{4-D} + \frac{1}{2} (D-1)(D-2) b e_p e_q r^{5-D}. \quad (111)
 \end{aligned}$$

In view of (13)–(22), the metric (99), (100) is thus of

- subtype II(a) $\Leftrightarrow b(u) = 0,$ (112)

- subtype II(b) $\Leftrightarrow \mathcal{R}_{pq} = \frac{h_{pq}}{D-2} \mathcal{R},$ (113)

- subtype II(c) $\Leftrightarrow C_{mpnq} = 0,$ (114)

- subtype II(d) always. (115)

Interestingly, it follows from the contraction of Bianchi identities and condition (113), which is identically satisfied in $D = 4$, that

$$0 = h^{mn} h^{pq} (\mathcal{R}_{mpnq\|k} + \mathcal{R}_{mpqk\|n} + \mathcal{R}_{mpkn\|q}) = \frac{D-4}{D-2} \mathcal{R}_{,k}. \quad (116)$$

Thus, any Robinson–Trautman $D > 4$ geometry (99), (100) of algebraic subtype II(b) must have $\mathcal{R}_{,k} = 0$. Moreover

$$C_{mpnq} = \mathcal{R}_{mpnq} - \frac{1}{(D-2)(D-3)} \mathcal{R} (h_{mn} h_{pq} - h_{mq} h_{np}). \quad (117)$$

The condition $C_{mpnq} = 0$ for subtype II(c) is always satisfied in $D = 4$ and $D = 5$.

We thus immediately infer that subtype II(bc) \equiv II(bcd) occurs if, and only if, the $(D-2)$ -dimensional transverse space has a *constant curvature*, that is

$$\mathcal{R}_{mpnq} = \frac{1}{(D-2)(D-3)} \mathcal{R}(u) (h_{mn} h_{pq} - h_{mq} h_{np}). \quad (118)$$

In such a case the metric can be written as $h_{pq} = P^{-2} \delta_{pq}$, where $P = 1 + \frac{1}{4} K \delta_{mn} x^m x^n$, $K = \frac{1}{(D-2)(D-3)} \mathcal{R}(u)$, see [7, 8].

The conditions for type III subtypes are then

- subtype III(a) $\Leftrightarrow \mathcal{R}_{,q} = 0$ and $c_{,q} = 2 a e_q + \frac{2}{D-3} X_q,$ (119)

where $X_q = h^{pm} X_{pmq}^{\text{RT}}, \quad X_{pmq}^{\text{RT}} = h_{p[m,u\|q]} + e_{[q,m]\|p},$

- subtype III(b) $\Leftrightarrow X_{pmq}^{\text{RT}} = \frac{1}{D-3} (h_{pm} X_q - h_{pq} X_m).$ (120)

The type N is obtained by applying *all* the conditions (112)–(115) and (119)–(120). In such a case, the only non-trivial function (111) reduces to

$$\begin{aligned}
W_{pq} = & \left(\frac{1}{2} c_{\|p\|q} - a e_{pq}^{\text{RT}} \right) r \\
& + \left(-\frac{1}{2} (e^n e_n)_{\|p\|q} - \frac{1}{2} c e_{pq}^{\text{RT}} + a e_p e_q - c_{,[p} e_{q]} \right. \\
& \left. + e_{(p,u\|q)} - \frac{1}{2} h_{pq,uu} + h^{mn} E_{mp}^{\text{RT}} E_{nq}^{\text{RT}} \right) r^2.
\end{aligned} \tag{121}$$

Such a geometry is of type O when $W_{pq} = \frac{1}{D-2} h_{pq} h^{mn} W_{mn}$.

The secondary alignment types II_i , III_i , D arise when the conditions (96)–(98) are satisfied, respectively, in which the key functions are given by (109)–(111).

We thus conclude that the Robinson–Trautman geometry of the form (99), (100) generally admits *all* the above mentioned algebraic types and subtypes. Of course, specific *field equations impose additional constraints* that may exclude some of the (sub)types. To illustrate this effect, let us now restrict ourselves to the most important case, namely vacuum spacetimes in the *Einstein theory*.

8.1. Most general Robinson–Trautman vacuum spacetimes

As shown in [9], a fully general Robinson–Trautman vacuum solution in the Einstein theory (including Λ) is given by (99), (100) where the metric functions are restricted by the constraints

$$\mathcal{R}_{pq} = \frac{1}{D-2} h_{pq} \mathcal{R}, \tag{122}$$

$$h_{pq,u} = 2 e_{(p\|q)} + c h_{pq}, \tag{123}$$

$$(D-4)\mathcal{R}_{,p} = 0, \tag{124}$$

$$h^{mn} a_{\|m\|n} + \frac{1}{2}(D-1)(D-2)b c + (D-2)b_{,u} = 0, \tag{125}$$

with a and c defined in (103) and (104).

Using conditions (122)–(125) and

$$e_{pq}^{\text{RT}} = -\frac{1}{2} c h_{pq}, \quad E_{pq}^{\text{RT}} = e_{p\|q} + \frac{1}{2} c h_{pq}, \tag{126}$$

$$h_{pq,uu} = 2c e_{(p\|q)} + 2e_{(p\|q),u} + (c^2 + c_{,u})h_{pq}, \tag{127}$$

$$X_{pmq}^{\text{RT}} = e^n \mathcal{R}_{npmq} + c_{,[q} h_{m]p}, \quad X_q = (D-3) \left(-a e_q + \frac{1}{2} c_{,q} \right), \tag{128}$$

functions (109)–(111) which, together with (106)–(108), characterize the algebraic structure of the Weyl tensor become

$$V_p = -a_{,p} r^{-1} + \frac{1}{2}(D-1)(D-2)b e_p r^{3-D}, \tag{129}$$

$$X_{pmq} = \left(e^n \mathcal{R}_{npmq} + c_{,[q} h_{m]p} \right) r^2, \tag{130}$$

$$\begin{aligned}
W_{pq} = & \frac{1}{2} a_{\parallel p \parallel q} + \left(\frac{1}{2} c_{\parallel p \parallel q} - 2a_{, (p} e_{q)} \right) r \\
& + e^m e^n C_{mpnq} r^2 + \frac{1}{2} (D-1)(D-2) b e_p e_q r^{5-D} \\
& + h_{pq} \left[\frac{1}{2} a c r + \left(a e^n e_n - \frac{1}{2} e^n c_{,n} - \frac{1}{2} c_{,u} \right) r^2 + \frac{1}{4} (D-1) b c r^{4-D} \right]. \quad (131)
\end{aligned}$$

To derive the last expression we have used the fact that $e_{(p,u\parallel q)} - e_{(p\parallel q),u} = e_m T_{pq}^m$, where the tensor $T_{pq}^m \equiv {}^S \Gamma_{pq,u}^m$ can be written as

$$T_{pq}^m = h^{mn} \left[e_{n\parallel(p\parallel q)} - e^k \mathcal{R}_{k(pq)n} \right] + \delta_{(p}^m c_{,q)} - \frac{1}{2} h^{mn} h_{pq} c_{,n}, \quad (132)$$

see appendix A of [9]. The non-vanishing Weyl tensor components with respect to the frame (7), sorted by the boost weight, are thus

$$\Psi_{2S} = -\frac{1}{2} (D-2)(D-3) b r^{1-D}, \quad (133)$$

$$\tilde{\Psi}_{2^{ijkl}} = m_i^m m_j^p m_k^n m_l^q C_{mpnq} r^2, \quad (134)$$

$$\Psi_{3T^i} = m_i^p \left[\frac{1}{2} (D-1)(D-3) b e_p r^{3-D} - \frac{D-3}{D-2} a_{,p} r^{-1} \right], \quad (135)$$

$$\tilde{\Psi}_{3^{ijk}} = m_i^p m_j^m m_k^q e^n C_{npmq} r^2, \quad (136)$$

$$\begin{aligned}
\Psi_{4^{ij}} = & m_i^p m_j^q \left\{ \frac{1}{2} \left(a_{\parallel p \parallel q} - \frac{1}{D-2} h_{pq} h^{mn} a_{\parallel m \parallel n} \right) + e^m e^n C_{mpnq} r^2 \right. \\
& + \left[\frac{1}{2} \left(c_{\parallel p \parallel q} - \frac{1}{D-2} h_{pq} h^{mn} c_{\parallel m \parallel n} \right) - 2 \left(a_{, (p} e_{q)} - \frac{1}{D-2} h_{pq} e^n a_{,n} \right) \right] r \\
& \left. + \frac{1}{2} (D-1)(D-2) b \left(e_p e_q - \frac{1}{D-2} h_{pq} e^n e_n \right) r^{5-D} \right\}. \quad (137)
\end{aligned}$$

It is now convenient to perform a null rotation of the frame (7) with the privileged null vector \mathbf{k} fixed, $\mathbf{l}' = \mathbf{l} + \sqrt{2} L^i \mathbf{m}_i + |\mathbf{L}|^2 \mathbf{k}$, $\mathbf{m}'_i = \mathbf{m}_i + \sqrt{2} L_i \mathbf{k}$, see (C1) in appendix C of [19], and the parameters $L_i \equiv -\frac{1}{\sqrt{2}} r^2 e_p m_i^p$. The new null frame is

$$\mathbf{k}' = \partial_r, \quad \mathbf{l}' = -\frac{1}{2} g^{rr} \partial_r + \partial_u - e^p \partial_p, \quad \mathbf{m}'_i = m_i^p \partial_p, \quad (138)$$

and using the expressions (C5) in [19] the non-vanishing irreducible Weyl scalars in such a frame simplify considerably to

$$\Psi'_{2S} = -\frac{1}{2} (D-2)(D-3) b r^{1-D}, \quad (139)$$

$$\tilde{\Psi}'_{2^{ijkl}} = m_i^m m_j^p m_k^n m_l^q C_{mpnq} r^2, \quad (140)$$

$$\Psi'_{3T^i} = -m_i^p \frac{D-3}{D-2} a_{,p} r^{-1}, \quad (141)$$

$$\Psi'_{4ij} = \frac{1}{2} m_i^p m_j^q \left[\left(a_{\parallel p \parallel q} + c_{\parallel p \parallel q} r \right) - \frac{1}{D-2} h_{pq} h^{mn} \left(a_{\parallel m \parallel n} + c_{\parallel m \parallel n} r \right) \right], \quad (142)$$

with $\tilde{\Psi}'_{3ijk} = 0$. Notice, interestingly, that the last term can be rewritten as

$$\Psi'_{4ij} = \frac{1}{2} m_i^p m_j^q \left[g_{\parallel p \parallel q}^{rr} - \frac{1}{D-2} h_{pq} h^{mn} g_{\parallel m \parallel n}^{rr} \right]. \quad (143)$$

The gravitational wave amplitude matrix Ψ'_{4ij} (which is symmetric and traceless) is thus directly determined by *the second spatial derivatives of the contravariant metric coefficient g^{rr}* , see (100).

To prove the *non-existence of type N and type II vacuum solutions in $D > 4$* it is now crucial to prove the identity

$$(D-4) \left(c_{\parallel p \parallel q} - \frac{1}{D-2} h_{pq} h^{mn} c_{\parallel m \parallel n} \right) = 0. \quad (144)$$

This follows from the u -derivative of the condition (122), namely

$$\mathcal{R}_{pq,u} = \frac{1}{D-2} \left(h_{pq,u} \mathcal{R} + \frac{1}{D-2} h_{pq} h_{mn} h^{mn}{}_{,u} \mathcal{R} + h_{pq} h^{mn} \mathcal{R}_{mn,u} \right), \quad (145)$$

which using $h^{mn}{}_{,u} = -h^{mp} h^{nq} h_{pq,u}$ and the constraint (123) can be rewritten as

$$\mathcal{R}_{pq,u} = \frac{1}{D-2} \left(2 e_{(p \parallel q)} \mathcal{R} - \frac{2}{D-2} h_{pq} e^n{}_{\parallel n} \mathcal{R} + h_{pq} h^{mn} \mathcal{R}_{mn,u} \right). \quad (146)$$

It remains to evaluate $\mathcal{R}_{pq,u}$. From the definition of the Ricci tensor it follows that

$$\mathcal{R}_{pq,u} = T_{pq \parallel m}^m - T_{pm \parallel q}^m, \quad (147)$$

where $T_{pq}^m \equiv {}^S \Gamma_{pq,u}^m$ is a *tensor* symmetric in p, q and given by (132). Using common relations for commutators of covariant derivatives and contracted Bianchi identities, see (3.2.3), (3.2.21), (3.2.16) in [21], we obtain the derivatives of (132)

$$T_{pq \parallel m}^m = e^n{}_{\parallel n \parallel p \parallel q} + \frac{2}{D-2} e_{(p \parallel q)} \mathcal{R} + \frac{1}{D-2} h_{pq} e^n \mathcal{R}_{,n} + c_{\parallel p \parallel q} - \frac{1}{2} h_{pq} h^{mn} c_{\parallel m \parallel n}, \quad (148)$$

$$T_{pm \parallel q}^m = e^n{}_{\parallel n \parallel p \parallel q} + \frac{1}{2} (D-2) c_{\parallel p \parallel q}. \quad (149)$$

Substituting into (147), the u -derivative of the Ricci tensor becomes

$$\mathcal{R}_{pq,u} = \frac{2}{D-2} e_{(p \parallel q)} \mathcal{R} + \frac{1}{D-2} h_{pq} e^n \mathcal{R}_{,n} - \frac{1}{2} (D-4) c_{\parallel p \parallel q} - \frac{1}{2} h_{pq} h^{mn} c_{\parallel m \parallel n}, \quad (150)$$

and its trace reads $h^{mn} \mathcal{R}_{mn,u} = \frac{2}{D-2} e^n{}_{\parallel n} \mathcal{R} + e^n \mathcal{R}_{,n} - (D-3) h^{mn} c_{\parallel m \parallel n}$. By inserting these two expressions into (146), the identity (144) is proven.

Using (139)–(142) with (124) and (144) it is now easy to determine explicitly the algebraic structure of all vacuum Robinson–Trautman spacetimes in the Einstein theory in any dimension D . The results are summarized in table 4.

In $D > 4$ it follows from (124) that $a_{,p} = 0$, which, together with (144), implies $\Psi'_{3T^i} = 0 = \Psi'_{4ij}$. This proves that *there are no* type N, type III and type II spacetimes in the Robinson–Trautman family in higher dimensions when the Einstein vacuum field equations are applied. Such a result is in full agreement with observations made in [7].

Thus, *all Robinson–Trautman vacuum solutions in $D > 4$ are of type D (or type O)*. The double degenerate WANDs are $k' = \mathbf{k} = \partial_r, l' = -\frac{1}{2} g^{rr} \partial_r + \partial_u - e^p \partial_p$, see (138). It is also

Table 4. The necessary and sufficient conditions for all possible algebraic (sub)types of the Robinson–Trautman vacuum solutions of Einstein’s field equations. Some of them are always satisfied. The admissible algebraic structures of the Weyl tensor differ significantly in the case $D = 4$ and in higher dimensions $D > 4$.

Type	$D = 4$	$D > 4$	
II(a)	$b = 0$	$b = 0$	\Leftrightarrow D(a)
II(b)	always	always	\Leftrightarrow D(b)
II(c)	always	$C_{mpnq} = 0$	\Leftrightarrow D(c)
II(d)	always	always	\Leftrightarrow D(d)
III	II(abcd)		
III(a)	$b = 0 = \mathcal{R}_{,p}$	equivalent to	O
III(b)	always for $b = 0$	equivalent to	O
N	III(ab)		
O	$b = 0 = \mathcal{R}_{,p}$ and $c_{\parallel p\parallel q} = \frac{1}{D-2} h_{pq} h^{mn} c_{\parallel m\parallel n}$	equivalent to	D(ac)
D	$\mathcal{R}_{,p} = 0$ and $c_{\parallel p\parallel q} = \frac{1}{D-2} h_{pq} h^{mn} c_{\parallel m\parallel n}$	always	D(bd)

straightforward to identify the possible subtypes, namely D(a) and D(c) and $O \equiv D(ac)$. In fact, the only non-vanishing Weyl scalars are

$$\Psi'_{2S} = -\frac{1}{2}(D - 2)(D - 3)b r^{1-D}, \tag{151}$$

$$\tilde{\Psi}'_{2^{ijkl}} = m_i^m m_j^p m_k^n m_l^q C_{mpnq} r^2, \tag{152}$$

so that all such spacetimes are of the subtype D(bd). Clearly, there are only two algebraically distinct cases possible, namely

- subtype D(a) \equiv D(abd) $\Leftrightarrow b = 0,$ (153)

- subtype D(c) \equiv D(bcd) $\Leftrightarrow C_{mpnq} = 0.$ (154)

The latter case (which necessarily occurs in dimension $D = 5$) admits just the scalar Ψ'_{2S} given by (151). Moreover, in view of (117) relation (118) must hold which means that the *transverse Riemannian space has a constant curvature*. Such a family of Robinson–Trautman vacuum spacetimes contains generalizations of the Schwarzschild black hole of mass proportional to b . When both conditions (153) and (154) are satisfied, the corresponding spacetime is of type O.

This generalizes, confirms and refines the conclusions of a previous work [7] where the Robinson–Trautman vacuum solutions with $e_p = 0$ were studied, i.e., assuming the metric functions e_p can be *globally removed*. Relation between the respective notations is $h_{pq} = P^{-2}(u, x)\gamma_{pq}(x)$ with $\det \gamma_{pq} = 1$, $b(u) = -\mu(u)$ and $c = -2(\log P)_{,u}$. It follows that the exceptional cases discussed in [7] with the functions μ and/or C_{mpnq} vanishing are, in fact, *algebraically distinct subtypes*.

9. Concluding summary

We investigated the algebraic structure of a fully general class on non-twisting and shear-free geometries in an arbitrary dimension D , that is, the complete Robinson–Trautman and Kundt family. In particular:

- Using the Christoffel symbols we derived all coordinate components of the Riemann, Ricci and Weyl curvature tensors in an explicit form. These are presented in appendix A.
- By projecting the Weyl tensor onto the natural null frame we evaluated the corresponding scalars of all boost weights. In contrast to a complicated form of the coordinate components, such Weyl scalars are, due to cancelation of many terms, surprisingly simple, see equations (13)–(22) with (23)–(28).
- Weyl scalars obtained in this manner directly determine the algebraic structure of the metric (1) with (6). Distinct algebraic types and subtypes are defined by the vanishing of these scalars (and their combinations), see tables 1 and 2.
- We proved that all non-twisting shear-free geometries are of type I(b), or more special, with the WAND aligned along the optically privileged null direction \mathbf{k} .
- We were able to explicitly derive the necessary and sufficient conditions of all principal alignment types such that the optically privileged null direction \mathbf{k} is a multiple WAND. These algebraically special (sub)types are II, II(a), II(b), II(c), II(d), their combinations, III, III(a), III(b), N and O. See the explicit conditions given in section 4 and table 3.
- In section 5 we also identified the secondary alignment types for which there exists an additional specific WAND \mathbf{l} distinct from the (multiple) WAND \mathbf{k} , namely I_i , II_i , III_i or D with a double \mathbf{l} . Moreover, there are various subtypes, namely $II(a)_i$, $II(b)_i$, $II(c)_i$, $II(d)_i$ (or their combinations), $III(a)_i$, $III(b)_i$ and $D(a)$, $D(b)$, $D(c)$, $D(d)$.
- The Kundt family, which is the non-expanding ($\Theta = 0$) subclass of the non-twisting shear-free geometries, is studied in section 6. The corresponding conditions for algebraic types and subtypes are simplified, and they fully agree with those obtained previously in [13].
- The algebraic structure of the general Robinson–Trautman class with an arbitrary expansion scalar $\Theta \neq 0$ is described in section 4. The special case $\Theta = 1/r$ is investigated in section 7. In fact, this is an important subcase, as such Robinson–Trautman geometries are of Riemann type I and also of Ricci type I.
- No field equations have been employed in these calculations and discussions. All results are thus ‘purely geometrical’, i.e., they can be applied in any metric theory of gravity that admits non-twisting and shear-free geometries.
- Of course, there are specific constraints on admissible algebraic types imposed by the field equations. To illustrate this, in section 8 we investigated an important example, namely the Robinson–Trautman vacuum solutions in Einstein’s theory. We proved that in all dimensions higher than four there exist only types $D(a) \equiv D(abd)$, $D(c) \equiv D(bcd)$ and O of such spacetimes. This is in striking contrast to the classical $D = 4$ case, which is much richer, see table 4.

Acknowledgements

JP has been supported by the grant GAČR P203/12/0118, RŠ by the project UNCE 204020/2012 and the Czech–Austrian MOBILITY grant 7AMB13AT003.

Appendix A. Explicit curvature tensors for a general non-twisting and shear-free geometry

After applying the shear-free condition (5) the Christoffel symbols for the general non-twisting geometry (1) are

$$\Gamma_{rr}^r = 0, \quad (\text{A.1})$$

$$\Gamma_{ru}^r = -\frac{1}{2}g_{uu,r} + \frac{1}{2}g^{rn}g_{un,r}, \quad (\text{A.2})$$

$$\Gamma_{rp}^r = -\frac{1}{2}g_{up,r} + \Theta g_{up}, \quad (\text{A.3})$$

$$\Gamma_{uu}^r = \frac{1}{2} \left[-g^{rr}g_{uu,r} - g_{uu,u} + g^{rn} \left(2g_{un,u} - g_{uu,n} \right) \right], \quad (\text{A.4})$$

$$\Gamma_{up}^r = \frac{1}{2} \left[-g^{rr}g_{up,r} - g_{uu,p} + g^{rn} \left(2g_{u[n,p]} + g_{np,u} \right) \right], \quad (\text{A.5})$$

$$\Gamma_{pq}^r = -\Theta g^{rr}g_{pq} - g_{u(p||q)} + \frac{1}{2}g_{pq,u}, \quad (\text{A.6})$$

$$\Gamma_{rr}^u = \Gamma_{ru}^u = \Gamma_{rp}^u = 0, \quad (\text{A.7})$$

$$\Gamma_{uu}^u = \frac{1}{2}g_{uu,r}, \quad (\text{A.8})$$

$$\Gamma_{up}^u = \frac{1}{2}g_{up,r}, \quad (\text{A.9})$$

$$\Gamma_{pq}^u = \Theta g_{pq}, \quad (\text{A.10})$$

$$\Gamma_{rr}^m = 0, \quad (\text{A.11})$$

$$\Gamma_{ru}^m = \frac{1}{2}g^{mn}g_{un,r}, \quad (\text{A.12})$$

$$\Gamma_{rp}^m = \Theta \delta_p^m, \quad (\text{A.13})$$

$$\Gamma_{uu}^m = \frac{1}{2} \left[-g^{rm}g_{uu,r} + g^{mn} \left(2g_{un,u} - g_{uu,n} \right) \right], \quad (\text{A.14})$$

$$\Gamma_{up}^m = \frac{1}{2} \left[-g^{rm}g_{up,r} + g^{mn} \left(2g_{u[n,p]} + g_{np,u} \right) \right], \quad (\text{A.15})$$

$$\Gamma_{pq}^m = -\Theta g^{rm}g_{pq} + {}^S\Gamma_{pq}^m. \quad (\text{A.16})$$

The Riemann curvature tensor components then read

$$R_{rprq} = -(\Theta_{,r} + \Theta^2)g_{pq}, \quad (\text{A.17})$$

$$R_{rpru} = -\frac{1}{2}g_{up,rr} + \frac{1}{2}\Theta g_{up,r}, \quad (\text{A.18})$$

$$R_{rpmq} = 2g_{p[m}\Theta_{,q]} - 2\Theta^2 g_{p[m}g_{q]u} + \Theta g_{p[m}g_{q]u,r}, \quad (\text{A.19})$$

$$R_{ruru} = -\frac{1}{2}g_{uu,rr} + \frac{1}{4}g^{mn}g_{um,r}g_{un,r}, \quad (\text{A.20})$$

$$R_{rpmq} = \frac{1}{2}g_{up,r||q} + \frac{1}{4}g_{up,r}g_{uq,r} - g_{pq}\Theta_{,u} - \frac{1}{2}\Theta \left(g_{pq,u} + g_{pq}g_{uu,r} + g_{uq}g_{up,r} - g_{pq}g^{rn}g_{un,r} + 2g_{u[p,q]} \right), \quad (\text{A.21})$$

$$R_{rupq} = g_{u[p,q],r} + \Theta \left(g_{u[p}g_{q]u,r} - 2g_{u[p,q]} \right), \quad (\text{A.22})$$

$$R_{mpnq} = {}^S R_{mpnq} - \Theta^2 g^{rr} \left(g_{mn}g_{pq} - g_{mq}g_{pn} \right) - \Theta \left(g_{mn}e_{pq} + g_{pq}e_{mn} - g_{mq}e_{pn} - g_{pn}e_{mq} \right), \quad (\text{A.23})$$

$$R_{ruup} = \frac{1}{2} \left(g_{uu,rp} - g_{up,ru} \right) + \frac{1}{4}g^{rn}g_{un,r}g_{up,r} - \frac{1}{2}g^{mn}g_{um,r}E_{np} + \Theta \left(g_{up,u} - \frac{1}{2}g_{uu,p} - \frac{1}{2}g_{up}g_{uu,r} \right), \quad (\text{A.24})$$

$$R_{upmq} = g_{p[m,u]||q} + g_{u[q,m]||p} + e_{p[m}g_{q]u,r} + \Theta \left(g^{rr}g_{p[m}g_{q]u,r} + g_{uu,[q}g_{m]p} - 2g^{rn}E_{n[q}g_{m]p} \right), \quad (\text{A.25})$$

$$R_{upuq} = -\frac{1}{2} \left(g_{uu} \right)_{||p||q} + g_{u(p,u]||q} - \frac{1}{2}g_{pq,uu} + \frac{1}{4}g^{rr}g_{up,r}g_{uq,r} - \frac{1}{2}g_{uu,r}e_{pq} + \frac{1}{2}g_{uu,(p}g_{q)u,r} - g^{rn}E_{n(p}g_{q)u,r} + g^{mn}E_{mp}E_{nq} - \frac{1}{2}\Theta g_{pq} \left[g^{rr}g_{uu,r} + g_{uu,u} - g^{rn} \left(2g_{un,u} - g_{uu,n} \right) \right]. \quad (\text{A.26})$$

The components of the Ricci tensor are

$$R_{rr} = -(D-2)(\Theta_{,r} + \Theta^2), \quad (\text{A.27})$$

$$R_{rp} = -\frac{1}{2}g_{up,rr} + g_{up}\Theta_{,r} - (D-3)\Theta_{,p} + (D-2)\Theta^2 g_{up} - \frac{1}{2}(D-4)\Theta g_{up,r}, \quad (\text{A.28})$$

$$R_{ru} = -\frac{1}{2}g_{uu,rr} + \frac{1}{2}g^{rn}g_{un,rr} + \frac{1}{2}g^{mn} \left(g_{um,r||n} + g_{un,r}g_{um,r} \right) - (D-2)\Theta_{,u} - \frac{1}{2}\Theta \left[g^{mn}g_{mn,u} - (D-4)g^{rn}g_{un,r} + (D-2)g_{uu,r} \right], \quad (\text{A.29})$$

$$\begin{aligned}
R_{pq} = & {}^S R_{pq} - f_{pq} - g_{pq} \left(g^{rr} \Theta_{,r} - 2\Theta_{,u} + 2g^{rn} \Theta_{,n} \right) + 2g_{u(p} \Theta_{,q)} \\
& + \Theta^2 \left[2g_{pq} g^{rn} g_{un} - (D-2) g_{pq} g^{rr} - 2g_{up} g_{uq} \right] \\
& + \Theta \left[2g_{u(p} g_{q)} + 2g_{u(p} g_{q)u,r} - (D-2) e_{pq} \right. \\
& \left. + g_{pq} \left(g_{uu,r} - 2g^{rn} g_{un,r} - g^{mn} e_{mn} \right) \right], \tag{A.30}
\end{aligned}$$

$$\begin{aligned}
R_{up} = & -\frac{1}{2} g^{rr} g_{up,rr} - \frac{1}{2} g_{uu,rp} + \frac{1}{2} g_{up,ru} + g^{rn} g_{u[n,p],r} - \frac{1}{2} g^{rn} \left(g_{up,r} g_{un} + g_{un,r} g_{up,r} \right) \\
& + g^{mn} \left(\frac{1}{2} g_{um,r} g_{un} g_{up} + g_{m[p,u]n} + g_{u[m,p]n} - \frac{1}{2} e_{mn} g_{up,r} \right) + g_{up} \Theta_{,u} \\
& + \Theta \left[g_{up} g_{uu,r} + \frac{1}{2} (D-4) \left(g_{uu} g_{up,r} - g_{uu,p} \right) - g_{up,u} - g^{rn} g_{un,r} g_{up} \right. \\
& \left. + (D-6) g^{rn} \left(g_{u[n,p]} - \frac{1}{2} g_{un} g_{up,r} \right) + \frac{1}{2} (D-2) g^{rn} g_{np,u} \right], \tag{A.31}
\end{aligned}$$

$$\begin{aligned}
R_{uu} = & -\frac{1}{2} g^{rr} g_{uu,rr} - g^{rn} g_{uu,rm} - \frac{1}{2} g^{mn} e_{mn} g_{uu,r} + g^{rn} g_{un,ru} - \frac{1}{2} g^{mn} g_{mn,uu} \\
& + g^{mn} \left(g_{um,u} g_{un} - \frac{1}{2} g_{uu} g_{m} g_{n} \right) + \frac{1}{2} \left(g^{rr} g^{mn} - g^{rm} g^{rn} \right) g_{um,r} g_{un,r} \\
& + 2g^{mn} g^{rp} g_{um,r} g_{un,p} + \frac{1}{2} g^{mn} g_{um,r} g_{un,n} + g^{mn} g^{pq} E_{pm} E_{qn} \\
& + \frac{1}{2} \Theta \left[(D-4) g^{rn} \left(2g_{un,u} - g_{uu,n} - g_{un} g_{uu,r} \right) \right. \\
& \left. + (D-2) \left(g_{uu} g_{uu,r} - g_{uu,u} \right) \right], \tag{A.32}
\end{aligned}$$

and the Ricci scalar is

$$\begin{aligned}
R = & {}^S R + g_{uu,rr} - 2g^{rn} g_{un,rr} - 2g^{mn} g_{um,r} g_{un} - \frac{3}{2} g^{mn} g_{um,r} g_{un,r} \\
& + 2\Theta_{,r} \left[(D-2) g_{uu} - (D-3) g^{rn} g_{un} \right] + 4(D-2) \Theta_{,u} - 4(D-3) g^{rn} \Theta_{,n} \\
& - \Theta^2 \left[(D-1)(D-2) g^{rr} - 2(2D-5) g^{rn} g_{un} \right] \\
& + \Theta \left[2(D-2) g_{uu,r} - 2(2D-7) g^{rn} g_{un,r} \right. \\
& \left. + (D-1) g^{mn} g_{mn,u} - 2(D-3) g^{mn} g_{um} g_{un} \right]. \tag{A.33}
\end{aligned}$$

These expressions enable us to calculate the explicit components of the Weyl tensor for any non-twisting and shear-free geometry of an arbitrary dimension D . After a straightforward but very lengthy calculation we obtain

$$C_{rprq} = 0, \tag{A.34}$$

$$C_{rpru} = \frac{D-3}{D-2} \left[-\frac{1}{2} g_{up,rr} + g_{up} \Theta_{,r} + \Theta_{,p} + \Theta g_{up,r} \right], \tag{A.35}$$

$$C_{rpmq} = \frac{2}{D-2} \left[-\frac{1}{2} g_{p[m} g_{q]u,rr} + g_{p[m} g_{q]u} \Theta_{,r} + g_{p[m} \Theta_{,q]} + \Theta g_{p[m} g_{q]u,r} \right], \quad (\text{A.36})$$

$$\begin{aligned} C_{ruru} = & -\frac{D-3}{D-1} \left[\left(\frac{1}{2} g_{uu,r} - \Theta g_{uu} \right)_{,r} + \frac{1}{(D-2)(D-3)} {}^S R \right. \\ & - \frac{1}{4} \frac{D-4}{D-2} g^{mn} g_{um,r} g_{un,r} + \frac{1}{D-2} \left(g^{rn} g_{un,rr} + g^{mn} g_{um,r||n} \right) \\ & - \frac{2}{D-2} g^{rn} g_{un} \Theta_{,r} - 2\Theta_{,u} - \frac{4}{D-2} g^{rn} \Theta_{,n} \\ & \left. - \Theta^2 \frac{D-4}{D-2} g^{rn} g_{un} + \Theta \left(\frac{D-6}{D-2} g^{rn} g_{un,r} - \frac{2}{D-2} g^{mn} g_{um||n} \right) \right], \quad (\text{A.37}) \end{aligned}$$

$$\begin{aligned} C_{rpq} = & \frac{1}{D-2} \left[{}^S R_{pq} - \frac{1}{D-1} g_{pq} {}^S R + \frac{1}{2} (D-2) g_{u[p,r||q]} + \frac{1}{2} (D-4) f_{pq} \right. \\ & + \frac{1}{2} \frac{D-3}{D-1} g_{pq} g_{uu,rr} - \frac{1}{2} g_{up} g_{uq,rr} - \frac{1}{2} \frac{D-5}{D-1} g_{pq} g^{rn} g_{um,rr} \\ & - \frac{1}{2} \frac{D-4}{D-1} g_{pq} g^{mn} g_{um,r} g_{un,r} - \frac{1}{2} \frac{D-5}{D-1} g_{pq} g^{mn} g_{um,r||n} \\ & + \Theta_{,r} \left(g_{up} g_{uq} - \frac{D-3}{D-1} g_{pq} g_{uu} + \frac{D-5}{D-1} g_{pq} g^{rn} g_{un} \right) - 2 \frac{D-3}{D-1} g_{pq} \Theta_{,u} \\ & + 2 \frac{D-5}{D-1} g_{pq} g^{rn} \Theta_{,n} - (D-5) g_{u(p} \Theta_{,q)} - (D-3) g_{u[p} \Theta_{,q]} \\ & + \Theta^2 (D-4) \left(g_{up} g_{uq} - \frac{2}{D-1} g_{pq} g^{rn} g_{un} \right) \\ & + \Theta \left(\frac{3D-13}{D-1} g_{pq} g^{rn} g_{un,r} - \frac{D-3}{D-1} g_{pq} g_{uu,r} - (D-5) g_{u(p} g_{q)u,r} + g_{u[p} g_{q]u,r} \right. \\ & \left. - (D-4) g_{u(p||q)} - (D-2) g_{u[p,q]} + \frac{D-5}{D-1} g_{pq} g^{mn} g_{um||n} \right) \Big], \quad (\text{A.38}) \end{aligned}$$

$$\begin{aligned} C_{rupq} = & g_{u[p,q],r} - \frac{1}{D-2} g_{u[p} g_{q]u,rr} \\ & - 2 \frac{D-3}{D-2} g_{u[p} \Theta_{,q]} - 2\Theta \left(g_{u[p,q]} - \frac{1}{D-2} g_{u[p} g_{q]u,r} \right), \quad (\text{A.39}) \end{aligned}$$

$$\begin{aligned} C_{mpnq} = & {}^S C_{mpnq} + \frac{2}{(D-2)(D-4)} \left(g_{mn} {}^S R_{pq} + g_{pq} {}^S R_{mn} - g_{mq} {}^S R_{pn} - g_{pn} {}^S R_{mq} \right) \\ & + \frac{1}{D-2} \left(g_{mn} f_{pq} + g_{pq} f_{mn} - g_{mq} f_{pn} - g_{pn} f_{mq} \right) \\ & - \frac{2}{D-2} \left[g_{mn} \left(g_{u(p} \Theta_{,q)} - \Theta^2 g_{up} g_{uq} + \Theta \left(g_{u(p||q)} + g_{u(p} g_{q)u,r} \right) \right) \right. \\ & \left. + g_{pq} \left(g_{u(m} \Theta_{,n)} - \Theta^2 g_{um} g_{un} + \Theta \left(g_{u(m||n)} + g_{u(m} g_{n)u,r} \right) \right) \right] \end{aligned}$$

$$\begin{aligned}
& -g_{mq} \left(g_{u(p}\theta_{,n)} - \theta^2 g_{up} g_{un} + \theta \left(g_{u(p||n)} + g_{u(p} g_{n)u,r} \right) \right) \\
& -g_{pn} \left(g_{u(m}\theta_{,q)} - \theta^2 g_{um} g_{uq} + \theta \left(g_{u(m||q)} + g_{u(m} g_{q)u,r} \right) \right) \\
& + \frac{1}{(D-1)(D-2)} (g_{mn} g_{pq} - g_{mq} g_{pn}) \left[g_{uu,rr} - 2g^{rs} g_{us,rr} \right. \\
& - 2g^{os} f_{os} - \frac{1}{2} g^{os} g_{uo,r} g_{us,r} - \frac{2(2D-5)}{(D-3)(D-4)} s_R \\
& + 2\theta_{,r} (-g_{uu} + 2g^{rs} g_{us}) - 4\theta_{,u} + 8g^{rs} \theta_{,s} \\
& \left. - 6\theta^2 g^{rs} g_{us} + 2\theta (-g_{uu,r} + 5g^{rs} g_{us,r} + 2g^{os} g_{uo||s}) \right], \tag{A.40}
\end{aligned}$$

$$\begin{aligned}
C_{ruup} = & \frac{1}{2} \frac{D-3}{D-2} (g_{uu,rp} - g_{up,ru}) + \frac{1}{4} \frac{D-4}{D-2} g^m g_{un,r} g_{up,r} - \frac{1}{2} \frac{D-3}{D-2} g^{mn} g_{um,r} E_{np} \\
& + \frac{1}{D-2} \left[g^{mn} e_{m[p} g_{n]u,r} + g^{mn} (g_{m[p,u||n]} + g_{u[m,p]||n}) \right] \\
& - \frac{1}{D-2} \left[\frac{1}{2} g^m g_{un} g_{up,rr} - g^m g_{u[n,p],r} + \frac{1}{2} g^m g_{up,r||n} \right] \\
& - \frac{1}{(D-1)(D-2)} g_{up} \left[s_R - \frac{1}{2} (D-3) g_{uu,rr} + \frac{1}{2} (D-4) g^{mn} g_{um,r} g_{un,r} \right. \\
& \left. + \frac{1}{2} (D-5) (g^m g_{un,rr} + g^{mn} g_{um,r||n}) \right] \\
& + \frac{D-3}{(D-1)(D-2)} \left[g_{up} (2g^m g_{un} - g_{uu}) \theta_{,r} + (D-3) g_{up} \theta_{,u} \right. \\
& \left. - (D-1) g_{uu} \theta_{,p} + 4g_{up} g^m \theta_{,n} \right] + \theta^2 \frac{(D-3)(D-4)}{(D-1)(D-2)} g^m g_{un} g_{up} \\
& - \frac{1}{D-2} \theta \left[(D-3) (g_{uu,p} - g_{up,u}) + \frac{1}{2} (D-6) g^m g_{un} g_{up,r} \right. \\
& \left. - \frac{1}{2} (D-2) g^m g_{np,u} - (D-6) g^m g_{u[n,p]} \right] \\
& - \frac{1}{(D-1)(D-2)} g_{up} \theta \left[\frac{1}{2} (D-5)(D-6) g^m g_{un,r} + \frac{1}{2} (D-1) g^{mn} g_{mn,u} \right. \\
& \left. + (D-3) (g_{uu,r} - 2g^{mn} g_{um||n}) \right], \tag{A.41}
\end{aligned}$$

$$\begin{aligned}
C_{upmq} = & g_{p[m,u||q]} + g_{u[q,m]||p} + e_{p[m} g_{q]u,r} + \frac{2}{D-2} (s_R p_{[m} g_{q]u} - f_{p[m} g_{q]u}) \\
& + \frac{1}{D-2} \left[(g_{uu} - g^m g_{un}) g_{p[m} g_{q]u,rr} - g_{uu,r} g_{[q} g_{m]p} + g_{p[m} g_{q]u,ru} \right. \\
& \left. + g^m (g_{pm} g_{u[n,q],r} - g_{pq} g_{u[n,m],r}) - g^m g_{un,r} g_{p[m} g_{q]u,r} - g^m g_{p[m} g_{q]u,r||n} \right]
\end{aligned}$$

$$\begin{aligned}
& + g^{ns} g_{un,r} g_{us} \llbracket [q g_m]_p + g^{ns} \left(g_{pm} g_{n[q,u]s} - g_{pq} g_{n[m,u]s} \right) \\
& + g^{ns} \left(g_{pm} g_{u[n,q]s} - g_{pq} g_{u[n,m]s} \right) - g^{ns} e_{ns} g_{p[m]g_q]u,r} \Big] \\
& - \frac{2}{(D-1)(D-2)} g_{p[m]g_q]u} \left[{}^S R + g_{uu,rr} - 2g^{rn} g_{un,rr} - \frac{3}{2} g^{ns} g_{un,r} g_{us,r} \right. \\
& - 2g^{ns} g_{un,r} \Big] + \left((D-3) g_{uu} - (D-5) g^{rn} g_{un} \right) \Theta_{,r} \\
& + (D-5) \Theta_{,u} - 2(D-5) g^{rn} \Theta_{,n} \Big] \\
& + \frac{2}{D-2} g_{up} \Theta_{,[m]g_q]u} - 4\Theta^2 \frac{D-4}{(D-1)(D-2)} g^{rn} g_{un} g_{p[m]g_q]u} - 2\Theta e_{p[m]g_q]u} \\
& + \frac{2}{D-2} \Theta \left[g_{u(p)g_q]u} - g_{u(p)g_q]u} - g_{up} g_{u[m]g_q]u,r} - g_{p[m]g_q]u,r} (g_{uu} - 2g^{rn} g_{un}) \right. \\
& - 2g^{rn} \left(g_{pm} g_{u[n,q]} - g_{pq} g_{u[n,m]} \right) - g_{p[m]g_q]u,u} + g_{uu,[q]g_m]p} \Big] \\
& + \frac{2}{(D-1)(D-2)} \Theta g_{p[m]g_q]u} \left[2g_{uu,r} + (D-11) g^{rn} g_{un,r} \right. \\
& \left. + (D-5) g^{ns} g_{un} \Big] - \frac{1}{2} (D-1) g^{ns} g_{ns,u} \Big], \tag{A.42}
\end{aligned}$$

$$\begin{aligned}
C_{upuq} = & -\frac{1}{2} g_{uu} \llbracket p \parallel q - \frac{1}{2} g_{pq,uu} + g_{u(p,u)q} - \frac{1}{2} g_{uu,r} e_{pq} + \frac{1}{2} g_{uu,(p]g_q]u,r} + g^{os} E_{op} E_{sq} \\
& - \frac{1}{D-2} g_{pq} g^{mn} \left(-\frac{1}{2} g_{uu} \llbracket m \parallel n - \frac{1}{2} g_{mn,uu} + g_{um,u} \parallel n \right. \\
& \left. - \frac{1}{2} g_{uu,r} e_{mn} + \frac{1}{2} g_{uu,m} g_{un,r} + g^{os} E_{om} E_{sn} \right) \\
& + \frac{1}{(D-1)(D-2)} \left(g_{uu} g_{pq} - g_{up} g_{uq} \right) \left({}^S R + g_{uu,rr} - 2g^{rn} g_{un,rr} \right. \\
& \left. - \frac{3}{2} g^{mn} g_{um,r} g_{un,r} - 2g^{mn} g_{um,r} \parallel n \right) \\
& - \frac{1}{2(D-2)} g_{uu} g_{pq} \left(g_{uu,rr} - g^{mn} g_{um,r} g_{un,r} \right) - \frac{1}{D-2} g_{uu} {}^S R_{pq} \\
& - \frac{1}{4} \left(\frac{D-4}{D-2} g_{uu} - g^{rn} g_{un} \right) g_{up,r} g_{uq,r} + \frac{1}{D-2} g_{uu} g_{u(p,r)q} - g^{rn} E_{n(p]g_q]u,r} \\
& + \frac{1}{D-2} g_{pq} g^{rn} \left[\frac{1}{2} g_{un} g_{uu,rr} + g_{uu,rm} - g_{un,ru} \right. \\
& \left. - \frac{1}{2} g_{uo,r} \left(g^{os} g_{us,r} g_{un} - g^{ro} g_{un,r} - 4 g^{os} g_{u[n,s]} \right) \right] \\
& + \frac{1}{D-2} \left[\left(g_{uu} - g^{rn} g_{un} \right) g_{u(q]g_p]u,rr} - g_{uu,r(p]g_q]u} + g_{u(q]g_p]u,ru} \right.
\end{aligned}$$

$$\begin{aligned}
& + g^m g_{un,r||p} g_{q} u - 2g^m g_{u(q} g_{p)u,r||n} - g^m g_{un,r} g_{u(q} g_{p)u,r} \Big] \\
& + \frac{1}{D-2} g^{mn} \left[g_{um,r} g_{u||p} g_{q} u - e_{mn} g_{u(q} g_{p)u,r} + g_{u(q} g_{p)m,u||n} - g_{mn,u||p} g_{q} u \right. \\
& + \left. \frac{1}{2} (g_{uq} g_{um||p||n} + g_{up} g_{um||q||n}) - g_{u(q} g_{p)u||m||n} \right] \\
& + \frac{1}{(D-1)(D-2)} \Theta_{,r} \left[g_{uu} g_{pq} ((D-3)g_{uu} - (D-5)g^m g_{un}) \right. \\
& - \left. 2g_{up} g_{uq} ((D-2)g_{uu} - (D-3)g^m g_{un}) \right] \\
& + \Theta_{,u} \frac{2(D-3)}{(D-1)(D-2)} (g_{uu} g_{pq} - g_{up} g_{uq}) - \frac{2}{D-2} g_{uu} g_{u(p} \Theta_{,q)} \\
& - 2g^m \Theta_{,n} \left(\frac{D-5}{(D-1)(D-2)} g_{uu} g_{pq} - \frac{2(D-3)}{(D-1)(D-2)} g_{up} g_{uq} \right) \\
& + \Theta^2 \left[\frac{2(D-4)}{(D-1)(D-2)} g^m g_{un} g_{uu} g_{pq} - \frac{D-4}{D-2} g_{uu} g_{up} g_{uq} + \frac{(D-3)(D-4)}{(D-1)(D-2)} g^m g_{un} g_{up} g_{uq} \right] \\
& + \Theta \left[\frac{1}{D-2} g_{pq} g^{rn} (2g_{un,u} - g_{uu,n} - g_{un} g_{uu,r}) + \frac{D-6}{D-2} (g_{uu} - g^m g_{un}) g_{u(q} g_{p)u,r} \right. \\
& + \frac{D-3}{(D-1)(D-2)} g_{uu,r} g_{uu} g_{pq} + \frac{2}{(D-1)(D-2)} g_{up} g_{uq} g_{uu,r} \\
& - \left. \frac{2(D-6)}{(D-1)(D-2)} g^m g_{un,r} (g_{uu} g_{pq} - g_{up} g_{uq}) \right. \\
& + g_{uu} e_{pq} + 2g^m E_{n(p} g_{q)u} - \frac{1}{D-2} g_{uu} (2g_{u(p||q)} - g_{pq} g^{mn} e_{mn}) \\
& - \frac{2}{D-2} g_{u(q} g_{p)u,u} - \frac{D-4}{D-2} g_{uu,(p} g_{q)u} - \frac{4}{D-2} g^m (g_{u||p} g_{q} u - g_{u(q} g_{p)u||n}) \\
& \left. + \frac{1}{(D-1)(D-2)} (g_{uu} g_{pq} - g_{up} g_{uq}) g^{mn} ((D-1)g_{mn,u} - 2(D-3)g_{um||n}) \right]. \quad (\text{A.43})
\end{aligned}$$

In the above expressions, ${}^S \Gamma_{pq}^m \equiv \frac{1}{2} g^{mn} (2g_{n(p,q)} - g_{pq,n})$ denote Christoffel symbols with respect to the spatial coordinates only, i.e., the coefficients of the covariant derivative on the transverse $(D-2)$ -dimensional Riemannian space. The symbol $_{||}$ denotes this covariant derivative with respect to g_{pq} . Similarly, ${}^S R_{mpnq}$, ${}^S C_{mpnq}$, ${}^S R_{pq}$ and ${}^S R$ are the Riemann tensor, Weyl tensor, Ricci tensor and Ricci scalar for the transverse-space metric g_{pq} , respectively. We have also introduced the following useful auxiliary quantities :

$$g_{up||q} \equiv g_{up,q} - g_{um} {}^S \Gamma_{pq}^m, \quad (\text{A.44})$$

$$g_{up,r||q} \equiv g_{up,rq} - g_{um,r} {}^S \Gamma_{pq}^m, \quad (\text{A.45})$$

$$g_{u[p,r||q]} = g_{u[p||q],r}, \quad (\text{A.46})$$

$$g_{p[m,u||q]} \equiv g_{p[m,q],u} + \frac{1}{2} ({}^S \Gamma_{pm}^n g_{nq,u} - {}^S \Gamma_{pq}^n g_{nm,u}), \quad (\text{A.47})$$

$$g_{u[q,m]||p} \equiv g_{u[q,m],p} - {}^S\Gamma_{pq}^n g_{u[n,m]} - {}^S\Gamma_{pm}^n g_{u[q,n]}, \quad (\text{A.48})$$

$$(g_{uu})_{||p||q} \equiv g_{uu,pq} - g_{uu,n} {}^S\Gamma_{pq}^n, \quad (\text{A.49})$$

$$g_{up,u||q} \equiv g_{up,uq} - g_{um,u} {}^S\Gamma_{pq}^m, \quad (\text{A.50})$$

$$e_{pq} \equiv g_{u(p||q)} - \frac{1}{2}g_{pq,u}, \quad (\text{A.51})$$

$$E_{pq} \equiv g_{u[p,q]} + \frac{1}{2}g_{pq,u}, \quad (\text{A.52})$$

$$f_{pq} \equiv g_{u(p,r||q)} + \frac{1}{2}g_{up,r}g_{uq,r}, \quad (\text{A.53})$$

where $g_{u[p,q]} = g_{u(p||q)}$. These are tensors on the transverse Riemannian space.

References

- [1] Robinson I and Trautman A 1960 Spherical gravitational waves *Phys. Rev. Lett.* **4** 431–2
- [2] Robinson I and Trautman A 1962 Some spherical gravitational waves in general relativity *Proc. R. Soc. A* **265** 463–73
- [3] Kundt W 1961 The plane-fronted gravitational waves *Z. Phys.* **163** 77–86
- [4] Kundt W 1962 Exact solutions of the field equations: twist-free pure radiation fields *Proc. R. Soc. A* **270** 328–34
- [5] Stephani H, Kramer D, MacCallum M, Hoenselaers C and Herlt E 2003 *Exact Solutions of Einstein's Field Equations* (Cambridge: Cambridge University Press)
- [6] Griffiths J B and Podolský J 2009 *Exact Space-Times in Einstein's General Relativity* (Cambridge: Cambridge University Press)
- [7] Podolský J and Ortaggio M 2006 Robinson–Trautman spacetimes in higher dimensions *Class. Quantum Grav.* **23** 5785–97
- [8] Ortaggio M, Podolský J and Žofka M 2008 Robinson–Trautman spacetimes with an electromagnetic field in higher dimensions *Class. Quantum Grav.* **25** 025006
- [9] Švarc R and Podolský J 2014 Absence of gyratons in the Robinson–Trautman class *Phys. Rev. D* **89** 124029
- [10] Podolský J and Žofka M 2009 General Kundt spacetimes in higher dimensions *Class. Quantum Grav.* **26** 105008
- [11] Coley A, Hervik S, Papadopoulos G and Pelavas N 2009 Kundt spacetimes *Class. Quantum Grav.* **26** 105016
- [12] Ortaggio M, Pravda V and Pravdová A 2013 Algebraic classification of higher dimensional spacetimes based on null alignment *Class. Quantum Grav.* **30** 013001
- [13] Podolský J and Švarc R 2013 Explicit algebraic classification of Kundt geometries in any dimension *Class. Quantum Grav.* **30** 125007
- [14] Coley A and Hervik S 2010 Higher dimensional bivectors and classification of the Weyl operator *Class. Quantum Grav.* **27** 015002
- [15] Coley A, Hervik S, Ortaggio M and Wylleman L 2012 Refinements of the Weyl tensor classification in five dimensions *Class. Quantum Grav.* **29** 155016
- [16] Ortaggio M, Pravda V and Pravdová A 2013 On the Goldberg–Sachs theorem in higher dimensions in the non-twisting case *Class. Quantum Grav.* **30** 075016
- [17] Hervik S, Ortaggio M and Wylleman L 2013 Minimal tensors and purely electric or magnetic spacetimes of arbitrary dimension *Class. Quantum Grav.* **30** 165014
- [18] Podolský J and Švarc R 2013 Physical interpretation of Kundt spacetimes using geodesic deviation *Class. Quantum Grav.* **30** 205016
- [19] Podolský J and Švarc R 2012 Interpreting spacetimes of any dimension using geodesic deviation *Phys. Rev. D* **85** 044057

- [20] Coley A, Milson R, Pravda V and Pravdová A 2004 Classification of the Weyl tensor in higher dimensions *Class. Quantum Grav.* **21** L35–41
- [21] Wald R M 1984 *General Relativity* (Chicago: University of Chicago Press)