

The advertisement role of major urinary proteins in mice

Pavel Stopka^{a,b,*}, Katerina Janotova^a, David Heyrovsky^c

^a Department of Zoology, Charles University, Vinicna 7, 128 44 Praha 2, Czech Republic

^b Institute of Animal Physiology and Genetics, Academy of Sciences of the Czech Republic, Rumburska 89, 277 21 Libechov, Czech Republic

^c Institute of Theoretical Physics, Charles University, V Holesovickach 2, 180 00 Praha 8, Czech Republic

Received 8 December 2006; received in revised form 27 March 2007; accepted 29 March 2007

Abstract

The variation of expression of major urinary proteins was studied in laboratory mice to further the understanding of the role of these proteins in social and reproductive contexts. Mouse major urinary proteins (MUPs) are known to carry volatile substances and protect them during their passage from the liver, through the kidneys into the urine. However, most studies on the role of MUPs were carried out on males. Using densitometry analysis of total MUP concentration in the urine, our present study clearly demonstrates that (i) individuals of both sex up-regulate MUPs during social contact, and that (ii) females use these proteins to advertise their reproductive state by varying the concentration of MUPs during the oestrous cycle. As the concentration of MUPs was normalized by the concentration of creatinine – a marker of glomerular filtration – the corrected concentration of MUPs represents instantaneous expression on the level of proteins. Cross-correlation analysis between oestrus quantification and MUP expression revealed that the oestrous curve is delayed by 1 day behind the MUP curve so that the expression of MUPs is up-regulated immediately at the beginning of oestrus. To conclude, the regulation of pheromone-carrying MUPs is directly linked to reproduction and, thus, enables female honest signalling.

© 2007 Elsevier Inc. All rights reserved.

Keywords: Oestrus; Oestrous cycle; Pheromone-carrying proteins; MUP; Major urinary protein; Mouse; Mice

1. Introduction

Pheromones are substances that trigger various physiological responses and are specific for each (signalling and/or receiving) sex. For example, the Whitten effect is a temporal change of female reproductive physiology that is switched on and off in response to the presence or absence of individual males or just their bedding material. This socially dependent mechanism has largely been demonstrated on the level of oestrus induction. Since the original discovery of this phenomenon [1] various authors have provided evidence that there exist specific oestrus-inducing pheromones and that these pheromones may have a different effect upon individuals of the same or the opposite sex. Female pheromones are involved in the suppression of oestrus and prolongation of the oestrous cycle is a typical response of females that are housed together [2,3]. This Lee-Boot effect has

been described in various species of mice of the genera *Mus* and *Apodemus* [4–6]. However, there is scarce scientific evidence on what female pheromones trigger in males and how.

The Whitten and Lee-Boot effects may also be viewed as two aspects of one phenomenon. Stopka and Macdonald [5] presented evidence that both effects can be demonstrated in one species and one experiment with different social setups or in varying social contexts. For example, female wood mice (*Apodemus sylvaticus*) shorten their oestrous cycle and prolong oestrus when housed with males (separated by a metal grid), but prolong the cycle and shorten the oestrus when caged alone and even more so when caged with other females. According to their study social modulation of the oestrous cycle has further implications regarding mating systems. In this promiscuous species [7] daily alternations of unfamiliar males behind the grid stimulated an accelerated onset of oestrus that was even more prolonged while the oestrous cycle was significantly shortened, so much that oestrus seemed almost continuous. Therefore, these physiological changes are dynamic responses to specific social environments, which vary across species with different

* Corresponding author. Department of Zoology, Charles University, Vinicna 7, 128 44 Praha 2, Czech Republic. Tel.: +420 221951856.

E-mail address: pstopka@natur.cuni.cz (P. Stopka).

mating systems and, thus, are clearly tuned to optimise individual fitness benefits across varying social environments.

Potential benefits of being informed and informing are classically viewed in terms of reproductive benefits, where transmitted information on sex, relatedness, reproductive and social status is an important prerequisite for mate choice. However, many studies on social modulation show that the same information that is used in the process of mate choice stimulates responses with extensive physiological and reproductive consequences. For example, female puberty may be accelerated in sub-adult mice simply in response to the presence of adult males [8] whilst the presence of unfamiliar males may cause spontaneous abortion of early embryos [9] in female mice. All the above described cases of social modulation, including the Whitten, Lee-Boot, Bruce, and Vandenbergh effects, are important physiological phenomena where social organization, mate competition and mate choice have presumably acted as important selective pressures that shaped the evolution of specific signalling and physiological responses.

Original research on socially modulating and modulated effects has been guided by a new paradigm following the discovery of the function of major urinary proteins and their role in transporting pheromones from the liver through the kidney into the urine [10,11]. After being excreted, MUPs release pheromones while drying up and, thus, individual odours are manifested even when the individual is temporarily absent from the territory [12]. Milestones in the understanding of MUPs in the social context include: the discovery of binding sites for pheromones that have specific effects upon targeted individuals [13,14], individually specific and behaviourally relevant expression patterns [15], tissue-specific expression and, therefore, their different function [16,17]. Furthermore, MUPs are an essential part of the odour delivery system that enhances the onset of ovulation in female mice [18]. To complement Morè's [18] finding that male MUPs are important in ovulation induction, this paper highlights the role of female MUPs in oestrus advertisement. We present evidence that during the oestrous cycle female MUPs undergo up-regulation of MUP expression which is tightly bound to the beginning of the oestrous cycle and, therefore, plays an important role in honest reproductive signalling.

2. Methods

2.1. Animals

We used strain C57BL/6 mice as subjects for this experiment. Twenty females and twenty males aged 7 weeks were kept on a 12:12-h light cycle with lights off at 1900 h. Water and food was provided ad libitum. All mice were caged individually in compartmented cages during individual treatment; in social treatment two individuals were separated by a metal grid to allow communication but suppress any behavioural interactions. Urine samples and vaginal smears were collected every day in the first half of the light period. Urine was obtained by gentle abdominal massage and frozen immediately at -20°C . The experiment lasted 17 days. Samples for cross-correlation analysis were collected from all females throughout the duration

of individual treatment. Of these 17 days day 10 was taken as a reference point for individual treatment. The first day of the 4-day social treatment was taken as a reference for the analysis of expression variation due to social treatment.

2.2. Oestrus identification

Vaginal smears were taken by vaginal lavage with 10 μl of Dulbecco's phosphate-buffered saline (Sigma-Aldrich, St. Louis, USA) and put into an Eppendorf tube containing 90 μl of PBS. The whole volume was pipetted into a plastic chamber, placed in a Cytospin 2 slide centrifuge (Shandon Southern Instruments, Sewickley, USA) and spun down (1000 rpm, 10 min) onto a microscope slide. The microscope preparations were stained by the Pappenheim method (May-Grünwald, Giemsa) and analysed under a light microscope.

Different oestrous stages were determined using a modified method of Natynczuk [19]. Randomly selected fields of fixed cells were counted to a total number of 200 cells per slide. The cell types counted were leukocytes (L), cornified non-nucleated epithelial cells (C) and nucleated epithelial cells (N). The index $E = C / (C + N + L)$ was calculated to obtain a continuous and smooth oestrus curve. Females were categorized as: oestrous if $E > 0.7$ or dioestrous if $E < 0.3$.

2.3. Protein analysis

Major urinary proteins from urine samples were separated on SDS-PAGE according to a modified Laemmli [20] protocol. Samples were thawed, stirred and centrifuged for 5 min at 16,000 g. The volume of 0.4 μl of urine was loaded onto a 15% Tris–Gly gel. To allow quantification of MUPs, four different concentrations of standard (Carbonic Anhydrase, 29 kDa, marker for SDS-PAGE, Sigma-Aldrich, St. Louis, USA) were run together with samples on each gel and used for the calculation of the regression curve. The gels were stained with Coomassie® G-250 (SimplyBlue™ Safe Stain, Invitrogen life technologies, Paisley, UK). Gel images were acquired using GS-800 Calibrated Densitometer and analysed with Quantity One 1-D Analysis Software (Bio-Rad Laboratories, Hercules, USA). Bands containing major urinary proteins were detected by western blotting with commercially available antibody YNGMMUP (Accurate chemical and Scientific Corporation, New York, USA). The antibody was tested against pooled normal mouse urine and purified mouse MUP and gives a reaction of full identity.

As the dilution of the urine can vary across different individuals or between days, we employed the creatinine assay to normalize all concentrations in the samples by using a LKreatinin kit (BioVendor, Brno, CZ). Creatinine is a reliable indicator of renal activity showing the volume of liquid filtered through the kidney. Based on the creatinine concentration, the dilution coefficient (kdil) was calculated using the formula $\text{kdil} = C_{\text{creat}} / C_{\text{ref}}$, where C_{creat} is the creatinine concentration of each sample and C_{ref} is the reference creatinine concentration. The reference concentration was defined as the highest concentration of all measured creatinine concentrations. Corrected volumes were obtained by division of




Fig. 1. Normalized expression of MUPs is sex-specific: Males produce approximately three times more MUPs than females.

each MUP volume with the dilution coefficient. These new normalized values, thus, represented the instantaneous level of expression per each mouse sample, whereas the original values reflect the actual concentration of MUPs in a given sample.

3. Results

3.1. Sex differences

Individuals of both sexes differed in the concentration of urinary MUPs ($t=16.13$, $d.f.=34$, $p<0.0000001$), and as depicted in Fig. 1, they also differed in the creatinine-normalized expression of MUPs $t=10.957$, $d.f.=34$, $p<0.0000001$).

3.2. Social modulation of MUPs

Sensitivity to chemosignals was measured in the compartmented setup, where individual expression of MUPs was tested using MANOVA under two treatments: INDIVIDUAL and SOCIAL. Sex was a factor that explained major differences in both MUP concentration ($F=246.1$, $d.f.=1$, $p<0.0001$) and MUP expression ($F=49.85$, $d.f.=1$, $p<0.0001$). Differences in the concentration ($F=6.7$, $d.f.=1$, $p=0.014$) and expression ($F=4.83$, $d.f.=1$, $p=0.035$) due to treatment were, however, also significant.

The combination of factors sex*treatment did not reveal significant results ($F=0.4$, $d.f.=1$, $p=0.521$), thus suggesting that there is no evidence that individuals of opposite sex would react differently in social setup. Furthermore, all individuals regardless of sex up-regulated MUPs in the social treatment.

3.3. Oestrus advertisement

For the analysis of MUP expression variation a cross-correlation procedure was employed, in which an oestrous time-series was tested against MUP expression cyclic variation. In

Fig. 2 a normalized cross-correlation coefficient close or equal to one represents a perfect match, whereas a poor match will yield a value close to zero. The error bars in Fig. 2 represent confidence intervals set at the level of $p=0.05$ and $n=20$ (females). When these confidence bars do not cross the x axis the cross-correlation analysis yields a significant effect. As depicted in Fig. 2, the maximum of MUP concentration precedes the day of oestrus yielding significant values for day 0 and even more for day +1 (maximum oestrus). Furthermore, our cross-correlation analysis also reveals a significant match with a harmonic period of 5 days — a typical length of oestrous cycle when females are caged individually. However, this is primarily the effect of oestrus periodicity. To conclude, the expression of MUPs is elevated at the beginning of oestrus,




Fig. 2. Cross-correlation analysis of time dependence of expression of MUPs and oestrous cyclicality: plot a presents the cross-correlation of expression of MUPs and oestrous cyclicality, with means and confidence intervals ($\alpha=0.05$, $n=20$) obtained by a statistical analysis of twenty studied females. Plot b presents partial results for one of the females.

reaching maximum expression values just 1 day before the phase during which ovulation occurs.

4. Discussion

The classical view of Whitten's effect focuses on female reproductive physiology being modulated as a consequence of exposure to males [1]. Similarly, the Lee-Boot effect focuses on bidirectional modulation of oestrus and the oestrous cycle between two or more females [3]. Furthermore, studies published to date largely and correctly assume that pheromones have the most important triggering effect on major reproductive events through the vomeronasal system of chemoperception [21,22]. In this study we have provided evidence that females vary both the concentration and the expression of MUPs during oestrous cycle and may, therefore, use this system of communication for the advertisement of their reproductive status. To our surprise, identical results were obtained when either MUP concentration or MUP expression was statistically tested. In this case it means that MUP concentration was a good predictor of MUP expression in females.

From the point of view of the costs and benefits of pheromone communication it is reasonable to assume that under different social contexts individuals of both sexes may need to modulate signalling to avoid substantial costs such as harassment from males and predation, and to influence reproduction of other individuals for essential (individual) fitness benefits. This study presents evidence that female mice use a system of pheromone-carrying and pheromone-protecting MUPs to tune pheromonal communication such that individuals may respond differentially to different social contexts. Previous studies [1], as well as the recent study by Morè [18], centred on the role of male pheromones in ovulation/oestrus induction, female receptivity, and puberty acceleration in both sexes [23]. This paper, however, demonstrates that individuals of either sex are simultaneously signallers and receivers and, therefore, their reproductive physiology is both socially modulating and socially modulated. This is an especially important finding because females of wild house mice (*Mus musculus domesticus* and *Mus musculus musculus*) have much lower expression of MUPs than males [24]. However, scarce information is still available on the role of the female pheromone delivery system in modulating male reproductive physiology.

To conclude, the importance of our results can be seen on two levels. First, individuals of both sexes of C57BL/6 mice up-regulate MUP expression in social contexts, thus supporting the assumption that the expression of MUPs is individually traded-off for reproductive benefits. Second, the expression of MUPs is cyclic, probably due to common steroid regulators as well as the oestrous cycle. Our cross-correlation analysis revealed a perfect match when the oestrous curve and MUP expression curve is shifted by 1 day. This demonstrates that oestrus is advertised by female mice prior to its onset in order to attract competing males and, thus, to increase individual fitness benefits.

Acknowledgments

We are very grateful to Jarmila Juklova and Ludmila Vitamvasova for lab assistance and to Petra Andrlíkova, Martin

Sandera, and Romana Stopkova for helpful discussions. The project was funded by: GACR 206/04/20493, MSMT VZ 0021620828, and by IRP IAPG No. A VOZ 50450515.

References

- [1] Whitten WK. Modification of the oestrous cycle of the mouse by external stimuli associated with the male. Changes in the oestrous cycle determined by vaginal smears. *J Endocrinol* 1958;17:307–13.
- [2] Whitten WK. Occurrence of anoestrus in mice caged in groups. *J Endocrinol* 1959;18:102–7.
- [3] Lee S, van der Boot LM. Spontaneous pseudopregnancy in mice. *Acta Physiol Pharmacol Neerl* 1955;4:442–3.
- [4] Cinquetti R, Rinaldi L. Effects of changed social conditions on modulation of the oestrous cycle in mice. *Boll Zool* 1989;56:137–42.
- [5] Stopka P, Macdonald DW. Signal interchange during mating in the wood mouse (*Apodemus sylvaticus*), the concept of active and passive signaling. *Behaviour* 1998;135:231–49.
- [6] Féron C, Gheusi G. Social regulation of reproduction in the female mound-builder mouse (*Mus spicilegus*). *Physiol Behav* 2003;78:717–22.
- [7] Polechova J, Stopka P. Geometry of social relationships in the Old World wood mouse, *Apodemus sylvaticus*. *Can J Zool* 2002;80:1383–8.
- [8] Vandenberg JG. Male odor accelerates female sexual maturation in mice. *Endocrinology* 1969;84:658–60.
- [9] Bruce HM, Parrott DMV. Role of olfactory sense in pregnancy block by strange males. *Science* 1960;131:1526.
- [10] Robertson DHL, Beynon RJ, Evershed RP. Extraction, characterization, and binding analysis of two pheromonally active ligands associated with major urinary protein of house mouse (*Mus musculus*). *J Chem Ecol* 1993;19:1405–16.
- [11] Cavaggioni A, Mucignat-Caretta C. Major urinary proteins, α_{2U} -globulins and aphrodisin. *BBA-Protein Struct M* 2000;1482:218–28.
- [12] Hurst JL, Robertson DHL, Tolladay U, Beynon RJ. Proteins in urine scent marks of male house mice extend the longevity of olfactory signals. *Anim Behav* 1998;55:1289–97.
- [13] Novotny MV, Harvey S, Jemiolo B, Alberts J. Synthetic pheromones that promote inter-male aggression in mice. *Proc Natl Acad Sci USA* 1985;82:2059–61.
- [14] Novotny MV. Pheromones, binding proteins and receptor responses in rodents. *Biochem Soc Trans* 2003;31:117–22.
- [15] Hurst JL, Payne CE, Nevison ChM, Marie AD, Humphries RE, Robertson DHL, et al. Individual recognition in mice mediated by major urinary proteins. *Nature* 2001;414:631–4.
- [16] Shaw PH, Held WA, Hastie ND. The gene family for major urinary proteins, expression in several secretory tissues of the mouse. *Cell* 1983;32:755–61.
- [17] Shahan K, Derman E. Tissue-specific expression of major urinary protein (MUP) genes in mice, characterization of MUP mRNAs by restriction mapping of cDNA and by in vitro translation. *Mol Cell Biol* 1984;4(11):2259–65.
- [18] Morè L. Mouse major urinary proteins trigger ovulation via the vomeronasal organ. *Chem Senses* 2006;31:393–401.
- [19] Natynczuk SE, MacDonald DW. Scent, sex, and self-calibrating rat. *J Chem Ecol* 1994;20:1843–57.
- [20] Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227(259):680–5.
- [21] Shepherd GM. Smells, brains and hormones. *Nature* 2006;439:149–51.
- [22] von Campenhausen H, Mori K. Convergence of segregated pheromonal pathways from the accessory olfactory bulb to the cortex in the mouse. *Eur J Neurosci* 2000;12:33–46.
- [23] Mucignat-Caretta CA, Caretta A, Baldini E. Protein-bound male urinary pheromones: differential responses according to age and gender. *Chem Senses* 1998;23:67–70.
- [24] Stopková R, Stopka P, Janotová K, Jedelský PL. Species-specific expression of major urinary proteins in the house mice (*Mus musculus musculus* and *Mus musculus domesticus*). *J Chem Ecol* 2007;33: 861–9.