

```
In[1]:= Clear["Global`*"];
```

Problem

Solve numerically the differential equation (in atomic units $\hbar = 1$, $m_e = 1$)

$$i \frac{\partial \psi(x, t)}{\partial t} = -\frac{1}{2\mu} \frac{\partial^2 \psi(x, t)}{\partial x^2} \quad (1)$$

with the following initial condition

$$\psi(x, 0) = \frac{1}{(2\pi\sigma^2)^{1/4}} e^{-(x-x_0)^2/(4\sigma^2) + i p x} \quad (2)$$

and Dirichlet boundary conditions

$$\begin{aligned} u(-\infty, t) &= 0 \\ u(+\infty, t) &= 0 \end{aligned} \quad (3)$$

Exact solution

Initial normalized Gaussian packet :

```
In[2]:= psi0[x_, x0_, σ_, p_] = 1 / (2 π σ^2)^(1/4) Exp[-(x - x0)^2 / (4 σ^2) + i p x];
Assuming[σ > 0,
Integrate[psi0[x, x0, σ, p] * psi0[x, x0, σ, -p], {x, -Infinity, Infinity}]]
```


Out[3]= 1

Exact solution:

```
In[4]:= psiExact[x_, t_, x0_, σ_, p_, μ_] = e^{-(x-x0)^2 μ - 2 i p^2 t σ^2 + p (-2 t x0 + 4 i x μ σ^2)} \left(\frac{2}{\pi}\right)^{1/4} \sqrt{\sigma} / \left(\sqrt{\frac{i t}{\mu} + 2 \sigma^2}\right);
```

Parameters:

```
In[5]:= pini = 3; mass = 1; xini = -10; σini = 1;
xmin = -15; xmax = 15; tmax = 5;
Plot3D[Abs[psiExact[x, t, xini, σini, pini, mass]]^2,
{x, xmin, xmax}, {t, 0, tmax}, PlotRange → All, PlotPoints → 100]
```



Numerical solution in the FEM-DVR basis

Gauss-Lobatto quadrature points and weights will be used for integration over one element

```
In[8]:= getGaussLobattoPointsAndWeights[n_, a_, b_] :=
Module[{x, w, p},
(* roots of the derivative of the (n-1)st Legendre polynomial are inner points
of the Gauss-Lobatto quadrature on [-1,1]*)
p[z_] = LegendreP[n - 1, z];
If[n == 2,
x = {-1.0, 1.0},
x = N[Flatten[{{-1.0, Sort[Re[z /. N[Solve[D[p[z], z] == 0, z], 16]]]}, 1.0}]]];
];
(* to get weights we need values of this polynomial *)
w = 2.0
Flatten[{1.0, Table[1.0 / (N[p[x[[i]]], 16])^2, {i, 2, n - 1}], 1.0}] / (n (n - 1));
(* shifting and scaling to the interval [a,b]*)
x = (b - a) x / 2 + (b + a) / 2;
w = (b - a) w / 2;
Return[{x, w}]
]
getGaussLobattoPointsAndWeights[5, 0, 1]

Out[9]= {{0., 0.1726731646, 0.5, 0.8273268354, 1.},
{0.05, 0.2722222222, 0.3555555556, 0.2722222222, 0.05}}
```

Full grid and weights

```
In[10]:= getFEMPointsAndWeights[nGL_, endPoints_] :=
Module[{nEl, nPoints, xGL, wGL, x, w},
nEl = Length[endPoints] - 1;
nPoints = nEl * (nGL - 1) + 1;
(* Print["Number of all points/basis functions is ", nPoints]; *)
x = ConstantArray[0.0, nPoints];
w = ConstantArray[0.0, nPoints];
Do[
{xGL, wGL} =
getGaussLobattoPointsAndWeights[nGL, endPoints[[i]], endPoints[[i + 1]]];
x[[[(i - 1) * (nGL - 1) + 1 ;; i * (nGL - 1) + 1]]] = xGL;
(* weights at points which are common to two elements are added up *)
w[[[(i - 1) * (nGL - 1) + 1 ;; i * (nGL - 1) + 1]]] += wGL,
{i, 1, nEl}
];
Return[{x, w}]
]
getFEMPointsAndWeights[4, {0, 1, 3, 6}]

Out[11]= {{0., 0.2763932023, 0.7236067977, 1., 1.552786405, 2.447213595, 3.,
3.829179607, 5.170820393, 6.}, {0.08333333333, 0.41666666667, 0.41666666667,
0.25, 0.83333333333, 0.83333333333, 0.41666666667, 1.25, 1.25, 0.25}}
```

Derivatives of the Lagrange polynomials at GL points on [-1,1] - result is a matrix nGL x nGL of $D[l_i(x), x = x_k]$

```
In[12]:= derivativesLagPol[nGL_] :=
Module[{xGL, wGL, dLP, hlp},
dLP = ConstantArray[0.0, {nGL, nGL}];
{xGL, wGL} = getGaussLobattoPointsAndWeights[nGL, -1.0, 1.0];
Do[
(* Diagonal terms *)
dLP[[i, i]] = 0.0;
Do[
If[i != s, dLP[[i, i]] = dLP[[i, i]] + 1.0 / (xGL[[i]] - xGL[[s]])],
{s, 1, nGL}
];
(* Off-diagonal terms *)
Do[
hlp = 1.0;
Do[
If[(j != i) && (j != k), hlp = hlp * (xGL[[k]] - xGL[[j]]) / (xGL[[i]] - xGL[[j]])],
{j, 1, nGL}
];
dLP[[i, k]] = hlp / (xGL[[i]] - xGL[[k]]);
dLP[[k, i]] = 1.0 / (hlp * (xGL[[k]] - xGL[[i]])),
{k, i + 1, nGL}
],
{i, 1, nGL}
];
Return[dLP];
];
derivativesLagPol[4]
Out[13]= {{-3., -0.8090169944, 0.3090169944, -0.5},
{4.045084972, -3.330669074 \times 10^{-16}, -1.118033989, 1.545084972},
{-1.545084972, 1.118033989, 2.220446049 \times 10^{-16}, -4.045084972},
{0.5, -0.3090169944, 0.8090169944, 3.}}
```

Construction of the stiffness matrix (ϕ'_i , ϕ'_j)

```

In[14]:= constructStiffnessMatrix[nGL_, endPoints_] :=
Module[{nEl, nPoints, xFEM, wFEM, xGL, wGL, dLP, dBf, k1, ii, jj, oldCorner, A},
nEl = Length[endPoints] - 1;
nPoints = nEl * (nGL - 1) + 1;
(* get weights for all points *)
{xFEM, wFEM} = getFEMPointsAndWeights[nGL, endPoints];
(* calculate derivatives of the Lagrange
interpolating polynomials at GL points on [-1,1] *)
dLP = derivativesLagPol[nGL];
(* build the stiffness matrix *)
A = ConstantArray[0.0, {nPoints, nPoints}];
oldCorner = 0.0;
Do[
{xGL, wGL} =
getGaussLobattoPointsAndWeights[nGL, endPoints[[k]], endPoints[[k + 1]]];
(* dilatation of derivatives of LP to be the derivatives
of the basis functions on the k-th element *)
dBf = 2.0 * dLP / (endPoints[[k + 1]] - endPoints[[k]]);
k1 = (k - 1) * (nGL - 1) + 1;
(* index of the first point of the k-th element in x *)
Do[
(* normalization factor of basis functions *)
dBf[[i, All]] = dBf[[i, All]] / Sqrt[wFEM[[k1 + i - 1]]],
{i, 1, nGL}
];
Do[
ii = k1 + i - 1; (* current row in the A matrix *)
Do[
jj = k1 + j - 1; (* current column in the A matrix *)
A[[ii, jj]] = Sum[wGL[[s]] * dBf[[i, s]] * dBf[[j, s]], {s, 1, nGL}];
A[[jj, ii]] = A[[ii, jj]],
{j, i, nGL}
],
{i, 1, nGL}
];
A[[k1, k1]] += oldCorner;
oldCorner = A[[k1 + nGL - 1, k1 + nGL - 1]],
{k, 1, nEl}
];
Return[A]
]
constructStiffnessMatrix[2, {0, 1, 2, 3, 4}] // MatrixForm

```

Out[15]//MatrixForm=

$$\begin{pmatrix} 2. & -1.414213562 & 0. & 0. & 0. \\ -1.414213562 & 2. & -1. & 0. & 0. \\ 0. & -1. & 2. & -1. & 0. \\ 0. & 0. & -1. & 2. & -1.414213562 \\ 0. & 0. & 0. & -1.414213562 & 2. \end{pmatrix}$$

Time evolution using the Crank-Nicolson implicit method

Parameters of numerical solution:

```
In[16]:= pini = 3.0; mass = 1.0; xini = -7.0; σini = 1.0;
{xmin, xmax} = {-15.0, 5.0};
{tmin, tmax} = {0.0, 1.0};
```

Set equidistant elements and calculate initial state and Hamiltonian matrix on the FEM-DVR grid:

```
In[19]:= nGL = 10;
nEl = 20;
endPoints = Table[N[xmin + i * (xmax - xmin) / nEl], {i, 0, nEl}];
{xFEM, wFEM} = getFEMPointsAndWeights[nGL, endPoints];
Nb = Length[xFEM];
Print["Number of points/basis functions: ", Nb];

(* coefficients of the initial wave packet ψ(x) in the FEM basis *)
psiini = N[psi0[xFEM, xini, σini, pini]] * Sqrt[wFEM];
(* stiffness matrix which is used to construct the Hamiltonian matrix *)
A = constructStiffnessMatrix[nGL, endPoints];
(* Hamiltonian matrix *)
H = A[[2 ;; Nb - 1, 2 ;; Nb - 1]] / (2.0 * mass);
Number of points/basis functions: 181
```

Time evolution:

```

In[28]:= {nx, nt} = {Nb - 2, 601}
dt = N[(tmax - tmin) / (nt - 1)];
T = N[Range[tmin, tmax, dt]];

(* Initialization of the array with zeroes - Dirichlet's boundary conditions *)
psi = ConstantArray[0.0, {nx, nt}];
error = ConstantArray[0.0, nt];

(* Initial state *)
psi[[All, 1]] = psiini[[2 ; Nb - 1]];

Do[
  (* Crank-Nicolson method *)
  psi[[All, n + 1]] = (IdentityMatrix[nx] - 0.5 * i * dt * H).psi[[All, n]];
  psi[[All, n + 1]] = LinearSolve[IdentityMatrix[nx] + 0.5 * i * dt * H, psi[[All, n + 1]]];
  {n, 1, nt - 1}
];


(* to get the functional values of the solution at grid
points we have to multiply the coefficients by Sqrt[w] *)
Do[
  psi[[All, n]] = psi[[All, n]] / Sqrt[wFEM[[2 ; Nb - 1]]],
  {n, 1, nt}
];

(* compare with the exact solution *)
Do[
  error[[n]] = 0.0;
  Do[
    error[[n]] = Max[error[[n]],
      Abs[psi[[j, n]] - psiExact[xFEM[[j + 1]], T[[n]], xini, oini, pini, mass]]],
    {j, 1, nx}
  ],
  {n, 1, nt}
];
ListLogPlot[{Table[{T[[n]], error[[n]]}, {n, 1, nt}]},
 PlotRange -> All]
Print["Maximal error: ", Max[error]]

(* fancy plotting *)
Manipulate[
 n = Round[t / dt] + 1;
 ListLinePlot[Table[{xFEM[[j + 1]], Abs[psi[[j, n]]^2]}, {j, 1, nx}],
 PlotRange -> {-0.5, 1.5}],
 {t, tmin, tmax, dt}
]

```

Out[28]= {179, 601}



Maximal error: 0.00002531399517

