

Static and radiating p -form black holes in the higher dimensional Robinson–Trautman class

Marcello Ortaggio*

Institute of Mathematics, Academy of Sciences of the Czech Republic
Žitná 25, 115 67 Prague 1, Czech Republic

Jiří Podolský† and Martin Žofka‡

Institute of Theoretical Physics, Faculty of Mathematics and Physics,
Charles University in Prague, V Holešovičkách 2, 180 00 Prague 8, Czech Republic

November 27, 2014

Abstract

We study Robinson–Trautman spacetimes in the presence of an aligned p -form Maxwell field and an arbitrary cosmological constant in $n \geq 4$ dimensions. As it turns out, the character of these exact solutions depends significantly on the (relative) value of n and p . In *odd* dimensions the solutions reduce to static black holes dressed with an electric and a magnetic field, with an Einstein space horizon (further constrained by the Einstein–Maxwell equations) — both the Weyl and Maxwell types are D. *Even* dimensions, however, open up more possibilities. In particular, when $2p = n$ there exist non-static solutions describing black holes gaining (or losing) mass by receiving (or emitting) electromagnetic radiation. In this case the Weyl type is II (D) and the Maxwell type can be II (D) or N. Conditions under which the Maxwell field is self-dual (for odd p) are also discussed, and a few explicit examples presented. Finally, the case $p = 1$ is special in all dimensions and leads to static metrics with a non-Einstein transverse space.

PACS 04.50.+h, 04.20.Jb, 04.40.Nr

Contents

1	Introduction	3
2	Robinson–Trautman geometry with aligned Maxwell fields	6

*ortaggio(at)math(dot)cas(dot)cz

†podolsky(at)mbox(dot)troja(dot)mff(dot)cuni(dot)cz

‡zofka(at)mbox(dot)troja(dot)mff(dot)cuni(dot)cz

3	Integration of the Einstein–Maxwell field equations	7
3.1	Maxwell equations, step one	7
3.2	Einstein equations for R_{ij} and R_{ur}	8
3.3	Maxwell equations, step two ($2p \neq n$)	10
3.4	Einstein equation for R_{ui} ($2p \neq n; p \neq 1, n-1$)	11
3.5	Einstein equation for R_{uu} ($2p \neq n; p \neq 1, n-1; n > 4$)	11
3.5.1	Case $2p \neq n, n \pm 1, n \pm 2$	11
3.5.2	Case $2p = n \pm 1$	12
3.5.3	Case $2p = n \pm 2$	12
4	Summary for the generic case $2p \neq n$ ($n > 4$): static black holes	13
4.1	Case $2p \neq n \pm 2$: static black holes	13
4.1.1	Metric and Maxwell field	13
4.1.2	Weyl and Maxwell types	14
4.2	Exceptional case $2p = n \pm 2$ ($n \geq 6$, even)	15
5	Special case $2p = n$ (n even): black holes with electromagnetic radiation	16
5.1	Remaining field equations for the case $2p = n$	16
5.1.1	Maxwell equations, step two	16
5.1.2	Einstein equation for R_{ui}	17
5.1.3	Einstein equation for R_{uu}	17
5.2	Summary and discussion	18
5.2.1	Maxwell type and self-duality	18
5.2.2	Weyl type	19
5.2.3	Examples	19
5.3	\mathbf{l} aligned with \mathbf{F} : static black holes ($n \geq 8$)	21
5.4	Type N Maxwell field	21
5.4.1	A special subcase: solutions with a factorized $h(u, x) = U(u)X(x)$	22
A	Robinson-Trautman spacetimes with an aligned Ricci tensor of type II	23
A.1	General metric	23
A.2	Ricci tensor components	24
A.2.1	Case $W^i = 0$ (\mathbf{k} is a multiple WAND — Weyl type II(bd))	25
A.3	Weyl tensor components in the case $R_{ij} \propto h_{ij}$ ($\Rightarrow W^i = 0$)	25
A.3.1	Conditions for the Weyl types II(abd), II(bcd) and III(b)	26
A.3.2	Conditions for the Weyl type N	26
A.3.3	Conditions for \mathbf{l} to be a multiple WAND (\Rightarrow Weyl type D(bd))	27
A.3.4	Conditions for the Weyl type D(bd) when \mathbf{l} is <i>not</i> a multiple WAND	27
B	Comments on the (Einstein) constraints (31) and (32) ($2 \leq p \leq n-2$)	28
B.1	Magnetic fields	28
B.2	Electric fields	29
B.3	Case $2p = n$ (n even)	30
C	Cases $p = 1$ and $p = n-1$	30
C.1	Case $n > 4$	31
C.2	Case $n = 4$	32

1 Introduction

Gravity in more than four spacetime dimensions has attracted a lot of interest in recent years. In particular, several properties of higher dimensional black holes have been elucidated (see, e.g., the reviews [1–3]). In addition to vacuum spacetimes, solutions of various theories with gauge fields have also been investigated extensively (numerous references being given in [1–3]). The simplest such theory one can consider is probably n -dimensional Einstein–Maxwell gravity in the presence of a single 2-form field $F_{\mu\nu}$. Among its solutions, an analog of the Reissner–Nordström spacetime has been known for a long time [4], together with its generalizations admitting an Einstein horizon [5] (in contrast, no $n > 4$ exact solution is known that would extend the Kerr–Newman metric, except in the vacuum limit [6], or in other theories [1–3]). However, since extended objects naturally couple to higher-rank forms, theories of gravity with $p > 2$ forms are also of interest, especially from the standpoint of supergravity and string theory. A direct generalization of Einstein–Maxwell gravity is thus the p -form theory defined by

$$S = \frac{1}{16\pi} \int d^n x \sqrt{-g} \left(R - 2\Lambda - \frac{\kappa_0}{p} F^2 \right), \quad (1)$$

where $F^2 = F_{\alpha_1 \dots \alpha_p} F^{\alpha_1 \dots \alpha_p}$ (and κ_0 is a constant taking into account different possible normalizations found in the literature, cf. also footnote 3), to which we shall restrict ourselves in the paper.

Although the theory (1) is in some respects very similar to standard Einstein–Maxwell gravity, the case $p = 2$ possesses some distinct features. For example, it has been shown recently that asymptotically flat static black holes cannot couple to electric p -form fields when $(n + 1)/2 \leq p \leq n - 1$ (and thus do not possess dipole hair) [7] and that, for any $p > 2$, static perturbations of the vacuum Schwarzschild–Tangherlini metric do not exist [8, 9]. In addition, results of [10] indicate that electromagnetic radiation may have properties different from those of standard $n = 4$, $p = 2$ electrovac general relativity (except possibly for $2p = n$), as confirmed in [11] in the case of test fields.

A relatively simple and yet rich class of exact solutions in $n = 4$ general relativity is given by the Robinson–Trautman family [12] (cf. the reviews [13, 14]), defined by the existence of a geodesic, shear-free, twist-free but expanding null vector field \mathbf{k} . It includes static black holes with an arbitrary cosmological constant, their accelerating counterpart (the C-metric) and other spacetimes containing both gravitational and electromagnetic radiation, as well as pure radiation solutions such as the Vaidya metric. Some of these solutions have found useful applications also beyond general relativity, for example to describe $2 + 1$ black holes on a brane [15], or in the context of the AdS/CFT correspondence [16]. Electrovac Robinson–Trautman spacetimes, in particular, have been investigated thoroughly in the presence of a Maxwell field $F_{\mu\nu}$ aligned with \mathbf{k} (already starting in [12]) and can be of Petrov type II, D and III (not N and O), while the Maxwell field can be both of type D (“non-null”) and N (“null”), but it must be null if the Petrov type is III (see section 28.2 of [13] and section 19.6 of [14] for reviews and for a number of original references). Within this class, solutions of Petrov type D with a null Maxwell field [12]¹ are of special interest, as they describe formation of black holes by gravitational collapse of purely electromagnetic radiation [21] (see [22–24] for related earlier studies).

¹It is interesting to observe that the “example” given in section 7 (ii) of [12] (eq. (28.43) of [13]) is in fact (up to adding a cosmological constant) the *unique* [17] solution of the 4D Einstein–Maxwell equations that is simultaneously both of Petrov type D and of Maxwell type N (in which case the Maxwell and Weyl tensor *necessarily* share a multiple principal null direction by the Mariot–Robinson and the (generalized) Goldberg–Sachs theorems [13], see also [18]) — cf. also [18–20] for various steps towards the complete proof [17] of this statement.

In [25] it was shown how the Robinson–Trautman line-element can be constructed in arbitrary dimension n (see also [26–30] for additional results). In the context outlined above, it is thus of interest to study Robinson–Trautman solutions of the theory (1) with arbitrary n and p , which is the purpose of the present paper. We observe that the case $p = 2$ has been already studied in detail in [27]. Similarly as in the vacuum case [25], it turned out that the Robinson–Trautman class is much more restricted when $n > 4$, and it essentially contains only static black hole spacetimes of Weyl type D (plus a few special non-static solutions [25–28]). However, the results of [10, 11] mentioned above suggest that it need not be so for the theory (1). As we will work out, this expectation turns out to be correct. In more detail, our results can be summarized as follows.

- In *odd* n dimensions, Robinson–Trautman spacetimes coupled to a p -form Maxwell field with $2 \leq p \leq n - 2$ reduce to *static black holes* specified by four independent parameters related to mass, electric and magnetic field strengths, and cosmological constant (section 4.1). These black holes have been already studied in [31], to which we add a few comments. Except for the case $p = 2$ ($p = n - 2$) of [4], they cannot be asymptotically flat (so there is no conflict with the results of [7–9]), but they can be asymptotically locally (A)dS in some cases. The horizon is an Einstein space but must also obey further constraints following from the field equations. The metric is (55) with (56) and the Maxwell field is (57) (see the text for more details). Both the Weyl and Maxwell tensors are of type D and share the same pair of doubly aligned (geodesic) null directions.
- In *even* n dimensions (with $2 \leq p \leq n - 2$), static black holes as those described above are also present (cf. also [31]) and, again, the horizon geometry is constrained by the field equations (for example, for $p = 2$ it must be almost-Kähler if the magnetic field is non-zero [27]). Moreover, in addition to static black holes, there are also some exceptional solutions when $2p = n \pm 2$ and $n \geq 6$ (as first noticed in [27] for the case $n = 6, p = 2$), cf. section 4.2 for details. Generically, these exceptional metrics are non-static and of Weyl type II, while the Maxwell type is still D (but now possesses also a non-geodesic aligned null direction).
- In even n dimensions, the unique rank p satisfying $2p = n$ (including, in particular, $n = 4$ with $p = 2$ [12–14]) gives rise to an additional (and more interesting) new class of solutions (section 5), consisting of the metric (55) with (77) and the Maxwell field (76). The Maxwell field is allowed to be of type II, D or N and in all these cases it can have a radiative term (thus giving concrete examples to the predictions of [10, 11]), while the Weyl tensor can be of type II or D (more precisely, II(bd)/II(bcd) or D(bd)/D(bcd)) but with no radiative term for $n > 4$. However, when the Maxwell type is N and $n > 4$ then the Weyl type can only be D(bd)/D(bcd) (this is a significant difference w.r.t the $n = 4, p = 2$ case, which can be traced back to the absence of gravitational radiation in the vacuum higher dimensional Robinson–Trautman class [25]). Similarly as in [21], some of these solutions can be used to describe black hole formation (or white hole evaporation, by time-reversal) by collapse (emission) of electromagnetic radiation. As in the static case, these black holes can be asymptotically locally (A)dS for certain choices of the parameters — again, the horizon is an Einstein space and can be flat, in particular. For *odd* p , some of these solutions possess a *self-dual* p -form field, a property which is of interest in supergravity and string theory [32].
- Both in odd and even dimensions, the rank $p = 1$ (or its dual $p = n - 1$) has special features and needs to be studied separately (appendix C). This results again in a family of static solutions of Weyl and Maxwell type D. In this case the transverse space cannot be an Einstein space since it “feels” the backreaction of the electromagnetic field (as opposed to the generic case $2 \leq p \leq n - 2$), which also defines a preferred direction here. Further, the transverse space cannot be a space with an everywhere non-negative Ricci scalar. There exists no $n = 4$ counterpart of these solutions.

The rest of the paper is organized as follows. In section 2 we describe our assumptions and set up the corresponding general form of the line-element (based on [25]) and of the Maxwell field. Section 3 is devoted to a systematic integration of the resulting Einstein–Maxwell equations — it can well be skipped by readers not interested in those technicalities. Our results are summarized in section 4 for the generic case $2p \neq n$, and in sections 5.2–5.4 for the special case $2p = n$ (section 5.1 contains integration of a subset of the Einstein–Maxwell equations that are special when $2p = n$). Appendix A, largely based on [25, 27, 29, 30], summarizes certain general properties of Robinson–Trautman spacetimes useful in the paper, but also contains a few new observations (sections A.3.1–A.3.4). Appendix B discusses some geometrical properties of the transverse metric h_{ij} of (55) (in particular, the black hole horizon) that follow from the Einstein equations (it partly overlaps with [27] when $p = 2$ or $p = n - 2$, and also summarizes certain observations of [31]). Appendix C studies the special ranks $p = 1$ and $p = n - 1$, also showing that these are forbidden in the four dimensional Robinson–Trautman class.

Notation and conventions

Throughout the paper we focus on $n > 4$ dimensions, but large part of the results applies also to the $n = 4$ case, on which we shall comment explicitly when important differences arise. We consider a p -form field $\mathbf{F} = \frac{1}{p!} F_{\alpha_1 \dots \alpha_p} dx^{\alpha_1} \wedge \dots \wedge dx^{\alpha_p}$ (with $1 \leq p \leq n - 1$)² that, in the theory (1), satisfies the source-free Maxwell equations $d^* \mathbf{F} = 0$, $d\mathbf{F} = 0$ or, in components,

$$(\sqrt{-g} F^{\mu\alpha_1 \dots \alpha_{p-1}})_{,\mu} = 0, \quad F_{[\alpha_1 \dots \alpha_p, \mu]} = 0. \quad (2)$$

The Maxwell field \mathbf{F} backreacts on the spacetime geometry via the energy-momentum tensor

$$T_{\mu\nu} = \frac{\kappa_0}{8\pi} \left(F_{\mu\alpha_1 \dots \alpha_{p-1}} F_{\nu}{}^{\alpha_1 \dots \alpha_{p-1}} - \frac{1}{2p} g_{\mu\nu} F^2 \right), \quad (3)$$

where $F^2 = F_{\alpha_1 \dots \alpha_p} F^{\alpha_1 \dots \alpha_p}$. Note that $8\pi T = \kappa_0(2p - n)F^2/(2p)$. With (3), Einstein’s equations with a cosmological constant $R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi T_{\mu\nu}$ (following from (1)) take the form

$$R_{\mu\nu} = \frac{2}{n-2} \Lambda g_{\mu\nu} + \kappa_0 \left[F_{\mu\alpha_1 \dots \alpha_{p-1}} F_{\nu}{}^{\alpha_1 \dots \alpha_{p-1}} - \frac{p-1}{p(n-2)} g_{\mu\nu} F^2 \right]. \quad (4)$$

It may be useful to recall the well-known fact that for any solution of (2) and (4) with a given p -form \mathbf{F} , a “discrete” duality transformation $\mathbf{F} \rightarrow * \mathbf{F}$ gives rise to a dual solution³ where the same metric is coupled to the $(n - p)$ -form $* \mathbf{F}$, defined by $* F_{b_1 \dots b_{n-p}} \equiv \frac{1}{p!} \epsilon^{a_1 \dots a_p}{}_{b_1 \dots b_{n-p}} F_{a_1 \dots a_p}$. In the special case $2p = n$, (anti-)self-dual p -form solutions $* \mathbf{F} = \pm \mathbf{F}$ (for which necessarily $\mathbf{F}^2 = 0 = * \mathbf{F} \cdot \mathbf{F}$, where total contraction is understood) may exist if p is *odd* (since \mathbf{F} is real and the signature Lorentzian, see, e.g., [32]). Instead, for $2p = n$ with *even* p (anti-)self-dual p -forms do not exist — on the other hand, in that case there is a continuous $SO(2)$ duality symmetry (in addition to the discrete one mentioned above) which maps solutions into solutions (cf., e.g., [34] and appendix A of [35]).

²The limiting value $p = n$ could also be included but, as well-known (cf., e.g., [33]), this is trivial in the sense that \mathbf{F} is simply given by the spacetime volume element (up to a constant rescaling) and acts on the geometry as an effective positive cosmological constant. Similarly, the dual case $p = 0$ reduces to a constant scalar field.

³To be precise, in order for $T_{\mu\nu}$ to be invariant under $\mathbf{F} \rightarrow * \mathbf{F}$, κ_0 should be replaced by $\kappa_0/(p - 1)!$ in (3) and (4) (which also shows how the limiting case $p = 0$ can be formally incorporated in the discussion) — this rescaling is not necessary in the case $2p = n$. While bearing this in mind, for compactness throughout the paper we will employ the simpler expressions (3) and (4).

2 Robinson–Trautman geometry with aligned Maxwell fields

We consider a n -dimensional spacetime that admits a non-twisting, non-shearing, expanding geodesic null vector field \mathbf{k} . The associated Robinson–Trautman line-element was obtained in adapted coordinates in [25]. The corresponding curvature has been fully computed recently in [30], showing in particular that the spacetime is generically of aligned Weyl type I(b).

It is the purpose of this paper to determine Robinson–Trautman spacetimes in the presence of a Maxwell p -form field in the theory (1), i.e., such that the Einstein equations (4) are satisfied, along with the Maxwell equations (2). However, we shall restrict to the case of Maxwell fields that are *aligned* with \mathbf{k} (so that \mathbf{F} is of type II or more special).⁴ This means that the components of \mathbf{F} of b.w. +1 vanish, which by (4) implies that the Ricci tensor components of b.w. +2 and +1 must also vanish, i.e., the Ricci type is II (or more special) aligned with \mathbf{k} . With this condition, we can take advantage of previous results of [25] (summarized in theorem A.1 of appendix A) asserting that the spacetimes in question can be represented by the line-element

$$\begin{aligned} ds^2 &= r^2 h_{ij} (dx^i + W^i du) (dx^j + W^j du) - 2 du dr - 2H du^2, \\ h_{ij} &= h_{ij}(u, x), \quad W^i = \alpha^i(u, x) + r^{1-n} \beta^i(u, x). \end{aligned} \quad (5)$$

Here the adapted coordinates $(u, r, x^1, \dots, x^{n-2})$ are used (Latin indices i, j, \dots or i_1, i_2, \dots etc. range over $1, \dots, n-2$ and label the spatial coordinates x^i , sometimes collectively denoted simply as x) such that

$$\mathbf{k} \equiv k^\mu \partial_\mu = \partial_r, \quad k_\mu dx^\mu = -du, \quad (6)$$

where r is an affine parameter along the generator \mathbf{k} of the null hypersurfaces $u = \text{const.}$ In such coordinates, the assumed alignment condition on \mathbf{F} takes the form

$$F_{ri_1 \dots i_{p-1}} = 0, \quad (7)$$

or, equivalently, $F^{ui_1 \dots i_{p-1}} = 0$, while the alignment conditions on the Ricci tensor (automatically satisfied by (5)) read $R_{rr} = 0 = R_{ri}$. By construction, the corresponding components of the Einstein equation (4) are thus identically satisfied (note also that (7) implies $T_{rr} = 0 = T_{ri}$, cf. (3)).

In the rest of the paper we thus need to study only the remaining Einstein equations for R_{ij} , R_{ur} , R_{ui} and R_{uu} . These will contain terms depending on the Maxwell field. Hence, in order to proceed it will be convenient to first fix the r -dependence of \mathbf{F} using Maxwell's equations (2).

For later calculations it is also useful to note that

$$\sqrt{-g} = r^{n-2} \sqrt{h}, \quad (8)$$

where $g \equiv \det g_{\mu\nu}$ and $h(u, x) \equiv \det h_{ij}$.

⁴We refer to the boost weight (b.w.) classification of a general tensor [36] — cf., e.g., [37–40] for further results in the particular case of 2-forms. Let us observe that since \mathbf{k} is shearfree and expanding here, a result of [10] implies that \mathbf{F} cannot be doubly aligned with it (i.e., it cannot be of type N), except possibly when $2p = n$ in even dimensions. We will prove that type N fields do indeed exist in that special case (section 5).

3 Integration of the Einstein–Maxwell field equations

3.1 Maxwell equations, step one

With (7), the “geometrical” equations $F_{[\alpha_1 \dots \alpha_p, \mu]} = 0$ give

$$F_{i_1 \dots i_p, r} = 0, \quad (9)$$

$$F_{[i_1 \dots i_p, j]} = 0, \quad (10)$$

$$F_{uj i_1 \dots i_{p-2}, r} = (p-1) F_{ur [i_1 \dots i_{p-2}, j]}, \quad (11)$$

$$F_{j i_1 \dots i_{p-1}, u} = p F_{u [i_1 \dots i_{p-1}, j]}. \quad (12)$$

Using relation (8), the “dynamical” equations $(\sqrt{-g} F^{\mu r i_1 \dots i_{p-3}})_{,\mu} = 0$ (not present for $p \leq 2$) and $(\sqrt{-g} F^{\mu u i_1 \dots i_{p-2}})_{,\mu} = 0$ (not present for $p \leq 1$) read, respectively,

$$(\sqrt{h} F^{ur j i_1 \dots i_{p-3}})_{,j} = 0, \quad (r^{n-2} F^{ur i_1 \dots i_{p-2}})_{,r} = 0. \quad (13)$$

As it turns out, the remaining equations $(\sqrt{-g} F^{\mu i_1 \dots i_{p-1}})_{,\mu} = 0$ and $(\sqrt{-g} F^{\mu r i_1 \dots i_{p-2}})_{,\mu} = 0$ become significantly simpler once some of the Einstein equations are enforced, and it is thus convenient to postpone their discussion to section 3.3.

The r -dependence of the Maxwell field is thus completely determined by (9), the second of (13) and (11), and can be summarized as

$$F_{i_1 \dots i_p} = b_{i_1 \dots i_p}, \quad (14)$$

$$F_{ur i_1 \dots i_{p-2}} = r^{2p-2-n} e_{i_1 \dots i_{p-2}}, \quad (15)$$

$$F_{uj i_1 \dots i_{p-2}} = r^{2p-1-n} \frac{p-1}{2p-1-n} e_{[i_1 \dots i_{p-2}, j]} + f_{j i_1 \dots i_{p-2}} \quad (2p \neq n+1), \quad (16)$$

where lowercase symbols b , e or f denote integration functions independent of r . Since we are restricting to $1 \leq p \leq n-1$, $b_{i_1 \dots i_p} = b_{[i_1 \dots i_p]}$ can be non-zero only for $1 \leq p \leq n-2$, and $e_{i_1 \dots i_{p-2}} = e_{[i_1 \dots i_{p-2}]}$ for $2 \leq p \leq n-1$ (for $p=2$ the term $e_{i_1 \dots i_{p-2}}$ obviously reduces to a scalar function e). From now on we will use the convention that indices of b , e and f are raised with the spatial metric h^{ij} so that, e.g.,

$$\begin{aligned} b^{i_1 \dots i_p} &= h^{i_1 j_1} \dots h^{i_p j_p} b_{j_1 \dots j_p}, & e^{i_1 \dots i_{p-2}} &= h^{i_1 j_1} \dots h^{i_{p-2} j_{p-2}} e_{j_1 \dots j_{p-2}}, \\ f^{i_1 \dots i_{p-1}} &= h^{i_1 j_1} \dots h^{i_{p-1} j_{p-1}} f_{j_1 \dots j_{p-1}}. \end{aligned} \quad (17)$$

In the special case with $2p = n+1$ ($n \geq 5$ odd, $p \geq 3$), eq. (16) is replaced by⁵

$$F_{uj i_1 \dots i_{p-2}} = \frac{n-1}{2} e_{[i_1 \dots i_{p-2}, j]} \ln r + f_{j i_1 \dots i_{p-2}} \quad (2p = n+1). \quad (18)$$

(However, we will see in section 3.3 that the above logarithmic term must in fact vanish.) From (14) we observe, in particular, that the magnetic components $F_{i_1 \dots i_p}$ are always r -independent, whereas the electric components $F_{ur i_1 \dots i_{p-2}}$ become r -independent only in the special case $2p = 2 + n$ (n even).

⁵Obviously, for dimensional reasons we should write $\ln(r/r_0)$ instead of $\ln r$, where r_0 is an r -independent coefficient with the dimension of length. However, r_0 can be absorbed in the following term $f_{j i_1 \dots i_{p-2}}$ and thus for brevity we will omit it. Similar comments apply to further logarithmic terms that will appear in other expressions in the following, and will not be repeated there.

Using (15) and (14), the first of (13) and (10) can be rewritten simply as

$$(\sqrt{h} e^{j i_1 \dots i_{p-3}})_{,j} = 0 \quad (p \geq 3), \quad b_{[i_1 \dots i_p, j]} = 0. \quad (19)$$

For later purposes, it will be useful to define the following r -independent quantities built out of the electric and magnetic parts of the Maxwell field (which will enter some of the Einstein equations)

$$\mathcal{E}_{ij}^2 \equiv e_{ik_1 \dots k_{p-3}} e_j^{k_1 \dots k_{p-3}}, \quad \mathcal{E}^2 \equiv h^{ij} \mathcal{E}_{ij}^2 \quad (p \geq 3), \quad (20)$$

$$\mathcal{B}_{ij}^2 \equiv b_{ik_1 \dots k_{p-1}} b_j^{k_1 \dots k_{p-1}}, \quad \mathcal{B}^2 \equiv h^{ij} \mathcal{B}_{ij}^2 \quad (p \leq n-2). \quad (21)$$

Clearly $\mathcal{E}_{ij}^2 = \mathcal{E}_{(ij)}^2$ and $\mathcal{B}_{ij}^2 = \mathcal{B}_{(ij)}^2$. In the case $p = 2$ the indices i, j disappear from $e_{i_1 \dots i_{p-2}}$ and \mathcal{E}_{ij}^2 and we simply have $\mathcal{E} = e$, while \mathcal{B}_{ij}^2 is identically zero for $p > n-2$. Obviously $\mathcal{E}^2 = 0 \Leftrightarrow e_{i_1 \dots i_{p-2}} = 0$ and $\mathcal{B}^2 = 0 \Leftrightarrow b_{i_1 \dots i_p} = 0$ since h_{ij} is positive definite.

In particular, the invariant F^2 (useful in the following) can be written as

$$F^2 = -r^{2(p-n)} p(p-1) \mathcal{E}^2 + r^{-2p} \mathcal{B}^2. \quad (22)$$

3.2 Einstein equations for R_{ij} and R_{ur}

Knowing the r -dependence of the Maxwell field, we can now consider the remaining Einstein equations. It is convenient to start from the equations for the Ricci components R_{ij} and R_{ur} (those of the highest remaining b.w., namely zero). From (4) with (5), (14), (15), (22) (using the definitions (20), (21)), these read

$$R_{ij} = \frac{2}{n-2} \Lambda r^2 h_{ij} + \kappa_0 \left\{ r^{2(p+1-n)} (p-1) \left[-(p-2) \mathcal{E}_{ij}^2 + \frac{p-1}{n-2} \mathcal{E}^2 h_{ij} \right] + r^{2(1-p)} \left[\mathcal{B}_{ij}^2 - \frac{p-1}{p(n-2)} \mathcal{B}^2 h_{ij} \right] \right\}, \quad (23)$$

$$-R_{ur} = \frac{2}{n-2} \Lambda - \kappa_0 \frac{p-1}{n-2} \left[r^{2(p-n)} \mathcal{E}^2 (n-p-1) + r^{-2p} \mathcal{B}^2 p^{-1} \right]. \quad (24)$$

The component R_{ij} of the metric (5) is reproduced in appendix A as eq. (A4). Comparing this with (23) immediately reveals that $\beta^i = 0$ (this follows by comparing various powers of r in (23) and (A4), which implies that $\beta^i \beta^j$ is either zero or proportional to h^{ij} — the latter option is however impossible, cf. also [25]).⁶ One can further perform a coordinate transformation (at least locally) to set $\alpha^i = 0$ [25]. From now on we shall thus have in (5)

$$W^i = 0, \quad (25)$$

and therefore $g^{ri} = 0 = g_{ui}$, which will simplify several expressions. In particular, the Weyl type will thus be II(d) or more special, aligned with \mathbf{k} (cf. theorem A.1).

Using (25), the component (A4) now simplifies drastically (cf. (A5)). By comparing its various powers of r with those of (23), after r -integration (and contraction with h^{ij} when necessary) one readily determines the r -dependence of the metric function H

⁶One might think that the (dual) cases $p = 1$ and $p = n - 1$ escape this conclusion since the $r^{2(2-n)}$ term of (A4) falls off in those cases as the term $r^{2(p+1-n)}$ or $r^{2(1-p)}$ of (23), respectively. However, for $p = 1$ [$p = n - 1$] the $r^{2(p+1-n)}$ [$r^{2(1-p)}$] term of (23) vanishes identically, cf. (20) and (21), so that the conclusion $\beta^i = 0$ remains true.

$$2H = \frac{\mathcal{R}}{(n-2)(n-3)} + \frac{2(\ln \sqrt{h})_{,u}}{n-2} r - \frac{2\Lambda}{(n-1)(n-2)} r^2 - \frac{\mu}{r^{n-3}} + \frac{\kappa_0}{n-2} \left[\frac{p-1}{n+1-2p} \frac{\mathcal{E}^2}{r^{2(n-p-1)}} - \frac{1}{p(n-1-2p)} \frac{\mathcal{B}^2}{r^{2(p-1)}} \right] \quad (2p \neq n \pm 1), \quad (26)$$

where μ is an arbitrary integration function independent of r .

In the special odd dimensional cases $2p = n \pm 1$ the last two terms of (26) should be replaced, respectively, by

$$+ \frac{\kappa_0}{n-2} \left[-\frac{n-1}{2} \mathcal{E}^2 \frac{\ln r}{r^{n-3}} + \frac{1}{n+1} \frac{\mathcal{B}^2}{r^{n-1}} \right] \quad (2p = n+1), \quad (27)$$

$$+ \frac{\kappa_0}{n-2} \left[\frac{n-3}{4} \frac{\mathcal{E}^2}{r^{n-1}} - \frac{2}{n-1} \mathcal{B}^2 \frac{\ln r}{r^{n-3}} \right] \quad (2p = n-1). \quad (28)$$

In addition, as a further consequence of the Einstein equation for R_{ij} , the following constraints (coming from terms of order r^0 , r , $r^{2(p+1-n)}$ and $r^{2(1-p)}$, respectively) must be satisfied (also when $2p = n \pm 1$)

$$\mathcal{R}_{ij} = \frac{\mathcal{R}}{n-2} h_{ij} \quad (p \neq 1, n-1), \quad (29)$$

$$h_{ij,u} = \frac{2(\ln \sqrt{h})_{,u}}{n-2} h_{ij}, \quad (30)$$

$$\mathcal{E}_{ij}^2 = \frac{\mathcal{E}^2}{n-2} h_{ij} \quad (3 \leq p \leq n-2, 2p \neq n), \quad (31)$$

$$\mathcal{B}_{ij}^2 = \frac{\mathcal{B}^2}{n-2} h_{ij} \quad (2 \leq p \leq n-2, 2p \neq n), \quad (32)$$

where \mathcal{R}_{ij} is the Ricci tensor associated with the spatial metric h_{ij} , \mathcal{R} is its Ricci scalar, and the general identity $h^{ij} h_{ij,u} = 2(\ln \sqrt{h})_{,u}$ has been used. For $p = 2$ eq. (31) becomes an identity. For $2p = n$ eqs. (31) and (32) are replaced by the following single equation

$$\frac{1}{4}(n-2)(n-4) \left(\mathcal{E}_{ij}^2 - \frac{\mathcal{E}^2}{n-2} h_{ij} \right) = \mathcal{B}_{ij}^2 - \frac{\mathcal{B}^2}{n-2} h_{ij} \quad (2p = n). \quad (33)$$

Eq. (33) is satisfied identically in the case $n = 4, p = 2$. Note that (31), (32) and (33) imply with (23) that in all cases $R_{ij} \propto h_{ij}$ (as can also be seen from (A5) with (29)).

As indicated above, equations (29), (31) and (32) do not hold in the limiting dual cases $p = 1$ (or $p = n-1$) — these are special since the \mathcal{B}_{ij}^2 (or \mathcal{E}_{ij}^2) terms behave as r^0 in (23), (24) (and therefore in (26)), and thus effectively act as sources for the transverse geometry. Since further differences will also arise in the remaining Einstein equations, we present all the results for $p = 1, n-1$ in appendix C and from now on we assume $p \neq 1, n-1$.

As in [25], relation (29) means that, at any given $u = u_0 = \text{const}$, the spatial metric $h_{ij}(x, u_0)$ must describe a $(n-2)$ -dimensional Riemannian Einstein space. It is well-known (see, e.g., p.76 of [41]) that for $n > 4$ (i.e., $n-2 > 2$) this implies that $\mathcal{R}_{,i} = 0$, so that \mathcal{R} can depend only on the coordinate u (additionally, for $n = 5$ the metric h_{ij} must be of constant curvature since it is 3-dimensional and Einstein). For $n = 4$ eq. (29) is instead an identity.

Equation (30) gives [25]

$$h_{ij} = h^{1/(n-2)} \gamma_{ij}(x) \quad \text{where } \det \gamma_{ij} = 1, \quad (34)$$

so that h_{ij} can depend on u only via the conformal factor $h^{1/(n-2)}$.

Eqs. (31) and (32) (or (33)) constrain both the Maxwell field and the metric h_{ij} , and the permitted relation between p and n — some comments are given in appendix B (see also [31], and [27] for the case $p = 2$).

The component R_{ur} of the metric (5) with (25) is reproduced in appendix A as eq. (A6). Substituting (26) into (A6) and comparing with (24), one finds that the corresponding Einstein equation is satisfied identically (including the cases $2p = n \pm 1$). We finally observe that (29) (with (25)) further restricts the Weyl tensor to be of type II(bd), aligned with \mathbf{k} (see theorem A.1).

3.3 Maxwell equations, step two ($2p \neq n$)

We can now turn to the remaining set of the Maxwell equations, namely $(\sqrt{-g} F^{\mu i_1 \dots i_{p-1}})_{,\mu} = 0$ and $(\sqrt{-g} F^{\mu r i_1 \dots i_{p-2}})_{,\mu} = 0$, which were not considered in section 3.1. As it turns out, from now on it will be necessary to consider the case $2p = n$ separately. Hereafter we thus restrict to the “generic” case $2p \neq n$, while the corresponding analysis for the special case $2p = n$ will be given later on in section 5.

Using (14), (16), and (25), $(\sqrt{-g} F^{\mu i_1 \dots i_{p-1}})_{,\mu} = 0$ contains three different powers of r , i.e., r^{n-2p-2} , r^{n-2p-1} , r^{-2} , generically leading, respectively, to

$$(\sqrt{h} b^{j i_1 \dots i_{p-1}})_{,j} = 0, \quad f_{j_1 \dots j_{p-1}} = 0, \quad e_{[j_2 \dots j_{p-1}, j_1]} = 0. \quad (35)$$

Note that the last two equations mean (see (16))

$$F_{u j_1 \dots j_{p-1}} = 0. \quad (36)$$

With (7) this implies that \mathbf{F} is aligned also with the null vector $\mathbf{l} = -\partial_u + H\partial_r$ (and is thus of type D), and will be important in the following (as a consequence, \mathbf{F} cannot be of type N, as found in [10]). Eq. (12) thus becomes

$$b_{i_1 \dots i_p, u} = 0. \quad (37)$$

Next, from $(\sqrt{-g} F^{\mu r i_1 \dots i_{p-2}})_{,\mu} = 0$ one gets (using (36))

$$(\sqrt{h} e^{i_1 \dots i_{p-2}})_{,u} = 0, \quad (38)$$

which with (34) simply gives

$$e_{i_1 \dots i_{p-2}} = h^{\frac{2p-n-2}{2(n-2)}} \tilde{e}_{i_1 \dots i_{p-2}}(x). \quad (39)$$

Note that the above results apply also in the $2p = n + 1$ case (18) with a logarithmic term, which thus in fact vanishes. For later purposes it is useful to observe that (39) and (37) imply, respectively (recall (20) and (21)), $\mathcal{E}_{ij}^2 = h^{(p-n+1)/(n-2)} \tilde{\mathcal{E}}_{ij}^2(x)$ and $\mathcal{B}_{ij}^2 = h^{(1-p)/(n-2)} \tilde{\mathcal{B}}_{ij}^2(x)$ (of course $\tilde{\mathcal{E}}_{ij}^2$ (or $\tilde{\mathcal{B}}_{ij}^2$) can be expressed in terms of $\tilde{e}_{i_1 \dots i_{p-2}}$ (or $\tilde{b}_{i_1 \dots i_p}$) and $\gamma^{ij}(x)$, if desired), and thus $\mathcal{E}^2 = h^{(p-n)/(n-2)} \tilde{\mathcal{E}}^2(x)$, $\mathcal{B}^2 = h^{-p/(n-2)} \tilde{\mathcal{B}}^2(x)$. These in turn imply

$$(n-2)(\mathcal{E}^2)_{,u} = -2(n-p)\mathcal{E}^2(\ln \sqrt{h})_{,u}, \quad (n-2)(\mathcal{B}^2)_{,u} = -2p\mathcal{B}^2(\ln \sqrt{h})_{,u}. \quad (40)$$

which will be useful later on.

We observe that above we have not employed eqs. (29)–(32), so that the results of the present section 3.3 apply also to the cases $p = 1, n - 1$.

3.4 Einstein equation for R_{ui} ($2p \neq n$; $p \neq 1, n-1$)

As remarked above, we now consider $2p \neq n$. Recalling (25) we have $g_{ui} = 0 = g^{ri}$. Using (7) and (36), we also obtain $F_{u\alpha_1 \dots \alpha_{p-1}} F_i^{\alpha_1 \dots \alpha_{p-1}} = 0$ (and thus $T_{ui} = 0$). The corresponding Einstein equation (4) thus simplifies considerably to

$$R_{ui} = 0, \quad (41)$$

where the explicit Ricci component is given by (A7). Employing (26) shows that this equation contains distinct powers of r , namely r^0 , r^{-1} , r^{2-n} , $r^{2p-2n+1}$, r^{1-2p} . The term of order r^0 vanishes identically thanks to (30), while the remaining terms immediately give, respectively, the following conditions

$$(n-4)\mathcal{R}_{,i} = 0 \quad (p \neq 1, n-1), \quad (42)$$

$$\mu_{,i} = 0, \quad (43)$$

$$(2p-n-2)(\mathcal{E}^2)_{,i} = 0 \quad (p \geq 2), \quad (44)$$

$$(2p-n+2)(\mathcal{B}^2)_{,i} = 0. \quad (45)$$

Eq. (42) is an identity due to (29) (as mentioned in section 3.2). One arrives at (42)–(45) also for $2p = n \pm 1$ after replacing (26) by the corresponding form of H containing the logarithmic terms (eqs. (27), (28)). Thus, generically, the Ricci curvature \mathcal{R} of the transverse $(n-2)$ -dimensional Riemannian space, the “mass” parameter μ , the electric scalar \mathcal{E}^2 and the magnetic scalar \mathcal{B}^2 must all be *independent of the spatial coordinates*. However, \mathcal{E}^2 and \mathcal{B}^2 can depend on x in the special cases $2p = n+2$ and $2p = n-2$, respectively (n necessarily even) — this will have some consequences in section 3.5. The case $n=4$ is also special in that \mathcal{R} can depend on the coordinates x — however, since the case $n=4, p=2$ is already well-known [12–14], only the cases $n=4, p=1, 3$ remain to be studied. These are precisely the ones dealt with in appendix C, so that from now on we can restrict in the main text to $n > 4$ with no loss of generality.

3.5 Einstein equation for R_{uu} ($2p \neq n$; $p \neq 1, n-1$; $n > 4$)

3.5.1 Case $2p \neq n, n \pm 1, n \pm 2$

Using $g_{uu} = -2H$, in view of (36) we obtain $F_{u\alpha_1 \dots \alpha_{p-1}} F_u^{\alpha_1 \dots \alpha_{p-1}} = 2H(p-1)r^{2(p-n)}\mathcal{E}^2$. With (22), the last Einstein equation (4) can thus be written in the simple form $R_{uu} = 2HR_{ur}$, where R_{ur} is the “on-shell” Ricci component (24). Using (A8) this means

$$-r^2(r^{-2}H)_{,r}(\ln \sqrt{h})_{,u} + (n-2)r^{-1}H_{,u} + r^{-2}\Delta H - (\ln \sqrt{h})_{,uu} - \frac{1}{4}h^{il}h^{jk}h_{ij,u}h_{kl,u} = 0, \quad (46)$$

where Δ is the Laplace operator associated with h_{ij} (cf. appendix A.2). Substituting (26) one easily sees that (for $2p \neq n \pm 1, n \pm 2$) this equation contains terms proportional to r^0 , r^{-1} , r^{2-n} , $r^{2p-2n+1}$, r^{1-2p} . Terms of order r^0 and r^{-1} do not contain the Maxwell field and, as shown in the vacuum case [25], vanish identically after using (30) and certain identities (see also the appendix of [29]). Terms of order $r^{2p-2n+1}$ and r^{1-2p} vanish thanks to (40). The only surviving r^{2-n} term fixes the u -dependence of $\mu(u)$ (as in vacuum)

$$(n-2)\mu_{,u} = -(n-1)\mu(\ln \sqrt{h})_{,u} \quad (2p \neq n \pm 2, n \pm 1, n > 4). \quad (47)$$

As in [27], taking the ∂_i derivative of (47) (with (43)) and of (40) (with (44) and (45)) gives immediately that $(\ln \sqrt{h})_{,ui} = 0$ (unless $\mu = \mathcal{E}^2 = \mathcal{B}^2 = 0$, which yields $\mathbf{F} = 0$ and thus a vacuum

spacetime [25]). This means $h(u, x) = U(u)X(x)$, but one can set $U(u) = 1$ by rescaling the coordinates u and r (which preserves the line element, see section 4 of [25] and section 4.1 of [27] for details). Hence, without loss of generality, from now on we can restrict to the case $h_{,u} = 0$. With (47), (43), (42), (40), (44), (45) this implies

$$h_{ij} = h_{ij}(x), \quad \mathcal{R} = \text{const}, \quad \mu = \text{const}, \quad \mathcal{E}^2 = \text{const}, \quad \mathcal{B}^2 = \text{const} \quad (n > 4), \quad (48)$$

so that H is a function of r only (cf. (26)) and (47) is satisfied identically. Up to a further (constant) coordinate rescaling, one can fix the normalization of \mathcal{R} such that $\mathcal{R} = 0, \pm(n-2)(n-3)$ [25, 27]. We also observe that (39) now implies $e_{i_1 \dots i_{p-2}} = e_{i_1 \dots i_{p-2}}(x)$.

3.5.2 Case $2p = n \pm 1$

For $2p = n \pm 1$ (n odd) there are additional terms of order r^{2-n} in (46) (due to the logarithmic terms in H , cf. (27), (28)) and (47) is replaced by

$$\mu_{,u} = -\frac{n-1}{n-2}(\ln \sqrt{h})_{,u} \left[\mu - \frac{\kappa_0}{2(n-2)} \mathcal{E}^2 \right] \quad (2p = n+1), \quad (49)$$

$$\mu_{,u} = -\frac{n-1}{n-2}(\ln \sqrt{h})_{,u} \left[\mu - \frac{2\kappa_0}{(n-1)^2(n-2)} \mathcal{B}^2 \right] \quad (2p = n-1). \quad (50)$$

However, here we still have $(\mathcal{E}^2)_{,i} = 0 = (\mathcal{B}^2)_{,i}$ (cf. (44), (45)), so that as in section 3.5.1 we can choose coordinates such that (48) holds (so that (49) and (50) are identically satisfied). In the end, the only difference in this case is thus the presence of the logarithmic terms in H .

3.5.3 Case $2p = n \pm 2$

Also for $2p = n \pm 2$ (n even) there are additional terms of order r^{2-n} in (46) (now due to (44), (45)) so that instead of (47) we obtain

$$\mu_{,u} = -\frac{n-1}{n-2} \mu (\ln \sqrt{h})_{,u} - \frac{\kappa_0 n}{(n-2)^2} \Delta(\mathcal{E}^2) \quad (n \geq 6, 2p = n+2), \quad (51)$$

$$\mu_{,u} = -\frac{n-1}{n-2} \mu (\ln \sqrt{h})_{,u} - \frac{2\kappa_0}{(n-2)^3} \Delta(\mathcal{B}^2) \quad (n \geq 6, 2p = n-2). \quad (52)$$

Recall that in the above two cases we generically have (eqs. (44) and (45)) $(\mathcal{B}^2)_{,i} = 0 \neq (\mathcal{E}^2)_{,i}$ (for $2p = n+2$) and $(\mathcal{E}^2)_{,i} = 0 \neq (\mathcal{B}^2)_{,i}$ (for $2p = n-2$). The argument of section 3.5.1 leading to $h(u, x) = U(u)X(x)$ (and then to $h_{ij} = h_{ij}(x)$) will thus still work in both cases provided $\mathcal{B}^2 \neq 0$ (for $2p = n+2$) or $\mathcal{E}^2 \neq 0$ (for $2p = n-2$). When these conditions are met, in suitable coordinates we thus arrive at (cf. a more detailed discussion in section 4.3 of [27] for the particular case $n = 6, p = 2$)

$$h_{ij} = h_{ij}(x), \quad \mathcal{R} = \text{const}, \quad \mu_{,u} = -\frac{\kappa_0 n}{(n-2)^2} \Delta(\mathcal{E}^2), \quad \mathcal{B}^2 = \text{const} \neq 0 \quad (2p = n+2), \quad (53)$$

$$h_{ij} = h_{ij}(x), \quad \mathcal{R} = \text{const}, \quad \mathcal{E}^2 = \text{const} \neq 0, \quad \mu_{,u} = -\frac{2\kappa_0}{(n-2)^3} \Delta(\mathcal{B}^2) \quad (2p = n-2). \quad (54)$$

In both cases we have $\mu_{,i} = 0$ and $(\mathcal{E}^2)_{,u} = 0 = (\mathcal{B}^2)_{,u}$ (see (40)), so that ∂_u is a Killing vector (twisting iff $H_{,i} \neq 0$). With the above equations this implies that $\mu_{,u}$ and $\Delta(\mathcal{E}^2)$ (respectively, $\Delta(\mathcal{B}^2)$) are both constants. If the transverse space with metric h_{ij} is assumed to be *compact* (as for

black hole spacetimes), then it follows (see, e.g., [42–44]) that \mathcal{E}^2 (respectively, \mathcal{B}^2) and thus μ are constant, and we thus again arrive at (48) (and thus $H = H(r)$).

However, in the special case $2p = n + 2$ with $\mathcal{B}^2 = 0 (\neq \mathcal{E}^2)$ (or $2p = n - 2$ with $\mathcal{E}^2 = 0 (\neq \mathcal{B}^2)$) one cannot in general conclude that $h(u, x) = U(u)X(x)$, and one has to consider the general equations (51), (52). If $h(u, x)$ is indeed non-factorized, then this metric describes an Einstein space that admits a conformal (non-homothetic) map on Einstein spaces [26, 27] and it is thus further constrained [45] (in particular, it must be of constant curvature when $n = 6$ and $p = 4$ or $p = 2$ in the electric and magnetic case, respectively [27]). One can still normalize $\mathcal{R} = 0, \pm(n - 2)(n - 3)$.

4 Summary for the generic case $2p \neq n$ ($n > 4$): static black holes

4.1 Case $2p \neq n \pm 2$: static black holes

4.1.1 Metric and Maxwell field

Keeping in mind also the concluding observations of section 3.5.1, the line-element is given by (cf. (5) with (25))

$$ds^2 = r^2 h_{ij} dx^i dx^j - 2 du dr - 2H du^2, \quad (55)$$

where $h_{ij} = h_{ij}(x)$ is the metric of a Riemannian Einstein space of dimension $(n - 2)$ and scalar curvature $\mathcal{R} = K(n - 2)(n - 3)$, and ((26) with (48))

$$2H = K - \frac{2\Lambda}{(n - 1)(n - 2)} r^2 - \frac{\mu}{r^{n-3}} + \frac{\kappa_0}{n - 2} \left[\frac{p - 1}{n + 1 - 2p} \frac{\mathcal{E}^2}{r^{2(n-p-1)}} - \frac{1}{p(n - 1 - 2p)} \frac{\mathcal{B}^2}{r^{2(p-1)}} \right] \quad (2p \neq n \pm 1), \quad (56)$$

where $\Lambda, \mu, \mathcal{E}^2, \mathcal{B}^2$ and $K = 0, \pm 1$ are all constants. The metric thus always admits the Killing vector field ∂_u , and it is static in regions where $H > 0$, while roots of $H(r)$ define Killing horizons (see also [31]). Recall that when $2p = n \pm 1$ (n odd), the second line of (56) should be replaced by (27) and (28), respectively.⁷

The ‘‘Coulombian’’ Maxwell field is given by (eqs. (7), (14), (15) and (36))

$$\mathbf{F} = \frac{1}{(p - 2)!} \frac{1}{r^{n+2-2p}} e_{i_1 \dots i_{p-2}}(x) du \wedge dr \wedge dx^{i_1} \wedge \dots \wedge dx^{i_{p-2}} + \frac{1}{p!} b_{i_1 \dots i_p}(x) dx^{i_1} \wedge \dots \wedge dx^{i_p}, \quad (57)$$

where $e_{i_1 \dots i_{p-2}}$ and $b_{i_1 \dots i_p}$ are harmonic forms (of respective rank $(p - 2)$ and p) in the transverse geometry h_{ij} , i.e., they obey the Euclidean source-free Maxwell equations in $(n - 2)$ dimensions (cf. (19) and the first and third of (35)). These forms are, however, further constrained by the conditions (31), (32) on the (constant) ‘‘square’’ of the field strengths, i.e.,

$$\mathcal{E}_{ij}^2 = \frac{\mathcal{E}^2}{n - 2} h_{ij} \quad (p \geq 3), \quad \mathcal{B}_{ij}^2 = \frac{\mathcal{B}^2}{n - 2} h_{ij} \quad (p \leq n - 2). \quad (58)$$

It is worth emphasizing again that conditions (58) do not only constrain the Maxwell field, but also impose severe restrictions on the Einstein metric h_{ij} . For instance, when $p = 2$ it must also be almost-Kähler if $\mathcal{B}^2 \neq 0$ and n must be even [27]. See [31] and appendix B for further comments.

⁷However, it can be seen that for $n = 5, p = 3$ and $n = 7, p = 4$ necessarily $\mathcal{E}^2 = 0$, and for $n = 5, p = 2$ and $n = 7, p = 3$ necessarily $\mathcal{B}^2 = 0$ (see appendix B) so that the logarithmic terms in (27) and (28) disappear in those cases. Therefore, effectively, logarithmic terms in both (27) and (28) can possibly arise only for an odd $n \geq 9$.

The above solutions can be seen as an extension to arbitrary p of the $p = 2$ ($n \neq 6$) solutions studied in [27] (including, when $\mathcal{B}^2 = 0$, the higher-dimensional Reissner-Nordström solution found by Tangherlini [4]) and possess similar qualitative features. In particular, they represent static black holes (at least for certain values of the parameters in (56)) dressed with electric and magnetic Maxwell fields. These solutions were previously obtained (starting from a static ansatz) and analyzed (including their thermodynamics) in [31], so that a detailed discussion is not necessary here. Recalling that in an $(n-2)$ -dimensional compact Riemannian space of positive constant curvature there exist no non-zero harmonic forms (except for a 0-form or a $(n-2)$ -form) [42, 43], we observe that the metric h_{ij} in (55) cannot describe a round sphere, except when $p = 2$ and $b_{i_1 i_2} = 0$ or, dually, when $p = n-2$ and $e_{i_1 \dots i_{n-4}} = 0$. Therefore, these static black holes cannot have a spherical horizon and cannot be asymptotically flat (in agreement with [7–9]), except in the electric $p = 2$ (or magnetic $p = n-2$) Reissner-Nordström solution of [4]. A flat and compact horizon metric h_{ij} is instead permitted (giving $K = 0$ in (56)), in which case the harmonic forms $e_{i_1 \dots i_{p-2}}$ and $b_{i_1 \dots i_p}$ must be covariantly constant [42, 43]. This allows for, e.g., asymptotically locally (A)dS black holes (see also [31]).

Similarly as for the case $p = 2$ [27], an additional “Vaidya-type” matter field representing pure radiation aligned with \mathbf{k} (i.e., adding an extra term to the component T_{uu} only) can easily be included by appropriately [27] modifying (47) and thus allowing for $\mu_{,u} \neq 0$ (see also a comment at the end of section 6 of [31]) — in the special case $2p = n$ this pure radiation can be sourced by the Maxwell field itself, as we show below in section 5.

4.1.2 Weyl and Maxwell types

As discussed in [27], the warped product structure of the metric (55) with $H = H(r)$ implies [28, 40] (see also [46] for a compact summary of these results) that the corresponding Weyl type is D(bd) and that

$$\mathbf{k} = \partial_r, \quad \mathbf{l} = -\partial_u + H\partial_r \quad (59)$$

are the (unique) double WANDs (no other WANDs — not even single ones — exist since the type is D(bd), cf. appendix A.3, in particular footnote 14; \mathbf{l} also defines a second Robinson–Trautman null direction, as follows by “time-reflection” symmetry [27, 28]). Since the type is D(bd), all the Weyl components are uniquely determined from (eqs. (A12)–(A14) of appendix (A) with (56))

$$\begin{aligned} \Phi &= -(n-2)(n-3) \frac{\mu}{2r^{n-1}} + \kappa_0 \frac{n-3}{(n-1)(n-2)} \\ &\quad \times \left[\frac{(n-p)(2n-2p-1)(p-1)}{n+1-2p} \frac{\mathcal{E}^2}{r^{2(n-p)}} - \frac{2p-1}{n-1-2p} \frac{\mathcal{B}^2}{r^{2p}} \right] \quad (2p \neq n \pm 1), \\ \tilde{\Phi}_{\hat{i}\hat{j}\hat{k}\hat{l}} &= r^{-2} \mathcal{C}_{\hat{i}\hat{j}\hat{k}\hat{l}}, \end{aligned} \quad (60)$$

where $\mathcal{C}_{\hat{i}\hat{j}\hat{k}\hat{l}}$ are the components of the Weyl tensor associated with h_{ij} in a frame of h_{ij} , and the notation of [10, 46] is employed (with a hat denoting components in a frame of the full spacetime metric $g_{\mu\nu}$, cf. appendix A.3). Since h_{ij} must be Einstein, $\mathcal{C}_{\hat{i}\hat{j}\hat{k}\hat{l}} = 0$ iff h_{ij} is of constant curvature (which is necessarily the case when $n = 5$), in which case the Weyl type becomes D(bcd). The scalar invariant $C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} = 4\Phi^2(n-1)/(n-3) + r^{-4} \mathcal{C}_{\hat{i}\hat{j}\hat{k}\hat{l}} \mathcal{C}_{\hat{i}\hat{j}\hat{k}\hat{l}}$ (cf. eqs. (69) and (70) of [47]) also signals a curvature singularity at $r = 0$. When $2p = n \pm 1$, the equation replacing (60) can be obtained by recalling (27), (28) and using (A12).

\mathbf{k} and \mathbf{l} are manifestly also aligned null directions of the Maxwell field (57) (\mathbf{k} is such by construction, recall (7), and \mathbf{l} then due to (36)), which is thus also of type D. It follows that also the Ricci tensor is of (aligned) type D (as can be explicitly verified thanks to $R_{ui} = 0$ and $R_{uu} = 2HR_{ur}$, cf. sections 3.4 and 3.5.1) — apart from (59), no other Ricci aligned null directions exist, not even

single ones, as can be seen recalling that $R_{ij} \propto h_{ij}$. One can easily see that in a frame parallelly transported along \mathbf{k} (and adapted to (59), see appendix A.3) the electric and magnetic components of (57) fall off, respectively, with the monopole rate $1/r^{n-p}$ and $1/r^p$ (as one could expect from a study of test fields [11]).

Examples with $p = 2$ (or, dually, $p = n - 2$) are given, e.g., in [4, 27]. Several other examples have been constructed in [31]. Using a construction described in [31] (see appendix B for a brief summary) one can produce more solutions, as we now exemplify.

Example ($n = 11, p = 3(8)$) A magnetic solution with $n = 11, p = 3$ (or electric with $p = 8$ after dualization) and a direct product transverse space is given by

$$\begin{aligned} h_{ij} &= h_{ij}^{(1)} + h_{ij}^{(2)} + h_{ij}^{(3)}, & 2H &= K - \frac{\Lambda}{45}r^2 - \frac{\mu}{r^8} - \frac{\kappa_0}{108}\frac{\mathcal{B}^2}{r^4}, \\ F_{123} &= F_{456} = F_{789} = \frac{\mathcal{B}}{3\sqrt{2}}, \end{aligned} \quad (61)$$

where $h_{ij}^{(1)}, h_{ij}^{(2)}$ and $h_{ij}^{(3)}$ are the metrics of three 3-dimensional spaces of constant curvature parameterized, respectively, by the coordinates $(x_1, x_2, x_3), (x_4, x_5, x_6)$ and (x_7, x_8, x_9) , and all of them having (constant) sectional curvature $4K$ (with $K = 0, \pm 1$). For a suitable choice of the parameters this represents magnetic static black holes with a direct product horizon. (Of course this solution can be dualized to an electric form with $p = 8$.)

4.2 Exceptional case $2p = n \pm 2$ ($n \geq 6$, even)

These special ranks of \mathbf{F} also fall into the discussion of section 4.1 if the *additional* assumptions $(\mathcal{E}^2)_{,i} = 0$ (for $2p = n + 2$) or $(\mathcal{B}^2)_{,i} = 0$ (for $2p = n - 2$) are made (or if the transverse space is assumed to be compact and the electric and magnetic fields are both non-zero, cf. section 3.5.3), in which case they again describe static black holes, as in the $n = 6, p = 2$ (or $p = 4$ after dualization) example given by eq. (66) of [27] (with $D = 6$), or as in the following example.

Example ($n = 8, p = 5(3)$) A solution describing an electric field with $n = 8, p = 5$ (or a magnetic field with $p = 3$ after dualization) is

$$\begin{aligned} h_{ij} &= \delta_{ij}, & 2H &= -\frac{\Lambda}{21}r^2 - \frac{\mu}{r^5} - \frac{\kappa_0}{3r^4}2\mathcal{E}^2, \\ F_{ur123} &= F_{ur456} = \frac{\mathcal{E}}{2\sqrt{3}}. \end{aligned} \quad (62)$$

Note that here the electric field is r -independent since $2p = 2 + n$ (first term of (57)). These are locally AdS (for $\Lambda < 0$) electric static black holes with a flat horizon.

However, more general solutions are now permitted for $2p = n + 2$ with $\mathcal{B}^2 = 0, \mathcal{E}^2 = \mathcal{E}^2(u, x)$ or for $2p = n - 2$ with $\mathcal{E}^2 = 0, \mathcal{B}^2 = \mathcal{B}^2(u, x)$ (or for a non-compact transverse space). The line-element is again given by (55), but H , given by (26), can generically depend on all coordinates (and the metric is thus *non-static*, in general). The Einstein metric $h_{ij} = h_{ij}(u, x)$ is generically non-factorized, and thus further constrained by the property of admitting conformal maps on Einstein spaces [45]. The Maxwell field is still given by (57) and satisfies the source-free Maxwell equations in the transverse geometry h_{ij} , but \mathcal{E}^2 (or \mathcal{B}^2) can now depend on x and u (eqs. (40)); the form $e_{i_1 \dots i_{p-2}}$ (or $b_{i_1 \dots i_p}$) is still u -independent (eq. (39) with $n = 2p + 2$, and (37)). The remaining equations to be satisfied are (51), (52) (see section 3.5.3 for more comments). Because (57) holds also here, the Maxwell and Ricci tensors obviously possess the same algebraic properties as discussed in section 4.1 and are

thus still of type D, doubly aligned with both null vectors (59). The Weyl type is generically II(bd) (see section 3.2 and appendix (A.3)) and the Weyl components of b.w. 0 (where Φ is necessarily non-zero) are still given by (60). We observe that $H_{,i} \neq 0$ implies that the vector \mathbf{l} is non-geodesic (appendix A.3) and not a multiple WAND, as can be seen using the results of appendix A.3.3 with (26) and (51), (52) (and vice versa, i.e., if $H_{,i} = 0$ then \mathbf{l} is a geodesic multiple WAND, as observed in section 4.1.2 for the case $2p \neq n \pm 2$).

It should also be observed that the ranks $2p = n \pm 2$ are special (even in the subcase of static metrics) because contributions from the electric and magnetic terms to the energy-momentum components T_{ij} cancel out (thanks to (31), (32)) for $2p = n + 2$ and for $2p = n - 2$, respectively. Related to this, for even p there can exist forms $e_{i_1 \dots i_{p-2}}$ (for $2p = n + 2$) or $b_{i_1 \dots i_p}$ (for $2p = n - 2$) that are self-dual in the transverse geometry h_{ij} , cf. also appendix B.

5 Special case $2p = n$ (n even): black holes with electromagnetic radiation

As we observed, the rank satisfying $2p = n$ has special properties and needs to be studied separately. This is not so surprising since this is the unique rank for which the Maxwell equations are conformally invariant and admit self-dual solutions (for odd p) or a continuous duality symmetry (for even p), cf., e.g., [32, 34, 35, 48].

The results of sections 3.1 and 3.2 are still valid also in the case $2p = n$ (but recall (33)). Instead, in sections 3.3, 3.4 and 3.5.1 we assumed $2p \neq n$ and those results are modified as follows.

5.1 Remaining field equations for the case $2p = n$

5.1.1 Maxwell equations, step two

Certain terms in the Maxwell equations studied in section 3.3 combine since they have the same r -dependence, and the equations of section 3.3 are thus replaced by⁸

$$(\sqrt{h} b^{kj_1 \dots j_{p-1}})_{,k} = \frac{1}{2}(n-2)\sqrt{h} h^{i_1 j_1} \dots h^{i_{p-1} j_{p-1}} e_{[i_2 \dots i_{p-1}, i_1]}, \quad (63)$$

$$b_{i_1 \dots i_p, u} = \frac{n}{2} f_{[i_2 \dots i_p, i_1]}, \quad (64)$$

$$(\sqrt{h} e^{i_1 \dots i_{p-2}})_{,u} = (\sqrt{h} f^{ki_1 \dots i_{p-2}})_{,k}. \quad (65)$$

One also obtains an additional equation $(\sqrt{h} h^{kl} h^{i_1 j_1} \dots h^{i_{p-2} j_{p-2}} e_{[i_1 \dots i_{p-2}, k]})_{,l} = 0$ ($\Leftrightarrow \star d \star d e = 0$), which is however identically satisfied by virtue of (63) and the antisymmetry of $b_{i_1 \dots i_p}$.

Note that the r.h.s. of (63) acts as a ‘‘current’’ for the p -form $b_{i_1 \dots i_p}$ so that $b_{i_1 \dots i_p}$ and $e_{[i_2 \dots i_{p-1}, i_1]}$ no longer satisfy the Euclidean source-free Maxwell equations in the transverse space. A fundamental difference from the generic case $2p \neq n$ is that now the components $F_{ui_1 \dots i_{p-1}}$ (eq. (16)) can be non-zero. Instead of (40) we now obtain from (65) and (64) (with (34))

$$(n-2)(\mathcal{E}^2)_{,u} = -n\mathcal{E}^2(\ln \sqrt{h})_{,u} + (n-2)\frac{2}{\sqrt{h}} e_{i_1 \dots i_{p-2}}(\sqrt{h} f^{ki_1 \dots i_{p-2}})_{,k}, \quad (66)$$

$$(n-2)(\mathcal{B}^2)_{,u} = -n\mathcal{B}^2(\ln \sqrt{h})_{,u} + n(n-2)b^{i_1 i_2 \dots i_p} f_{[i_2 \dots i_p, i_1]}. \quad (67)$$

⁸For later purposes it may be useful to observe that in an index-free notation the first of these equations reads $\star d \star b = -de$, where \star is the Hodge dual in the transverse space of h_{ij} (not to be confused with the n -dimensional Hodge dual \star in the full spacetime geometry $g_{\mu\nu}$, defined in section 1).

5.1.2 Einstein equation for R_{ui}

Since in general $F_{ui_1\dots i_{p-1}} \neq 0$, also $F_{u\alpha_1\dots\alpha_{p-1}} F_i^{\alpha_1\dots\alpha_{p-1}} \neq 0$ and the Einstein equation (4) for R_{ui} thus becomes

$$R_{ui} = \frac{\kappa_0}{4} \left[-(n-2) \delta_i^k e^{j_1\dots j_{p-2}} - 2b_i^{kj_1\dots j_{p-2}} \right] \left[r^{1-n}(n-2)e_{[j_1\dots j_{p-2},k]} - 2r^{2-n}f_{kj_1\dots j_{p-2}} \right]. \quad (68)$$

Instead of (43)–(45) this now gives

$$\mu_{,i} = \kappa_0 \left[(n-2) \delta_i^k e^{j_1\dots j_{p-2}} + 2b_i^{kj_1\dots j_{p-2}} \right] f_{kj_1\dots j_{p-2}}, \quad (69)$$

$$\left(\mathcal{E}^2 + \frac{4}{n(n-2)} \mathcal{B}^2 \right)_{,i} = \frac{n-2}{2} \left[(n-2) \delta_i^k e^{j_1\dots j_{p-2}} + 2b_i^{kj_1\dots j_{p-2}} \right] e_{[j_1\dots j_{p-2},k]}, \quad (70)$$

while $(n-4)\mathcal{R}_{,i} = 0$ still holds.

5.1.3 Einstein equation for R_{uu}

When $2p = n$ this equation is different for various reasons. First, since generically $F_{ui_1\dots i_{p-1}} \neq 0$, the r.h.s of (46) is not zero but

$$\begin{aligned} & \kappa_0 h^{i_1 j_1} \dots h^{i_{p-1} j_{p-1}} \left[r^{-n} \frac{(n-2)^2}{4} e_{[i_2\dots i_{p-1}, i_1]} e_{[j_2\dots j_{p-1}, j_1]} \right. \\ & \left. + r^{2-n} f_{i_1\dots i_{p-1}} f_{j_1\dots j_{p-1}} - r^{1-n}(n-2) e_{[i_2\dots i_{p-1}, i_1]} f_{j_1\dots j_{p-1}} \right]. \end{aligned} \quad (71)$$

Further, we now generically have $\Delta\mathcal{E} \neq 0$, $\Delta\mathcal{B} \neq 0$, $\Delta\mu \neq 0$ (cf. section 5.1.2), and (40) are not satisfied (section 5.1.1), giving rise to further terms of order r^{-n} and r^{1-n} in R_{uu} (cf. the l.h.s. of (46)). Keeping (71) and all these terms in mind, instead of (47) one obtains (at order r^{2-n} , r^{-n} , r^{1-n} , respectively)

$$(n-2)\mu_{,u} = -(n-1)\mu(\ln\sqrt{h})_{,u} - 2\kappa_0 \mathcal{F}^2 \quad (n > 4), \quad (72)$$

$$\begin{aligned} \Delta\mu = \frac{\kappa_0}{2} \left[(n-2) \left(\mathcal{E}^2 + \frac{4}{n(n-2)} \mathcal{B}^2 \right)_{,u} + n \left(\mathcal{E}^2 + \frac{4}{n(n-2)} \mathcal{B}^2 \right) (\ln\sqrt{h})_{,u} \right] \\ + 2\kappa_0 (n-2) e_{[i_2\dots i_{p-1}, i_1]} f^{i_1\dots i_{p-1}}, \end{aligned} \quad (73)$$

$$\Delta \left(\mathcal{E}^2 + \frac{4}{n(n-2)} \mathcal{B}^2 \right) = (n-2)^2 h^{i_1 j_1} \dots h^{i_{p-1} j_{p-1}} e_{[i_2\dots i_{p-1}, i_1]} e_{[j_2\dots j_{p-1}, j_1]}, \quad (74)$$

where we have defined the r -independent quantity

$$\mathcal{F}^2 \equiv h^{i_1 j_1} \dots h^{i_{p-1} j_{p-1}} f_{i_1\dots i_{p-1}} f_{j_1\dots j_{p-1}} = f^{i_1\dots i_{p-1}} f_{i_1\dots i_{p-1}}. \quad (75)$$

We observe that (73) and (74) are identically satisfied by virtue of (69) and (70), respectively (after using (63)–(67)) — this will be understood from now on. Note that if $\mathcal{F}^2 = 0$ ($\Leftrightarrow f_{i_1\dots i_{p-1}} = 0$) then (69) gives $\mu_{,i} = 0$ and thus $\mu(\ln\sqrt{h})_{,ui} = 0$ by (72), which implies (as explained in section (3.5.1)) that if $\mu \neq 0$ one can rescale the coordinates so as to have $h_{ij} = h_{ij}(x)$, $\mathcal{R} = \text{const}$, $\mu = \text{const}$. Vice versa, $\mathcal{F}^2 \neq 0$ clearly requires that $\mu_{,u}$ and $(\ln\sqrt{h})_{,u}$ cannot both vanish simultaneously, so that ∂_u cannot be a Killing vector field (as opposed to the case of static black holes, cf. section 4.1.1). For $n = 4$ ($p = 2$) the l.h.s. of (72) contains an additional term $-\frac{1}{2}\Delta\mathcal{R}$ [12, 13] (cf. also appendix B of [27]⁹) and this argument does not apply.

⁹Eq. (B.13) of [27] contains a typo: its r.h.s. should read $8P^2(Q_{,1}\xi_1 + Q_{,2}\xi_2)$.

5.2 Summary and discussion

We first observe that static black hole solutions clearly exist also for the special rank $2p = n$ (see section 5.3 below), to which a discussion similar to that section 4.1 still applies. However, there exists now also a new family of solutions in the case $F_{u j_1 \dots j_{p-1}} \neq 0$. The Maxwell field is given by

$$\begin{aligned} \mathbf{F} = & \frac{1}{\left(\frac{n}{2} - 2\right)!} \frac{1}{r^2} e_{i_1 \dots i_{p-2}} du \wedge dr \wedge dx^{i_1} \wedge \dots \wedge dx^{i_{p-2}} + \frac{1}{\left(\frac{n}{2}\right)!} b_{i_1 \dots i_p} dx^{i_1} \wedge \dots \wedge dx^{i_p} \\ & + \frac{1}{2\left(\frac{n}{2} - 1\right)!} \left(-\frac{n-2}{r} e_{[i_2 \dots i_{p-1}, i_1]} + 2f_{i_1 \dots i_{p-1}} \right) du \wedge dx^{i_1} \wedge \dots \wedge dx^{i_{p-1}}. \end{aligned} \quad (76)$$

The forms $e_{i_1 \dots i_{p-2}}(u, x)$ and $b_{i_1 \dots i_p}(u, x)$ are generally not harmonic in the transverse space, but satisfy the “modified” Euclidean Maxwell equations in $(n-2)$ dimensions (19) and (63). In addition, they can depend on u , cf. eqs. (64), (65). The latter further tells us that the $(p-1)$ -form $f_{i_1 \dots i_{p-1}}(u, x)$ is also generically non-harmonic. Notice that the Maxwell equations do not specify the u -dependence of $f_{i_1 \dots i_{p-1}}$, which can be interpreted as a freedom in the choice of a “wave profile”.

The line-element is given by (55) with

$$2H = K + \frac{2(\ln \sqrt{h})_{,u}}{n-2} r - \frac{2\Lambda}{(n-1)(n-2)} r^2 - \frac{\mu}{r^{n-3}} + \frac{\kappa_0}{2} \left(\mathcal{E}^2 + \frac{4}{n(n-2)} \mathcal{B}^2 \right) \frac{1}{r^{n-2}}, \quad (77)$$

where \mathcal{E}^2 , \mathcal{B}^2 and μ are generically functions of u and x , cf. the corresponding equations (66), (67), (69), (70), (72). The metric $h_{ij}(u, x) = h^{1/(n-2)}(u, x) \gamma_{ij}(x)$ is again Einstein, with scalar curvature $\mathcal{R} = K(n-2)(n-3)$ (where $K = 0, \pm 1$). One further has the constraint (33).

If h_{ij} is taken to be the metric of a *compact* space, then by (74) necessarily [42–44] $e_{[i_2 \dots i_{p-1}, i_1]} = 0$ (since the r.h.s of (74) is non-negative), so that $\left(\mathcal{E}^2 + \frac{4}{n(n-2)} \mathcal{B}^2 \right)_{,i} = 0$ (cf. (70)), and $e_{i_1 \dots i_{p-2}}$ and $b_{i_1 \dots i_p}$ must both be harmonic (eqs. (19), (63)). In particular, it follows [42, 43] that h_{ij} *cannot describe a round sphere*, and no asymptotically flat spacetimes are thus to be found in this class of solutions (note that if $e_{i_1 \dots i_{p-2}} = 0 = b_{i_1 \dots i_p}$, as in section 5.4 below, the argument still applies since in that case it is $f_{i_1 \dots i_{p-1}}$ that must be harmonic, cf. (64), (65)).

These solutions also include the standard electrovac Robinson–Trautman solutions with $2p = n = 4$. However, in that case an extra term proportional to $\Delta \mathcal{R}$ is present in (72) [12, 13, 27].

5.2.1 Maxwell type and self-duality

The Maxwell field (76) is in general of type II (aligned with \mathbf{k} by construction) and, in a parallelly transported frame adapted to (59) (appendix A.3), it peels off as (in agreement with test-field results [11])

$$\mathbf{F} = \frac{\mathbf{N}}{r^{\frac{n}{2}-1}} + \frac{\mathbf{II}}{r^{\frac{n}{2}}}, \quad (78)$$

where the symbols \mathbf{II} and \mathbf{N} specify the algebraic type of the corresponding terms, with the “radiative” \mathbf{N} term proportional to $f_{i_1 \dots i_{p-1}}$. The Maxwell type becomes D when there exists a second null direction aligned with \mathbf{F} , i.e. (using a null rotation (A21)), iff the following equation admits a solution¹⁰ for the null rotation parameter $z_{\bar{i}_p} \equiv r^{-1} z_{\hat{i}_p}$

$$-\frac{n-2}{2} e_{[\bar{i}_2 \dots \bar{i}_{p-1}, \bar{i}_1]} + r f_{\bar{i}_1 \dots \bar{i}_{p-1}} = -\frac{n-2}{2} r e_{[\bar{i}_2 \dots \bar{i}_{p-1}, z_{\bar{i}_1}]} + r z_{\bar{i}_p} b_{\bar{i}_p \bar{i}_1 \dots \bar{i}_{p-1}}, \quad (79)$$

¹⁰After contraction with $z_{\bar{i}_1}$ eq. (79) gives for $n > 4$ (i.e., $p > 2$) the necessary condition $f_{\bar{i}_1 \dots \bar{i}_{p-1}} z_{\bar{i}_1} = 0$, which generically will not be satisfied, thus showing that the generic Maxwell type is indeed II here (while for $n = 4$ ($p = 2$) the Maxwell types II and D are always equivalent, cf., e.g., [13]).

see solution (86) below for an example. As a special case, this occurs (with $z_{\hat{i}_1} = 0$) when the frame vector \mathbf{l} (59) is aligned with the Maxwell field, i.e., iff $f_{i_1 \dots i_{p-1}} = 0 = e_{[i_2 \dots i_{p-1}, i_1]}$ (which implies that the radiative term vanishes). On the other hand, the Maxwell type is N iff \mathbf{k} is doubly aligned, i.e., $e_{i_1 \dots i_{p-2}} = 0 = b_{i_1 \dots i_p}$ — these special cases are discussed below in sections 5.3 and 5.4. We observe that for $n = 6$ ($p = 3$) only the Maxwell type N is possible (cf. appendix B.3). Let us further notice that the b.w. -2 component $8\pi T_{\mu\nu} l^\mu l^\nu = R_{\mu\nu} l^\mu l^\nu = R_{uu} - 2HR_{ur}$ (with (59) and $R_{rr} = 0$) of the energy-momentum tensor, representing the flux of electromagnetic energy along \mathbf{k} , equals expression (71) and is characterized by the leading term $\kappa_0 r^{2-n} \mathcal{F}^2$ — an invariant quantity taking the same value in any frame parallelly transported along \mathbf{k} (i.e., it is invariant under a null rotation (A21) of the frame (A11) with $z_{\hat{i}}$ independent of r , and under spins).

When p is *odd*, straightforward calculations show that the field (76) is *self-dual* (${}^* \mathbf{F} = \mathbf{F}$) iff $\mathbf{e} = \star \mathbf{b}$, $\star d\mathbf{e} = -d\mathbf{e}$ and $\star \mathbf{f} = -\mathbf{f}$ (recall the definition of \star in footnote 8), whereas self-duality is not possible for an even p [32] (in particular, the condition $\mathbf{e} = \star \mathbf{b}$ implies $p(p-1)\mathcal{E}^2 = \mathcal{B}^2$, and that the l.h.s. and r.h.s. of (33) must vanish separately). Examples satisfying these conditions (with $\mathcal{E}^2 = 0 = \mathcal{B}^2$) are given by (100) and (101) (under the conditions described there).

5.2.2 Weyl type

In general, the Weyl tensor is of type II(bd), see section 3.2 and appendix A.3. The nontrivial Weyl components read (cf. (A12)–(A14) with (77))

$$\Phi = -\frac{1}{2}(n-2)(n-3)\mu r^{1-n} + \frac{1}{4}n(n-3)\kappa_0 \left(\mathcal{E}^2 + \frac{4}{n(n-2)} \mathcal{B}^2 \right) r^{-n}, \quad (80)$$

$$\tilde{\Phi}_{\hat{i}\hat{j}\hat{k}\hat{l}} = r^{-2} \mathcal{C}_{\hat{i}\hat{j}\hat{k}\hat{l}}, \quad (81)$$

$$\Psi'_{\hat{i}} = m_{(\hat{i})}^k \frac{n-3}{n-2} \left[\frac{1}{2}(n-1)\mu_{,k} r^{1-n} - \frac{1}{4}n\kappa_0 \left(\mathcal{E}^2 + \frac{4}{n(n-2)} \mathcal{B}^2 \right)_{,k} r^{-n} \right], \quad (82)$$

$$\Omega'_{\hat{i}\hat{j}} = m_{(\hat{i})}^k m_{(\hat{j})}^l \left\{ -\frac{1}{2} \left(\mu_{||kl} - \Delta \mu \frac{h_{kl}}{n-2} \right) r^{1-n} + \frac{1}{4} \kappa_0 \left[\left(\mathcal{E}^2 + \frac{4}{n(n-2)} \mathcal{B}^2 \right)_{||kl} - \Delta \left(\mathcal{E}^2 + \frac{4}{n(n-2)} \mathcal{B}^2 \right) \frac{h_{kl}}{n-2} \right] r^{-n} \right\}, \quad (83)$$

where (83) *holds only for* $n > 4$ since the identity (A10) has been employed (see [13] for $n = 4$), and the terms appearing in (82) and (83) are determined by (69), (70) (and (73), (74)). Eqs. (80) and (81) (which coincide with (60) for $2p = n$) imply that \mathbf{k} can never be a triple (or quadruple) WAND, since this would require $\mu = \mathcal{E}^2 = \mathcal{B}^2 = \mathcal{C}_{\hat{i}\hat{j}\hat{k}\hat{l}} = 0$ (and thus $f_{i_1 \dots i_{p-1}} = 0$ thanks to (72)), leading to a vacuum spacetime of constant curvature — except for this trivial case, the Weyl types III, N and O are thus *forbidden*. The Weyl type D(bd) (or D(bcd)) is possible in special cases, for example for metrics (84) and (85) below, when \mathbf{l} is aligned with \mathbf{F} (section 5.3), or when \mathbf{k} is doubly aligned with \mathbf{F} (section 5.4). As in section 4.1.2, $C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} = 4\Phi^2(n-1)/(n-3) + r^{-4} \mathcal{C}_{\hat{i}\hat{j}\hat{k}\hat{l}} \mathcal{C}_{\hat{i}\hat{j}\hat{k}\hat{l}}$.

5.2.3 Examples

Some explicit examples of various Weyl and Maxwell types are provided below.

Example ($n = 8, p = 4$) It is easy to construct examples if the transverse space is taken to be flat, the forms $e_{i_1 \dots i_{p-2}}, b_{i_1 \dots i_p}$ and $f_{i_1 \dots i_{p-1}}$ to be harmonic — in fact x -independent — and $\mu = \mu(u)$

(with $(\mathcal{E}^2)_{,u} = 0 = (\mathcal{B}^2)_{,u}$ by (66), (67)). A simple example with $n = 8$, $p = 4$ is given by (55) and (76) with

$$\begin{aligned} h_{ij} &= \delta_{ij}, & 2H &= -\frac{\Lambda}{21}r^2 - \frac{\mu(u)}{r^5} + \frac{\kappa_0}{2r^6} \left(\mathcal{E}^2 + \frac{\mathcal{B}^2}{12} \right), & \mu(u) &= \mu_0 - \frac{\kappa_0}{3} \int \mathcal{F}^2 du, \\ F_{ur12} &= F_{ur34} = F_{ur56} = \frac{\mathcal{E}}{\sqrt{6}} \frac{1}{r^2}, & F_{1234} &= F_{1256} = F_{3456} = \frac{\mathcal{B}}{6\sqrt{2}}, \\ F_{uijk} &= f_{ijk}(u) & \text{with } F_{ui12} &= F_{ui34} = F_{ui56} = 0, \end{aligned} \quad (84)$$

where \mathcal{E} and \mathcal{B} are constants. This spacetime generically represents black holes with a flat horizon in the presence of an electromagnetic field which consists of both a static component and of u -dependent expanding radiation.¹¹ Correspondingly, the mass parameter μ (and thus the location of horizons) is u -dependent and monotonically increases or decreases according to the time-orientation of \mathbf{k} (thus corresponding to received or emitted radiation, cf. [21]). These metrics are asymptotically locally (A)dS if $\Lambda \neq 0$, but if $\mu > 0$ they clearly possess a horizon when $\Lambda = 0$, too. The Weyl type is D(bcd) because here $H = H(u, r)$ and h_{ij} is flat [28, 40, 46] (as can also be seen explicitly from (80)–(83)). Generically, the Maxwell field is of genuine type II. Here \mathcal{E} and \mathcal{B} are independent parameters and, in particular, can vanish independently (in other words, (31) and (32) are satisfied separately). However, since here $2p = n$, solutions exist that satisfy only the weaker constraint (33) (i.e., the electric and magnetic fields must both be non-zero), as illustrated by the following example.

Example ($n = 8$, $p = 4$) To obtain a different solution with $n = 8$, $p = 4$, one can specify the metric functions and electromagnetic field by

$$\begin{aligned} h_{ij} &= \delta_{ij}, & 2H &= -\frac{\Lambda}{21}r^2 - \frac{\mu(u)}{r^5} + \kappa_0 \frac{3\mathcal{E}^2}{4r^6}, & \mu(u) &= \mu_0 - \frac{\kappa_0}{3} \int \mathcal{F}^2 du, \\ F_{ur12} &= F_{ur34} = \frac{\mathcal{E}}{2} \frac{1}{r^2}, & F_{1234} &= \frac{\mathcal{E}}{2}, \\ F_{uijk} &= f_{ijk}(u) & \text{with } F_{ui12} &= 0 = F_{ui34}, \end{aligned} \quad (85)$$

where \mathcal{E} is a constant. Clearly (31) and (32) are not satisfied in this case. If $\mathcal{E} \neq 0 \neq f_{ijk}$ the Maxwell field is of genuine type II. Comments similar to those given for (84) apply also here, in particular the Weyl type is again D(bcd).

Example ($n = 8$, $p = 4$) An example in which μ has also a non-trivial x -dependence is given by

$$\begin{aligned} h_{ij} &= \delta_{ij}, & 2H &= -\frac{\Lambda}{21}r^2 - \frac{\mu(u, x_1)}{r^5} + \frac{\kappa_0 \mathcal{B}^2}{24r^6}, & \mu(u, x_1) &= \mu_0 - 2\kappa_0 f_0 (\sqrt{2}\mathcal{B}x_1 + 2f_0 u), \\ F_{1234} &= F_{1256} = F_{3456} = \frac{\mathcal{B}}{6\sqrt{2}}, & F_{u234} &= F_{u256} = f_0, \end{aligned} \quad (86)$$

where \mathcal{B} and f_0 are constants (for $f_0 = 0$ this trivially reduces to a subcase of (84)). Here $\Phi \neq 0 \neq \Psi'_i$ while $\tilde{\Phi}_{\hat{i}\hat{j}\hat{k}\hat{l}} = 0 = \Omega'_{\hat{i}\hat{j}}$ (see (80)–(83)), so that the Weyl tensor is of genuine type II(bcd) (by the argument in appendix A.3.4, since here (A24) admits no solutions). The vector field \mathbf{l} is clearly not aligned with \mathbf{F} , nevertheless the Maxwell type is D, a second aligned null direction being given by a null rotation (A21) with $\mathbf{m}_{(\hat{1})} = r^{-1}\partial_{x_1}$ and the only non-zero parameter $z_{\hat{1}} = -6\sqrt{2}K_0\mathcal{B}^{-1}r$ (cf. (79)).

¹¹It should be observed that no direct four-dimensional analog of such solutions exists since in 4D the presence of non-vanishing b.w. 0 components of the Maxwell field would imply (by (69)) that $\mu_{,i} \neq 0$ (cf. (28.37e) of [13], or (B.11) of [27]).

5.3 l aligned with F : static black holes ($n \geq 8$)

The null vector field l of (59) is uniquely defined by certain geometrical properties (appendix A.3). Furthermore, as noticed above, it is aligned with F iff

$$f_{i_1 \dots i_{p-1}} = 0, \quad e_{[i_2 \dots i_{p-1}, i_1]} = 0, \quad (87)$$

which implies that the Maxwell field is of type D and non-radiative (as mentioned above and in appendix B.3, for $n = 6$ this would lead to $F = 0$, so we can restrict here to $n \geq 8$) — indeed here $T_{\mu\nu} l^\mu l^\nu = 0$. Equations (63), (64), (65), (66), (67), (69) and (72) now take exactly the same form as in the generic case $2p \neq n$ (in particular, (66), (67) reduce to (40)). Instead of the conditions $\mathcal{E}_{,i}^2 = 0 = \mathcal{B}_{,i}^2$ found in the $2p \neq n$ case, one has the weaker condition (from (70))

$$\left(\mathcal{E}^2 + \frac{4}{n(n-2)} \mathcal{B}^2 \right)_{,i} = 0. \quad (88)$$

However, together with (40) this suffices to show that, again, one can rescale away the u -dependence of h_{ij} (and thus also of μ and $e_{i_1 \dots i_{p-2}}$), and the discussion of section 4.1 then applies (with the only difference that (58) are replaced by (33), and that (88) replaces $\mathcal{E}_{,i}^2 = 0 = \mathcal{B}_{,i}^2$). The metric is thus (55) with $H(r)$ given by (77) but without the $\frac{2}{n-2}(\ln \sqrt{h})_{,u} r$ term (all the coefficients of the powers of r appearing in H are indeed constants) and represents static black holes. Again it follows that the Weyl type is D(bd) (possibly, D(bcd)) and l is also a double WAND. The Maxwell field is given by (57) with $2p = n$. Here (78) clearly reduces to $F = D r^{-\frac{n}{2}}$. These $2p = n$ solutions with F of type D were previously studied in [31]. Two examples with $n = 8$, $p = 4$ are given by (84) and (85) with $f_{ijk}(u) = 0$ (and thus $\mu = \text{const}$), see [31] for others.

5.4 Type N Maxwell field

The field (76) is of type N iff

$$e_{i_1 \dots i_{p-2}} = 0, \quad b_{i_1 \dots i_p} = 0, \quad (89)$$

with k being the unique aligned null direction, so that

$$F = \frac{1}{\left(\frac{n}{2} - 1\right)!} f_{i_1 \dots i_{p-1}} du \wedge dx^{i_1} \wedge \dots \wedge dx^{i_{p-1}}, \quad (90)$$

and the peeling (78) becomes simply $F = N r^{1-\frac{n}{2}}$.

In this case eqs. (63)–(72) reduce to

$$f_{[i_2 \dots i_p, i_1]} = 0, \quad (\sqrt{h} f^{k i_1 \dots i_{p-2}})_{,k} = 0, \quad (91)$$

$$\mu_{,i} = 0, \quad (92)$$

$$(n-2)\mu_{,u} = -(n-1)\mu(\ln \sqrt{h})_{,u} - 2\kappa_0 \mathcal{F}^2 \quad (n > 4). \quad (93)$$

Eqs. (91) mean that $f_{i_1 \dots i_{p-1}}(u, x)$ effectively defines a Maxwell $\left(\frac{n}{2} - 1\right)$ -form in the $(n-2)$ -dimensional transverse space (i.e., it is harmonic).

The line-element is (55), where $h_{ij}(u, x) = h^{1/(n-2)}(u, x) \gamma_{ij}(x)$ is Einstein and with $H(u, r, x)$ given by

$$2H = K + \frac{2(\ln \sqrt{h})_{,u}}{n-2} r - \frac{2\Lambda}{(n-1)(n-2)} r^2 - \frac{\mu(u)}{r^{n-3}}. \quad (94)$$

The term $(\ln \sqrt{h})_{,u}$ can be reexpressed in terms of μ and \mathcal{F}^2 using (93), if desired. Note that by (93) necessarily $\mu \neq 0$.¹² Since $f_{i_1 \dots i_{p-1}}$ is harmonic, the metric h_{ij} cannot be of positive constant curvature [42, 43].

Here the energy-momentum tensor possesses only the (b.w. -2) component describing a flux of radiation along \mathbf{k}

$$8\pi T_{\mu\nu} l^\mu l^\nu = 8\pi T_{uu} = \kappa_0 r^{2-n} \mathcal{F}^2, \quad (95)$$

and thus the Maxwell form acts as an aligned *pure radiation* field. Therefore these special Robinson–Trautman metrics are contained in the pure radiation family of solutions studied in [25] (but the corresponding Maxwell equations were not considered there). In particular, from appendix A of [25] it immediately follows that the Weyl type is D(bd) (D(bcd) if h_{ij} is of constant curvature), with (59) being the two multiple WANDs and

$$\Phi = -(n-2)(n-3) \frac{\mu(u)}{2r^{n-1}} \neq 0, \quad \tilde{\Phi}_{\hat{i}\hat{j}\hat{k}\hat{l}} = r^{-2} \mathcal{C}_{\hat{i}\hat{j}\hat{k}\hat{l}}, \quad (96)$$

as can also be seen explicitly from (80)–(83) with $\mathcal{E}^2 = 0 = \mathcal{B}^2$.

5.4.1 A special subcase: solutions with a factorized $h(u, x) = U(u)X(x)$

In the special case of a factorized $h(u, x) = U(u)X(x)$, one can set $U(u) = 1$ by a coordinate transformation, and thus obtain a special subclass of solutions with metric (55) with $h_{ij} = h_{ij}(x)$ and $H(u, r)$ given by

$$2H = K - \frac{2\Lambda}{(n-1)(n-2)} r^2 - \frac{\mu(u)}{r^{n-3}}, \quad (97)$$

$$\mu(u) = \mu_0 - \frac{2\kappa_0}{n-2} \int \mathcal{F}^2 du. \quad (98)$$

Here necessarily $\mu_{,u} \neq 0$. The Maxwell field is given by (90), where $f_{i_1 \dots i_{p-1}}$ must satisfy (91) and, by (93), also

$$(\mathcal{F}^2)_{,i} = 0. \quad (99)$$

These solutions will in general describe black holes in the presence of electromagnetic radiation with non-zero expansion, with a monotonically increasing (or decreasing, according to the choice of time-orientation of \mathbf{k}) mass parameter μ .

A simple solution can be obtained by taking the transverse metric to be flat, i.e., $h_{ij} = \delta_{ij}$ and $f_{i_1 \dots i_{p-1}} = f_{i_1 \dots i_{p-1}}(u)$ (obviously with such a choice (91) and (99) are identically satisfied). To be specific, we give the following explicit example for $n = 6$.

Example ($n = 6, p = 3$) The metric and the Maxwell field are

$$\begin{aligned} h_{ij} &= \delta_{ij}, & 2H &= -\frac{\Lambda}{10} r^2 - \frac{\mu(u)}{r^3}, & \mu(u) &= \mu_0 - \frac{\kappa_0}{2} \int \mathcal{F}^2 du, \\ F_{uij} &= f_{ij}(u). \end{aligned} \quad (100)$$

For $\Lambda < 0$ this spacetime represents asymptotically locally AdS black holes with electromagnetic radiation. By a rotation, one can always simplify the Maxwell field so as to only have non-zero

¹²It is worth observing that this is not true in 4D, due precisely to the additional term proportional to $\Delta\mathcal{R}$ entering (93) when $n = 4$. Thanks to this extra term, 4D solutions of Petrov type III (with $\mu = 0$ and a null Maxwell field) are thus possible, see section 28.2.2 of [13] and references therein.

components $F_{u12} = f_{12}(u)$, $F_{u34} = f_{34}(u)$, in which case \mathbf{F} is self-dual when $f_{34}(u) = -f_{12}(u)$ (or anti-self-dual when $f_{34}(u) = +f_{12}(u)$), cf. section 5.2. Solution (100) is an extension of a solution given in 4D (for $\Lambda = 0$) in [12] (also reproduced in eq. (28.43) of [13]) and recently discussed in [21] (see also footnote 1). It should be observed that for $n = 6$ this example in fact comprises all the possible Robinson–Trautman solutions with a Maxwell field of type N *if* the transverse space is assumed to be of constant curvature (this follows from (91) and (99) after using coordinates adapted to the constant curvature space, e.g., those employed in [25] — it also follows that the (constant) curvature must necessarily be zero). A similar example with $n = 8$, $p = 4$ is given by (84) with $\mathcal{E} = 0 = \mathcal{B}$. An $n = 6$ example where, instead, h_{ij} is not of constant curvature follows.

Example ($n = 6$, $p = 3$) As another example in 6D one can take h_{ij} to be a direct product of two S^2 or H^2 , namely

$$\begin{aligned} h_{ij} dx^i dx^j &= [1 - 3Kx_1^2]^{-1} dx_1^2 + [1 - 3Kx_1^2] dx_2^2 + [1 - 3Kx_3^2]^{-1} dx_3^2 + [1 - 3Kx_3^2] dx_4^2, \\ 2H &= K - \frac{\Lambda}{10} r^2 - \frac{\mu(u)}{r^3}, \quad K = \pm 1, \quad \mu(u) = \mu_0 - \kappa_0 \int (f_{12}^2(u) + f_{34}^2(u)) du, \\ F_{u12} &= f_{12}(u), \quad F_{u34} = f_{34}(u). \end{aligned} \quad (101)$$

As above, (anti-)self-duality holds iff $f_{34}(u) = \mp f_{12}(u)$. For $K = 0$ this solutions reduces to (100) (up to a space rotation).

Acknowledgments

M.O. and M.Ž. have been supported by the Albert Einstein Center for Gravitation and Astrophysics, Czech Science Foundation GAČR 14-37086G. M.O. has also been supported by research plan RVO: 67985840. J.P. has been supported by the research grant GAČR P203/12/0118.

A Robinson–Trautman spacetimes with an aligned Ricci tensor of type II

For the purposes of the present paper and for possible future reference it is useful to summarize some of the results of [25, 27, 29, 30] in the present appendix. Note that we restrict here to the Robinson–Trautman spacetimes in which the Ricci tensor is of type II aligned with the privileged vector field \mathbf{k} . This includes vacuum solutions as well as solutions with aligned matter content (as we assume in the main text).

A.1 General metric

The line-element and its Weyl type are specified by the following theorem.

Theorem A.1. *If a n -dimensional spacetime ($n \geq 4$) admits a non-twisting, non-shearing, expanding geodesic null vector field \mathbf{k} and the Ricci tensor is of aligned type II, adapted coordinates $(u, r, x^1, \dots, x^{n-2})$ can be chosen such that [25]*

$$ds^2 = r^2 h_{ij} (dx^i + W^i du) (dx^j + W^j du) - 2 du dr - 2H du^2, \quad (A1)$$

$$h_{ij} = h_{ij}(u, x), \quad W^i = \alpha^i(u, x) + r^{1-n} \beta^i(u, x), \quad (A2)$$

$$\mathbf{k} = \partial_r, \quad \theta = 1/r, \quad (A3)$$

where H is an arbitrary function of all coordinates. \mathbf{k} is automatically a WAND, such that the Weyl tensor is in general of aligned type I(b). It is a multiple WAND iff $\beta^i = 0$ [30], in which case the Weyl tensor is of aligned type II(d) (or more special). When $\beta^i = 0$, one can locally set $W^i = 0$ (after a coordinate transformation giving $\alpha^i = 0$) [25]. The Weyl type further specializes to II(bd) iff h_{ij} is an Einstein metric (with still $W^i = 0$) [30].

The vector field \mathbf{k} is the generator of null hypersurfaces $u = \text{const}$ such that $k_\mu dx^\mu = -du$, r is an affine parameter along \mathbf{k} , θ is its expansion scalar, and $x \equiv (x^i) \equiv (x^1, \dots, x^{n-2})$ are spatial coordinates on a “transverse” $(n-2)$ -dimensional Riemannian manifold. We observe that the condition that the Ricci tensor is doubly aligned with \mathbf{k} can be expressed in a frame-independent form as $R_{\mu\nu}k^\nu = \alpha k_\mu$, which in the coordinates of (A1) means $R_{rr} = 0 = R_{ri}$. For certain calculations it may be useful to observe that $2H = g^{rr} = -g_{uu}$ and $W^i = g^{ri}$ (such that $W^i = 0 \Leftrightarrow g^{ri} = 0 \Leftrightarrow g_{ui} = 0$).

The first part of the theorem was proven in sections 3.1 and 3.2 of [25]. The results on Weyl alignment follow immediately from eqs. (13)–(15), (17) and (19) (with (24), (25)) of [30] together with (A1).¹³ In vacuum or with aligned pure radiation (i.e., the only non-zero energy-momentum tensor component being T_{uu}) necessarily $\beta^i = 0$ and h_{ij} is Einstein for any $n \geq 4$, and the Weyl tensor further specializes to type D(bd) (possibly, D(bcd) or D(acd)) if $n > 4$ [25] (the latter result was also rederived in [30] without the coordinate choice $\alpha^i = 0$), the type O being possible only in the trivial case of constant curvature spacetimes. In [29] it was shown that for any $n \geq 4$ Robinson–Trautman spacetimes cannot support aligned gyratonic matter (i.e., an energy-momentum tensor with *both* T_{uu} and T_{ui} — and no other components — being non-zero). Let us finally observe that if one relaxes the assumptions of the theorem by requiring only the aligned Ricci type I (i.e., $R_{rr} = 0$), one obtains the same form of the metric, except that the $W^i(u, r, x)$ are arbitrary functions [25] (the Weyl type remains I(b) [30]).

A.2 Ricci tensor components

The Ricci tensor component R_{ij} for the general metric (A1), (A2) was given explicitly in eq. (A.1) of [27] and reads

$$\begin{aligned} R_{ij} = & \mathcal{R}_{ij} - r^{4-n} (r^{n-3} 2H)_{,r} h_{ij} - r^{2(2-n)} \frac{(n-1)^2}{2} h_{ik} h_{jl} \beta^k \beta^l \\ & - r \left[\frac{n-2}{2} \left(2h_{k(i} \alpha^k_{,j)} + \alpha^k h_{ij,k} - h_{ij,u} \right) + \left(\alpha^k_{,k} + \alpha^k (\ln \sqrt{h})_{,k} - (\ln \sqrt{h})_{,u} \right) h_{ij} \right] \\ & + r^{2-n} \left[\frac{1}{2} \left(2h_{k(i} \beta^k_{,j)} + \beta^k h_{ij,k} \right) - \left(\beta^k_{,k} + \beta^k (\ln \sqrt{h})_{,k} \right) h_{ij} \right], \end{aligned} \quad (\text{A4})$$

where \mathcal{R}_{ij} is the Ricci tensor associated with the spatial metric h_{ij} (as defined in section 3.2), and a partial derivative w.r.t. (e.g.) x^j is simply denoted by a comma followed by j . For the purposes of the present paper we need the remaining Ricci components only in the special case $W^i = 0$, which we now present (but see [29,30] for the Ricci tensor components of the most general Robinson–Trautman metric, i.e., without not even enforcing $R_{rr} = 0 = R_{ri}$).

¹³Recall that the Weyl type I(b) in fact characterizes *all* Robinson–Trautman geometries, independently of any assumptions on the Ricci tensor [30]. This can also be seen from the Ricci identities (11g) and (11k) (and their trace) of [49] using the fact that \mathbf{k} is geodesic, shearfree and twistfree.

A.2.1 Case $W^i = 0$ (\mathbf{k} is a multiple WAND — Weyl type II(bd))

If one further assumes $W^i = 0$ in (A1) (i.e., \mathbf{k} is a multiple WAND, as we indeed find in section 3.2), eq. (A4) and the remaining Ricci components, given in eqs. (26), (27) and (31) of [25], reduce to

$$R_{ij} = \mathcal{R}_{ij} - r^{4-n} (r^{n-3} 2H)_{,r} h_{ij} + r \left[\frac{n-2}{2} h_{ij,u} + (\ln \sqrt{h})_{,u} h_{ij} \right], \quad (\text{A5})$$

$$R_{ur} = r^{2-n} (r^{n-2} H_{,r})_{,r} - r^{-1} (\ln \sqrt{h})_{,u}, \quad (\text{A6})$$

$$R_{ui} = r^{4-n} (r^{n-4} H_{,i})_{,r} + \frac{1}{2} (h^{jk} h_{ik,u})_{,j} + \frac{1}{2} h^{jk} h_{ik,u} (\ln \sqrt{h})_{,j} - \frac{1}{4} h^{jk} h^{lm} h_{kl,u} h_{jm,i} - (\ln \sqrt{h})_{,ui}, \quad (\text{A7})$$

$$R_{uu} = 2HR_{ur} - r^2 (r^{-2} H)_{,r} (\ln \sqrt{h})_{,u} + (n-2)r^{-1} H_{,u} + r^{-2} \Delta H - (\ln \sqrt{h})_{,uu} - \frac{1}{4} h^{il} h^{jk} h_{ij,u} h_{kl,u}, \quad (\text{A8})$$

where in the last expression the first quantity on the r.h.s. has been written in terms of (A6) for brevity and for convenience in the calculations of section 3.5.1, and $\Delta \equiv \frac{1}{\sqrt{h}} \partial_j (\sqrt{h} h^{ij} \partial_i)$ is the Laplace operator in the $(n-2)$ -dimensional space with metric h_{ij} . Note that $2(n-2)R_{ur} = -r^{-1} (h^{ij} R_{ij})_{,r}$. For general purposes it is useful to observe that the component R_{ui} can be rewritten more compactly as $R_{ui} = r^{4-n} (r^{n-4} H_{,i})_{,r} + \frac{1}{2} h^{kl} (h_{ki,u||l} - h_{kl,u||i})$ [29], where the lower double bar $||$ denotes a covariant derivative w.r.t. h_{ij} — but in the computations of the present paper the form (A7) can be used more readily.

A.3 Weyl tensor components in the case $R_{ij} \propto h_{ij}$ ($\Rightarrow W^i = 0$)

For the purposes of the present paper, we further restrict here to the case when the condition $R_{ij} \propto h_{ij}$ holds for the metric (A1) (this is also true in various other cases of physical interest, e.g., in vacuum, or whenever the traceless Ricci tensor is of aligned type III or more special), while we refer to [30] for the Weyl tensor components of the most general Robinson–Trautman metric. Similarly as in section 3.2, together with (A4) this restriction gives $\beta^i = 0$, and thus $W^i = 0$ (theorem A.1). By (A5) we further have

$$\mathcal{R}_{ij} = \frac{\mathcal{R}}{n-2} h_{ij}, \quad h_{ij,u} = \frac{2(\ln \sqrt{h})_{,u}}{n-2} h_{ij}, \quad (\text{A9})$$

so that, in particular, h_{ij} is Einstein and therefore $(n-4)\mathcal{R}_{,i} = 0$. (Note that, indeed, these conditions are obtained in section 3.2 as a consequence of the Einstein equations — except in the cases $p = 1, n = 1$, for which see appendix C.) Using (A9) one can further prove the identity (cf. eq. (144) of [30])

$$(n-4) \left[[(\ln \sqrt{h})_{,u}]_{||ij} - \frac{\Delta [(\ln \sqrt{h})_{,u}] h_{ij}}{n-2} \right] = 0, \quad (\text{A10})$$

needed in the following.

We choose a frame

$$\mathbf{k} = \partial_r, \quad \mathbf{l} = -\partial_u + H\partial_r, \quad \mathbf{m}_{(\hat{i})} = \frac{1}{r} m_{(\hat{i})}^j \partial_j, \quad (\text{A11})$$

where the functions $m_{(\hat{i})}^j$ do not depend on r . The spacelike vectors $\mathbf{m}_{(\hat{i})}$ span the transverse space of constant u and r , and obviously $\bar{\mathbf{m}}_{(\hat{i})} \equiv r\mathbf{m}_{(\hat{i})} = m_{(\hat{i})}^j \partial_j$ defines an orthonormal frame for the

metric h_{ij} . Before proceeding, it is useful to observe that since \mathbf{k} and \mathbf{l} span the 2-space (u, r) we have the relation $\tau_i - \tau'_i = 0$ (in the notation of [10, 46] for the Ricci rotation coefficients). Using (55), it immediately follows that \mathbf{l} is parallelly transported along \mathbf{k} , i.e., $\tau'_i = 0$, so that also $\tau_i = 0$. There cannot be other null directions \mathbf{l}' with these properties (as follows from the transformation properties of τ_i and τ'_i under null rotations [10, 49]). It is also easy to see that \mathbf{l} is geodesic iff $H_{,i} = 0$ [28]. In fact, not only \mathbf{l} but the full frame (A11) is parallelly transported along \mathbf{k} .

Recall that, thanks to $W^i = 0$ and the first of (A9), the Weyl type here is II(bd) or more special (Theorem A.1). In the frame (A11) one further finds $\tilde{\Psi}'_{\hat{i}\hat{j}\hat{k}} = 0$, so that all the non-zero components are determined by (using eqs. (16), (18), (20) and (22) of [30], or eqs. (A.1) of [25])

$$\Phi = \frac{n-3}{2(n-1)} \left\{ r^2 \left[r^{-2} \left(2H - \frac{\mathcal{R}}{(n-2)(n-3)} \right) \right]_{,r} \right\}, \quad \tilde{\Phi}_{\hat{i}\hat{j}\hat{k}\hat{l}} = r^{-2} \mathcal{C}_{\bar{i}\bar{j}\bar{k}\bar{l}}, \quad (\text{A12})$$

$$\Psi'_i = r m_{(i)}^k \frac{n-3}{n-2} \left[r^{-2} \left(H - \frac{(\ln \sqrt{h})_{,u}}{n-2} r \right) \right]_{,kr}, \quad (\text{A13})$$

$$\Omega'_{\hat{i}\hat{j}} = \frac{1}{r^2} m_{(i)}^k m_{(j)}^l \left(H_{||kl} - \frac{\Delta H}{n-2} h_{kl} \right), \quad (\text{A14})$$

where $\mathcal{C}_{\bar{i}\bar{j}\bar{k}\bar{l}}$ are the frame components of the Weyl tensor associated with h_{ij} (in the corresponding frame $\bar{\mathbf{m}}_{(\bar{i})}$). The above components are ordered by b.w. and expressed in the notation of [10, 46] (cf. table 2 of [46]). We observe that the alternative notation of [30] corresponds to $\Psi_{2S} = -\Phi$, $\tilde{\Psi}_{2^{ijk}l} = \tilde{\Phi}_{\hat{i}\hat{j}\hat{k}\hat{l}}$, $\Psi_{3T^i} = \Psi'_i$ and $\Psi_{4^{ij}} = \Omega'_{\hat{i}\hat{j}}$.

A.3.1 Conditions for the Weyl types II(abd), II(bcd) and III(b)

As noticed above, here the Weyl type is generically II(bd). It becomes II(abd) iff $\Phi = 0$, i.e. (using the first of (A12)), when

$$2H = \frac{\mathcal{R}}{(n-2)(n-3)} + c_1(u, x)r + c_2(u, x)r^2, \quad (\text{A15})$$

where c_1 and c_2 are integration functions independent of r (recall that \mathcal{R} is also independent of r and $(n-4)\mathcal{R}_{,i} = 0$, as noticed above).

On the other hand, the type becomes II(bcd) iff $\tilde{\Phi}_{\hat{i}\hat{j}\hat{k}\hat{l}} = 0$, namely (using the second of (A12)) for

$$\mathcal{C}_{\bar{i}\bar{j}\bar{k}\bar{l}} = 0, \quad (\text{A16})$$

which (with the first of (A9)) means that h_{ij} is a constant curvature metric.

The Weyl type further specializes to III(b) (recall that here we have $\tilde{\Psi}'_{\hat{i}\hat{j}\hat{k}} = 0$ identically) iff (A15) and (A16) hold simultaneously.

A.3.2 Conditions for the Weyl type N

The Weyl type becomes N when $\Phi = 0$, $\tilde{\Phi}_{\hat{i}\hat{j}\hat{k}\hat{l}} = 0$ and $\Psi'_i = 0$ which gives (using (A13) and (A15))

$$2H = \frac{\mathcal{R}}{(n-2)(n-3)} + \frac{2(\ln \sqrt{h})_{,u}}{n-2} r + c_2(u, x)r^2, \quad (\text{A17})$$

which must hold together with and (A16). In this case, for $n > 4$ the only non-zero Weyl components are (using also (A10))

$$\Omega'_{\hat{i}\hat{j}} = \frac{1}{2} m_{\hat{i}}^k m_{\hat{j}}^l \left(c_{2||kl} - \frac{\Delta c_2}{n-2} h_{kl} \right) \quad (n > 4). \quad (\text{A18})$$

See [13] for $n = 4$.

A.3.3 Conditions for l to be a multiple WAND (\Rightarrow Weyl type D(bd))

It is clear from (A13) and (A14) that, in general, the vector field l of (A11) is not a WAND (not even a single one). The conditions for l being a multiple WAND read

$$2H = d_1(u, r) + \frac{2(\ln \sqrt{h})_{,u}}{n-2} r + c_2(u, x) r^2, \quad (\text{A19})$$

$$c_{2||ij} = \frac{\Delta c_2}{n-2} h_{ij}, \quad [(\ln \sqrt{h})_{,u}]_{||ij} = \frac{\Delta [(\ln \sqrt{h})_{,u}]}{n-2} h_{ij}. \quad (\text{A20})$$

For $n > 4$ the latter of these is identically satisfied thanks to (A10). When all these conditions are met the Weyl type becomes D(bd). This happens, for example, in the special case $c_{2,i} = 0$.

A.3.4 Conditions for the Weyl type D(bd) when l is *not* a multiple WAND

Even when the vector field l of (A11) is not a multiple WAND (i.e., (A19), (A20) are not simultaneously satisfied) the Weyl type can still be D(bd) provided a second multiple WAND (in addition to k , and different from l) exists. In order to find conditions for this to happen, it is necessary to perform a null rotation about k , i.e.,

$$k \mapsto k, \quad l \mapsto l + z_i m_i - \frac{1}{2} z_i z^i k, \quad m_i \mapsto m_i - z_i k, \quad (\text{A21})$$

such that, in the transformed frame, $\Psi'_i \mapsto 0$, $\tilde{\Psi}'_{\hat{i}\hat{j}\hat{k}} \mapsto 0$ and $\Omega'_{\hat{i}\hat{j}} \mapsto 0$. The transformation laws under (A21) for the negative b.w. Weyl components are given by eqs. (2.33)–(2.35) of [10] (while non-negative b.w. components are unchanged under (A21) since k is a multiple WAND). Using the fact that the Weyl type is II(bd), the condition $\Psi'_i \mapsto 0$ uniquely fixes the parameters z_i by

$$\Phi z_i = \frac{n-2}{n-1} \Psi'_i, \quad (\text{A22})$$

except in the case $\Phi = 0 = \Psi'_i$, for which the z_i remain arbitrary.

Next, requiring $\tilde{\Psi}'_{\hat{i}\hat{j}\hat{k}} \mapsto 0$ and using (A22) further gives (also in the case $\Phi = 0 = \Psi'_i$)

$$\tilde{\Psi}'_{\hat{i}\hat{j}\hat{k}} = -\tilde{\Phi}_{\hat{i}\hat{j}\hat{k}} z_i. \quad (\text{A23})$$

Finally, imposing $\Omega'_{\hat{i}\hat{j}} \mapsto 0$ gives

$$\Omega'_{\hat{i}\hat{j}} = \frac{2(n-2)}{n-3} \left(z_{(\hat{j}} \Psi'_{\hat{i})} - \frac{(z_{\hat{k}} \Psi'_{\hat{k}})}{n-2} \delta_{\hat{i}\hat{j}} \right) + 2z_{\hat{k}} \tilde{\Psi}'_{(\hat{i}\hat{j})\hat{k}} - \frac{n-1}{n-3} \Phi \left(z_i z_j - \frac{z^2}{n-2} \delta_{ij} \right) - z_{\hat{k}} z_{\hat{l}} \tilde{\Phi}_{\hat{k}\hat{l}\hat{i}\hat{j}}, \quad (\text{A24})$$

which after using (A22), (A23) (and multiplying by Φ) gives a constraint among the Weyl tensor components¹⁴

$$\Phi \Omega'_{\hat{i}\hat{j}} = \frac{n-2}{n-1} \left[\frac{n-2}{n-3} \left(\Psi'_{\hat{i}} \Psi'_{\hat{j}} - \frac{(\Psi'_{\hat{k}} \Psi'_{\hat{k}})}{n-2} \delta_{\hat{i}\hat{j}} \right) + \tilde{\Psi}'_{(\hat{i}\hat{j})\hat{k}} \Psi'_{\hat{k}} \right]. \quad (\text{A25})$$

Recalling that for the metric (A1) with $W^i = 0$ and (A9) one has $\tilde{\Psi}'_{\hat{i}\hat{j}\hat{k}} = 0$, and using (A12), eq. (A23) reduces to

$$\mathcal{C}_{\bar{l}\bar{j}\bar{k}} z_{\hat{l}} = 0. \quad (\text{A26})$$

This is a restriction on the Weyl tensor of h_{ij} that will not be true for a *generic* Einstein metric h_{ij} , therefore we can conclude that *generically the metric (A1) with $W^i = 0$ and (A9) is of genuine type II(bd)*. However, the “genericity” conditions may be violated if a special choice of H (e.g., in vacuum [25]) or of h_{ij} is made in (A1), in which case the Weyl types D(bd), D(bcd), and D(abd) are all possible (see, e.g., [25, 27, 30, 46]). The specific form of (A25) for spacetimes (A1) with $W^i = 0$ and (A9) can be obtained by substituting (A12)–(A14) and $\tilde{\Psi}'_{\hat{i}\hat{j}\hat{k}} = 0$ into (A25).

B Comments on the (Einstein) constraints (31) and (32) ($2 \leq p \leq n-2$)

The b.w. 0 components of \mathbf{F} can be divided into an electric and a magnetic part described, respectively, by $e_{i_1 \dots i_{p-2}}$ and $b_{i_1 \dots i_p}$ (cf. (14), (15)). The latter live in the transverse geometry of h_{ij} and must obey the constraints (31) and (32) (replaced by (33) if $2p = n$). In this appendix we discuss some general consequences of those constraints.

Recall that the form $e_{i_1 \dots i_{p-2}}$ is defined for $2 \leq p \leq n$ while $b_{i_1 \dots i_p}$ for $0 \leq p \leq n-2$. The cases $p = 0, n$ are trivial (footnote 2) while $p = 1, n-1$ require a special discussion (appendix C), so here we restrict for both $e_{i_1 \dots i_{p-2}}$ and $b_{i_1 \dots i_p}$ to the ranks $2 \leq p \leq n-2$. Let us also observe that the equations obeyed by $e_{i_1 \dots i_{p-2}}$ and $b_{i_1 \dots i_p}$ ((31) and (32)) are identical, so the algebraic constraints derived from them which apply to $b_{i_1 \dots i_p}$ for a certain p will also automatically apply to $e_{i_1 \dots i_{p'-2}}$ for $p' = p+2$ (and vice versa). Additionally, by duality constraints on $b_{i_1 \dots i_p}$ for a certain p will also apply on $e_{i_1 \dots i_{p'-2}}$ for $p' = n-p$ (and vice versa). In the magnetic and electric cases the following ranks p are worth mentioning.

B.1 Magnetic fields

- $p = 2$: this is the case studied in [27] and $b_{ij} = 0$ if n is odd, so that n must be even (thus, in particular, the $\ln r$ term in $2H$ (eq. (28)) vanishes for $n = 5$ [27]). If \mathbf{F} is assumed to be regular (and non-zero) on the transverse space, then Maxwell’s equations (19), (35) for b_{ij}

¹⁴In passing, let us observe that (A22), (A23) and (A25) give the necessary and sufficient conditions under which *any* Weyl tensor of type II(bd) admits a second multiple WAND (so being in fact of type D(bd)), since no special features of the Robinson-Trautman class have been used in their derivation. For the same reason, (A24) alone determines the conditions under which a Weyl tensor of type II(bd) admits a single WAND (in addition to a multiple one). This implies that a Weyl tensor of *proper* type D(bd) or D(bcd) admits *precisely two* WANDs (necessarily double) since (A24) has no non-zero solution if $\Omega'_{\hat{i}\hat{j}} = \Psi'_{\hat{i}\hat{j}\hat{k}} = \Psi'_i = 0$, $\Phi \neq 0$ (as easily seen after contraction with $z_{\hat{j}}$). However, Weyl tensors of type D(abd) may admit an infinity of multiple WANDs, and this occurs precisely when the equation $\tilde{\Phi}_{\hat{i}\hat{j}\hat{k}} z_{\hat{l}} = 0$ (cf. (A23)) admits a solution (in this case (A22) and (A25) are satisfied trivially) — see, e.g., [46, 50] for examples of such spacetimes. Similarly, an infinity of *single* WANDs exists for type D(abd) if the weaker condition $z_{\hat{k}} z_{\hat{l}} \tilde{\Phi}_{\hat{k}\hat{l}\hat{j}} = 0$ is satisfied. We observe that L. Wylleman has obtained more general results on the structure of multiple WANDs for any Weyl type D (private communication), some of which are mentioned in [40, 46].

further imply [27] that h_{ij} must be almost-Kähler (almost-Hermitian if $n = 6$). See eq. (66) of [27] for an example.

- $p = n - 4$: this is dual to the electric $p = 4$ case discussed in section B.2 so that again n must be even and h_{ij} almost-Kähler (almost-Hermitian if $n = 6$).
- $p = n - 3$: it is easy to see that in this case eq. (32) is impossible (unless $b_{i_1 \dots i_{n-3}} = 0$). Using also the previous observations one sees, e.g., that for $n = 5$ only $p = 3$ is permitted, for $n = 6$ only $p = 2, 4$, for $n = 7$ only $p = 5$, etc. This implies, in particular, that the $\mathcal{B}^2 r^{3-n} \ln r$ term in $2H$ (eq. (28)) can occur only for $p = (n + 1)/2 \geq 4$ with $n \geq 9$ (n odd).
- $p = n - 2$: $F_{i_1 \dots i_p}$ is proportional to the h -volume element. For example, this includes the magnetic counterpart of the electrovac Schwarzschild-Tangherlini black hole [4].
- $p = \frac{n}{2} - 1$ (n even): in this case if h_{ij} is a direct product of two $(\frac{n}{2} - 1)$ -dimensional spaces then a possible magnetic field can be written as a sum of their volume elements, see also [31]; for even p it is h -self-dual. This case is also special in that the magnetic field does not contribute to T_{ij} . In particular, this includes the case $n = 6$, $p = 2$ (again eq. (66) of [27] — with $D = 6$ — gives an example). See (62) for an example in 8D (after dualization).
- $p = md$, $n - 2 = Nd$: when p and $n - 2$ are both multiples of the same integer $d \geq 2$ (m and N are also positive integers, with $m \leq N$), eq. (3.18) of [31] gives a method of constructing $F_{i_1 \dots i_p}$ that works when h_{ij} is a direct product of $N = (n - 2)/d$ d -dimensional Einstein spaces (in particular they are all flat if $\mathcal{R} = 0$). (For $N = 2$, $m = 1$ this reduces to the case $p = \frac{n}{2} - 1$ discussed above.) For example: $n = 8$, $p = 3$, $d = 3$, $N = 2$, $m = 1$ (note that here $p = \frac{n}{2} - 1$); $n = 8$, $p = 4$, $d = 2$, $N = 3$, $m = 2$ (here $p = \frac{n}{2}$); $n = 10$, $p = 4$, $d = 2$, $N = 4$, $m = 2$ (here $p = \frac{n}{2} - 1$); $n = 11$, $p = 6$ [3], $d = 3$, $N = 3$, $m = 2$ [1] etc. Cf. (61) for an explicit example. In particular, this always works when $p = \frac{n}{2}$ with p even, i.e., when n is a multiple of 4 (in this case $d = 2$, $N = p - 1$), see example (84).
- $p = \frac{n}{2} - 2$ with p even: these solutions can be constructed as explained for the electric forms with $p = \frac{n}{2}$ in section B.2.

B.2 Electric fields

- $p = 2$: this is the standard case studied in [27] (eq. (31) becomes an identity) containing, e.g., Schwarzschild-Tangherlini black holes with electric charge [4].
- $p = 3$: here e_i is a 1-form and this case is obviously forbidden by (31) (indeed dual to a magnetic field with $p = n - 3$, also forbidden as already discussed), i.e., $e_i = 0$.
- $p = 4$: here e_{ij} is a 2-form and (as for a magnetic field with $p = 2$ [27] — cf. also section B.1) n must be even and h_{ij} almost-Kähler (almost-Hermitian if $n = 6$).
- $p = n - 2$: this is dual to a magnetic 2-form so again n must be even and h_{ij} almost-Kähler (almost-Hermitian if $n = 6$). Using also the previous observations one sees, e.g., that for $n = 5$ only $p = 2$ is permitted, for $n = 6$ only $p = 2, 4$, for $n = 7$ only $p = 2$, etc. This implies, in particular, that the $\mathcal{E}^2 r^{3-n} \ln r$ term in $2H$ (eq. (27)) can occur only for $p = (n + 1)/2 \geq 5$ with $n \geq 9$ (n odd).
- $p = \frac{n}{2} + 1$ (n even): in this case (dual to a magnetic field with $p = \frac{n}{2} - 1$) the electric field does not depend on r and if h_{ij} is a direct product of two $(\frac{n}{2} - 1)$ -dimensional spaces then a possible $e_{i_1 \dots i_{p-2}}$ can be written as $(du \wedge dr) \wedge$ (a sum of the volume elements of the two subspaces), cf. also [31]; for even p it is h -self-dual. The electric field does not contribute to T_{ij} . See example (62).

- $p - 2 = m'd$, $n - 2 = Nd$ ($d \geq 2$): this is dual to an already discussed magnetic field with $p = md$ (with $m = N - m'$), and one can similarly use eq. (3.25) of [31]. As in the magnetic case, this always works when $p = \frac{n}{2}$ with p even (with $d = 2$, $N = p - 1$), see example (84). (For $N = 2$, $m' = 1$ this reduces to the case $p = \frac{n}{2} + 1$ discussed above.) By combining the constructions for the electric and magnetic cases, for $d = 2$ ($m = m' + 1$) one can also construct dyonic fields [31].
- $p = \frac{n}{2} + 2$ with p even: these solutions can be constructed as explained for the magnetic forms with $p = \frac{n}{2}$ in section B.1.

B.3 Case $2p = n$ (n even)

It should be recalled that in the case $p = \frac{n}{2}$, if both an electric and a magnetic field are present then they generically obey the weaker constraint (33) (the corresponding ‘‘Maxwell’’ equations are also generically modified, as discussed in section 5.2). The construction with $p = \frac{n}{2}$ (with p even) mentioned above in sections B.1 and B.2 still provides examples in the special case when (31) and (32) are satisfied separately (and not just (33)), but more general solutions also exist, see for example (85). Note, however, that in the case $n = 6$, $p = 3$, not only (31) and (32) (as discussed in sections B.1 and B.2) but also the weaker constraint (33) can be satisfied only trivially by $\mathcal{E}^2 = 0 = \mathcal{B}^2$, as can be seen by directly substituting into (33) the most general possible form of a 1-form \mathbf{e} and a 3-form \mathbf{b} (examples are given by (100), (101)). Therefore, solutions $2p = n$ for which \mathbf{e} and \mathbf{b} are not both zero require $n \geq 8$ ($p \geq 4$) (see, e.g., (84), (85)).

C Cases $p = 1$ and $p = n - 1$

Here we analyze the special ranks $p = 1$ and $p = n - 1$. The results of section 3.1 apply also here and need not be repeated. Just note that there is no electric term $e_{i_1 \dots i_{p-2}}$ for $p = 1$ (while $\mathcal{B}_{ij}^2 = b_i b_j$), and no magnetic term $b_{i_1 \dots i_p}$ for $p = n - 1$.

As observed in section 3.2, the first difference appears in the Einstein equation for R_{ij} . Instead of (29) and (32) [(31)] we now have the trace-free equations

$$\mathcal{R}_{ij} - \frac{\mathcal{R}}{n-2} h_{ij} = \kappa_0 \left(b_i b_j - \frac{\mathcal{B}^2}{n-2} h_{ij} \right) \quad (p = 1), \quad (\text{C1})$$

$$\mathcal{R}_{ij} - \frac{\mathcal{R}}{n-2} h_{ij} = -(n-2)(n-3)\kappa_0 \left(\mathcal{E}_{ij}^2 - \frac{\mathcal{E}^2}{n-2} h_{ij} \right) \quad (p = n - 1), \quad (\text{C2})$$

while (30) remains true, so that, again, $h_{ij} = h^{1/(n-2)} \gamma_{ij}(x)$ (with $\det \gamma_{ij} = 1$).

As noticed, the results of section 3.3 are still valid also here. In particular, we have $F_u = 0$ for $p = 1$ and $F_{u_{i_1 \dots i_{n-2}}} = 0$ for $p = n - 1$, and (40) are still true.

As for section 3.4, we still obtain

$$\mu_{,i} = 0, \quad (\text{C3})$$

but due to coincidence of certain powers of r , eqs. (42), (44) and (45) are replaced by

$$(n-4) [\mathcal{R}_{,i} - \kappa_0 (\mathcal{B}^2)_{,i}] = 0 \quad (p = 1), \quad (\text{C4})$$

$$(n-4) [\mathcal{R}_{,i} - \kappa_0 (n-2) (\mathcal{E}^2)_{,i}] = 0 \quad (p = n - 1). \quad (\text{C5})$$

The above two equations in fact also follow from, respectively, (C1) and (C2) once the contracted Bianchi identity and the Maxwell equations in the geometry of h_{ij} are employed.

Finally, the equation discussed in section 3.5.1 contains now an additional r^{-2} term that, however, vanishes identically thanks to (C4), (C5) (except for $n = 4$, see section C.2 below). Hence one still obtains $(n-2)\mu_{,u} = -(n-1)\mu(\ln\sqrt{h})_{,u}$ (for $n \neq 4$). As in section 3.5.1, the latter equation together with (40) implies that we can choose coordinates such as

$$h_{ij} = h_{ij}(x), \quad \mu = \text{const}, \quad (\mathcal{B}^2)_{,u} = 0 \quad (p = 1), \quad (\mathcal{E}^2)_{,u} = 0 \quad (p = n - 1) \quad (n \neq 4), \quad (\text{C6})$$

which with (C4), (C5) for $n > 4$ give

$$\mathcal{R} = \kappa_0 \mathcal{B}^2 + \mathcal{R}_0 \quad (p = 1), \quad \mathcal{R} = \kappa_0(n-2)\mathcal{E}^2 + \mathcal{R}_0 \quad (p = n - 1), \quad (\text{C7})$$

where \mathcal{R}_0 is a constant (corresponding to the Ricci scalar of the transverse geometry in the vacuum limit).

C.1 Case $n > 4$

To summarize, for $n > 4$ the metric is given by (55) with

$$2H = \frac{\mathcal{R}_0}{(n-2)(n-3)} - \frac{2\Lambda}{(n-1)(n-2)} r^2 - \frac{\mu}{r^{n-3}}, \quad (\text{C8})$$

where \mathcal{R}_0 , Λ and μ are constants, and h_{ij} is a Riemannian metric satisfying (C1) or (C2) (and thus having a Ricci scalar given by (C7)). The Maxwell field is given by

$$\mathbf{F} = b_i(x) dx^i \quad (p = 1), \quad (\text{C9})$$

$$\mathbf{F} = \frac{1}{(n-3)!} r^{n-4} e_{i_1 \dots i_{n-3}}(x) du \wedge dr \wedge dx^{i_1} \wedge \dots \wedge dx^{i_{n-3}} \quad (p = n - 1), \quad (\text{C10})$$

where $b_i(x)$ and $e_{i_1 \dots i_{n-3}}(x)$ are harmonic forms in the geometry of h_{ij} . In contrast to the case $2 \leq p \leq n - 2$, here the Maxwell field does not enter the metric function H , but instead “back-reacts” on the transverse geometry h_{ij} , which thus cannot be Einstein. The Weyl type is D(bd) (since $H = H(r)$), the Maxwell type is D, and (59) are doubly aligned null directions for both the Weyl and Maxwell tensors. The metric is static (at least in regions where $H > 0$). We further observe that (C1) gives $(n-2)\mathcal{R}_{ij}b^i b^j = \mathcal{B}^2[\mathcal{R} + \kappa_0(n-3)\mathcal{B}^2]$, which is certainly non-negative if $\mathcal{R} \geq 0$. Since b_i is harmonic, this implies (see [42, 51]; also theorem 2.9 of [43]) that h_{ij} cannot describe a compact space having $\mathcal{R} \geq 0$ everywhere (unless, trivially, $\mathcal{R} = 0 = \mathcal{B}^2$).

Thanks to (C1) and (C2), it is easy to see that any $(n-2)$ -dimensional solution of the Euclidean Einstein–Maxwell theory (i.e., a metric h_{ij} coupled to a 1-form $b_i(x)$ or a $(n-3)$ -form $e_{i_1 \dots i_{n-3}}(x)$) can be used to generate an n -dimensional Robinson–Trautman spacetime coupled to a 1-form (or a $(n-1)$ -form). For example, by taking as a “seed” the Euclidean version of the 3D charged BTZ metric (with “ $J = 0$ ”) [52] we obtain the following 5D example.

Example ($n = 5$, $p = 1$)

$$h_{ij} dx^i dx^j = -(\lambda \rho^2 + m + \kappa_0 b_x^2 \ln \rho) d\tau^2 - \frac{d\rho^2}{\lambda \rho^2 + m + \kappa_0 b_x^2 \ln \rho} + \rho^2 dx^2, \\ 2H = \lambda - \frac{\Lambda}{6} r^2 - \frac{\mu}{r^2}, \quad \mathbf{F} = b_x dx, \quad (\text{C11})$$

where $\lambda = \frac{\mathcal{R}_0}{6}$, m and b_x are constants (respectively, the cosmological constant, mass parameter and field strength of the 3D BTZ solution), and $\mathcal{R} = 6\lambda + \kappa_0 b_x^2 \rho^{-2}$. Having a Lorentzian signature

requires (at least for a large ρ) that $\lambda < 0$. The dualization to $p = 4$ is obvious. We observe however that the $\rho = 0$ curvature singularity of the BTZ metric extends to the full 5D solution, interestingly also beyond the horizon(s). Euclideanization of this 5D solution can be used, in turn, to produce a 7D Robinson–Trautman solutions, and so on to higher odd dimensions.

C.2 Case $n = 4$

We observe that for $n = 4$ the l.h.s. of both (C1) ($p = 1$) and (C2) ($p = 3$) is identically zero (since h_{ij} is 2-dimensional), which implies, respectively, $b_i = 0$ and $e_i = 0$. Since we also have $F_u = 0$ for $p = 1$ and $F_{ui_1\dots i_{n-2}} = 0$ for $p = n - 1$, we conclude that \mathbf{F} vanishes identically and one is left only with vacuum Robinson–Trautman spacetimes, obeying the standard equation $-\Delta\mathcal{R} + 4\mu_{,u} + 6\mu(\ln\sqrt{h})_{,u} = 0$ (this contains, in particular, the Schwarzschild metric, which has $\mathbf{F} = 0$ and yet can describe a black hole with non-zero axionic charge in the presence of a non-zero Kalb–Ramond field — in fact it is the only static and asymptotically flat solution [53]). Therefore, in 4D the only electrovac spacetimes with $\mathbf{F} \neq 0$ are obtained for $p = 2$, which is the well-known Einstein–Maxwell case [12–14]. We observe that certain Robinson–Trautman 4D solutions with aligned $p = 3$ forms (axionic black holes) have been discussed in [31], but these involve multiple p -form fields and thus do not contradict our result.

References

- [1] R. Emparan and H. S. Reall, *Black holes in higher dimensions*, *Living Reviews in Relativity* **11** (2008), no. 6.
- [2] S. Hollands and A. Ishibashi, *Black hole uniqueness theorems in higher dimensional spacetimes*, *Class. Quantum Grav.* **29** (2012) 163001.
- [3] G. T. Horowitz, ed., *Black holes in higher dimensions*. Cambridge University Press, Cambridge, 2012.
- [4] F. R. Tangherlini, *Schwarzschild field in n dimensions and the dimensionality of space problem*, *Nuovo Cimento* **27** (1963) 636–651.
- [5] G. W. Gibbons and D. L. Wiltshire, *Space-time as a membrane in higher dimensions*, *Nucl. Phys. B* **287** (1987) 717–742.
- [6] R. C. Myers and M. J. Perry, *Black holes in higher dimensional space-times*, *Ann. Phys. (N. Y.)* **172** (1986) 304–347.
- [7] R. Emparan, S. Ohashi, and T. Shiromizu, *No-dipole-hair theorem for higher-dimensional static black holes*, *Phys. Rev. D* **82** (2010) 084032.
- [8] R. Güven, *Hertz potentials in higher dimensions*, *Class. Quantum Grav.* **6** (1989) 1961–1980.
- [9] T. Shiromizu, S. Ohashi, and K. Tanabe, *Perturbative no-hair property of form fields for higher dimensional static black holes*, *Phys. Rev. D* **83** (2011) 084016.
- [10] M. Durkee, V. Pravda, A. Pravdová, and H. S. Reall, *Generalization of the Geroch–Held–Penrose formalism to higher dimensions*, *Class. Quantum Grav.* **27** (2010) 215010.
- [11] M. Ortogio, *Asymptotic behaviour of Maxwell fields in higher dimensions*, *Phys. Rev. D* (2014), to appear [1406.3186].
- [12] I. Robinson and A. Trautman, *Some spherical gravitational waves in general relativity*, *Proc. R. Soc. A* **265** (1962) 463–473.

- [13] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers, and E. Herlt, *Exact Solutions of Einstein's Field Equations*. Cambridge University Press, Cambridge, second ed., 2003.
- [14] J. B. Griffiths and J. Podolský, *Exact Space-Times in Einstein's General Relativity*. Cambridge University Press, Cambridge, 2009.
- [15] R. Emparan, G. T. Horowitz, and R. C. Myers, *Exact description of black holes on branes*, *JHEP* **01** (2000) 007.
- [16] G. Bernardi de Freitas and H. S. Reall, *Algebraically special solutions in AdS/CFT*, *JHEP* **06** (2014) 148.
- [17] N. Van den Bergh, *Einstein–Maxwell null fields of Petrov type D*, *Class. Quantum Grav.* **6** (1989) 1871–1878.
- [18] M. Cahen and J. Sengier, *Espaces de classe D admettant un champ électromagnétique*, *Bull. Acad. Roy. Belg. Cl. Sci.* **53** (1967) 801–816.
- [19] J. Leroy, *Champs électromagnétiques à rayons intégrables, divergents et sans distorsion*, *Bull. Acad. Roy. Belg. Cl. Sci.* **62** (1976) 259–282.
- [20] R. Debever, N. Van den Bergh, and J. Leroy, *Diverging Einstein–Maxwell null fields of Petrov type D*, *Class. Quantum Grav.* **6** (1989) 1373–1378.
- [21] J. M. M. Senovilla, *Black hole formation by incoming electromagnetic radiation*, [gr-qc/1408.2778](#).
- [22] J. P. S. Lemos, *Gravitational collapse to toroidal, cylindrical, and planar black holes*, *Phys. Rev. D* **57** (1998) 4600–4605.
- [23] J. P. S. Lemos, *Collapsing shells of radiation in anti-de Sitter spacetimes and the hoop and cosmic censorship conjectures*, *Phys. Rev. D* **59** (1999) 044020.
- [24] P. Dadras, J. Firouzjaee, and R. Mansouri, *A concrete anti-de Sitter black hole with dynamical horizon having toroidal cross-sections and its characteristics*, *Europhys.Lett.* **100** (2012) 39001.
- [25] J. Podolský and M. Ortaggio, *Robinson–Trautman spacetimes in higher dimensions*, *Class. Quantum Grav.* **23** (2006) 5785–5797.
- [26] M. Ortaggio, *Higher dimensional spacetimes with a geodesic, shearfree, twistfree and expanding null congruence*, in *Proceedings of the XVII SIGRAV Conference (Torino, September 4–7, 2006)*, [gr-qc/0701036](#).
- [27] M. Ortaggio, J. Podolský, and M. Žofka, *Robinson–Trautman spacetimes with an electromagnetic field in higher dimensions*, *Class. Quantum Grav.* **25** (2008) 025006.
- [28] V. Pravda, A. Pravdová, and M. Ortaggio, *Type D Einstein spacetimes in higher dimensions*, *Class. Quantum Grav.* **24** (2007) 4407–4428.
- [29] R. Švarc and J. Podolský, *Absence of gyratons in the Robinson–Trautman class*, *Phys. Rev. D* **89** (2014) 124029.
- [30] J. Podolský and R. Švarc, *Algebraic structure of Robinson–Trautman and Kundt geometries in arbitrary dimension*, *Class. Quantum Grav.* (2014), to appear [[gr-qc/1406.3232](#)].
- [31] Y. Bardoux, M. M. Caldarelli, and C. Charmousis, *Shaping black holes with free fields*, *JHEP* **05** (2012) 054.
- [32] M. Henneaux and C. Teitelboim, *Dynamics of chiral (self-dual) p-forms*, *Phys. Lett. B* **206** (1988) 650–654.
- [33] M. Duff and P. van Nieuwenhuizen, *Quantum inequivalence of different field representations*, *Phys. Lett. B* **94** (1980) 179–182.

- [34] S. Deser, A. Gomberoff, M. Henneaux, and C. Teitelboim, *Duality, self-duality, sources and charge quantization in abelian N -form theories*, *Phys. Lett. B* **400** (1997) 80–86.
- [35] M. Bremer, H. Lü, C. N. Pope, and K. S. Stelle, *Dirac quantization conditions and Kaluza-Klein reduction*, *Nucl. Phys. B* **529** (1998) 259–294.
- [36] R. Milson, A. Coley, V. Pravda, and A. Pravdová, *Alignment and algebraically special tensors in Lorentzian geometry*, *Int. J. Geom. Meth. Mod. Phys.* **2** (2005) 41–61.
- [37] G. Bergqvist and J. Senovilla, *Null cone preserving maps, causal tensors and algebraic Rainich theory*, *Class. Quantum Grav.* **18** (2001) 5299–5326.
- [38] A. Coley, R. Milson, V. Pravda, and A. Pravdová, *Vanishing scalar invariant spacetimes in higher dimensions*, *Class. Quantum Grav.* **21** (2004) 5519–5542.
- [39] R. Milson, *Alignment and the classification of Lorentz-signature tensors*, [gr-qc/0411036](#).
- [40] S. Hervik, M. Ortaggio, and L. Wylleman, *Minimal tensors and purely electric or magnetic spacetimes of arbitrary dimension*, *Class. Quantum Grav.* **30** (2013) 165014.
- [41] A. Z. Petrov, *Einstein Spaces*. Pergamon Press, Oxford, translation of the 1961 Russian ed., 1969.
- [42] S. Bochner, *Curvature and Betti numbers*, *Ann. of Math.* **49** (1948) 379–390.
- [43] K. Yano and S. Bochner, *Curvature and Betti numbers*. No. 32 in Annals of Mathematics Studies. Princeton University Press, Princeton, 1953.
- [44] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry*, vol. 2. Interscience, New York, 1969.
- [45] H. W. Brinkmann, *Einstein spaces which are mapped conformally on each other*, *Math. Ann.* **94** (1925) 119–145.
- [46] M. Ortaggio, V. Pravda, and A. Pravdová, *Algebraic classification of higher dimensional spacetimes based on null alignment*, *Class. Quantum Grav.* **30** (2013) 013001.
- [47] M. Ortaggio, V. Pravda, and A. Pravdová, *Higher dimensional Kerr-Schild spacetimes*, *Class. Quantum Grav.* **26** (2009) 025008.
- [48] M. Henneaux and C. Teitelboim, *p -form electrodynamics*, *Found. Phys.* **16** (1986) 593–617.
- [49] M. Ortaggio, V. Pravda, and A. Pravdová, *Ricci identities in higher dimensions*, *Class. Quantum Grav.* **24** (2007) 1657–1664.
- [50] M. Ortaggio, V. Pravda, and A. Pravdová, *On the Goldberg-Sachs theorem in higher dimensions in the non-twisting case*, *Class. Quantum Grav.* **30** (2013) 075016.
- [51] S. Bochner, *Vector fields and Ricci curvature*, *Bull. Amer. Math. Soc.* **52** (1946) 776–797.
- [52] M. Bañados, C. Teitelboim, and J. Zanelli, *Black hole in three-dimensional spacetime*, *Phys. Rev. Lett.* **69** (1992) 1849–1851.
- [53] M. J. Bowick, S. B. Giddings, J. A. Harvey, G. T. Horowitz, and A. Strominger, *Axionic black holes and an Aharonov-Bohm effect for strings*, *Phys. Rev. Lett.* **61** (1988) 2823–2826.