

Seznam vzorců pro zápočtovou písemku.

$$i\hbar \frac{|\psi\rangle}{\partial t} = \hat{H}|\psi\rangle$$

Důležité komutátory

$$[\hat{x}_\alpha, \hat{p}_\beta] = i\hbar\delta_{\alpha,\beta} \quad [\hat{J}_\alpha, \hat{V}_\beta] = i\hbar\varepsilon_{\alpha,\beta,\gamma}\hat{V}_\gamma$$

Moment hybnosti a sférické harmoniky

$$\begin{aligned} \hat{J}^2|jm\rangle &= \hbar^2 j(j+1)|jm\rangle & \hat{J}_z|jm\rangle &= \hbar m|jm\rangle & \hat{J}_\pm &= \hat{J}_x \pm i\hat{J}_y \\ \hat{J}_\pm|jm\rangle &= \hbar\sqrt{(j \mp m)(j \pm m + 1)}|jm \pm 1\rangle \end{aligned}$$

Pauliho matice

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Několik prvních sférických harmonik:

$$\begin{array}{lll} Y_{00} = \frac{1}{\sqrt{4\pi}} & Y_{22} = \sqrt{\frac{15}{32\pi}} \frac{(x+iy)^2}{r^2} & Y_{33} = -\sqrt{\frac{35}{64\pi}} \frac{(x+iy)^3}{r^3} \\ = & Y_{21} = -\sqrt{\frac{15}{8\pi}} \frac{z(x+iy)}{r^2} & Y_{32} = \sqrt{\frac{105}{32\pi}} \frac{z(x+iy)^2}{r^3} \\ = & Y_{20} = \sqrt{\frac{5}{16\pi}} \frac{2z^2-x^2-y^2}{r^2} & Y_{31} = -\sqrt{\frac{21}{64\pi}} \frac{(4z^2-x^2-y^2)(x+iy)}{r^3} \\ Y_{11} = -\sqrt{\frac{3}{8\pi}} \frac{x+iy}{r} & Y_{2-1} = \sqrt{\frac{15}{8\pi}} \frac{z(x-iy)}{r^2} & Y_{30} = \sqrt{\frac{7}{16\pi}} \frac{z(2z^2-3x^2-3y^2)}{r^3} \\ Y_{10} = \sqrt{\frac{3}{4\pi}} \frac{z}{r} & Y_{2-2} = \sqrt{\frac{15}{32\pi}} \frac{(x-iy)^2}{r^2} & Y_{3-1} = \sqrt{\frac{21}{64\pi}} \frac{(4z^2-x^2-y^2)(x-iy)}{r^3} \\ Y_{1-1} = \sqrt{\frac{3}{8\pi}} \frac{x-iy}{r} & = & Y_{3-2} = \sqrt{\frac{105}{32\pi}} \frac{z(x-iy)^2}{r^3} \\ = & = & Y_{3-3} = \sqrt{\frac{35}{64\pi}} \frac{(x-iy)^3}{r^3} \end{array}$$

Harmonický oscilátor

$$\begin{aligned} x_0 &= \sqrt{\frac{\hbar}{m\omega}} & p_0 &= \frac{\hbar}{x_0} & \hat{a} &= \frac{\hat{x}/x_0 + i\hat{p}/p_0}{\sqrt{2}} \\ \hat{H} &= \frac{\hat{p}^2}{2m} + \frac{m\omega^2}{2}\hat{x}^2 = \hbar\omega(\hat{a}^\dagger\hat{a} + 1/2) & [\hat{a}, \hat{a}^\dagger] &= 1 \\ \hat{a}|n\rangle &= \sqrt{n}|n-1\rangle & \hat{a}^\dagger|n\rangle &= \sqrt{n+1}|n+1\rangle \\ \langle x|n\rangle &= \sqrt{\frac{1}{\sqrt{\pi}x_0 n! 2^n}} H_n(x/x_0) \exp(-(x/x_0)^2/2) \end{aligned}$$

Hermitovy polynomy

$$H_0(x) = 1, H_1(x) = 2x, H_2(x) = 4x^2 - 2, H_3(x) = 8x^3 - 12x.$$

Coulombický potenciál

$$V(x) = \gamma/r, \quad E_n = -\frac{m\gamma^2}{2\hbar^2 n^2}, \quad a = \frac{\hbar^2}{m|\gamma|}$$

Radiální vlnová $R_{nl}(r)$ funkce pro prvních pár stavů:

$$\begin{aligned} R_{10}(r) &= 2\sqrt{\frac{1}{a^3}} \exp(-r/a) \\ R_{20}(r) &= \sqrt{\frac{1}{2a^3}} (1 - r/2a) \exp(-r/2a) \\ R_{21}(r) &= \frac{1}{2} \sqrt{\frac{1}{6a^3}} r/a \exp(-r/2a) \end{aligned}$$