

Gyratons on Melvin spacetime

Hedvika Kadlecová* and Pavel Krtouš†

Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles University, V Holešovičkách 2, 180 00 Prague 8, Czech Republic

(Received 9 June 2010; published 25 August 2010)

We present and analyze new exact gyraton solutions of algebraic type II on a background which is a static, cylindrically symmetric Melvin universe of type D. For a vanishing electromagnetic field it reduces to previously studied gyratons on Minkowski background. We demonstrate that the solutions are a member of a more general family of the Kundt spacetimes. We show that the Einstein equations reduce to a set of mostly linear equations on a transverse 2-space and we discuss the properties of polynomial scalar curvature invariants which are generally nonconstant but unaffected by the presence of gyratons.

DOI: 10.1103/PhysRevD.82.044041

PACS numbers: 04.20.Jb, 04.30.-w, 04.40.Nr

I. INTRODUCTION

Gyraton solutions represent the gravitational field of a localized matter source with an intrinsic rotation which is moving at the speed of light. Such an idealized ultrarelativistic source can describe a pulse of a spinning radiation beam and it is accompanied by a sandwich or impulsive gravitational wave.

The gravitational fields generated by (nonrotating) light pulses and beams were already studied by Tolman [1] in 1934, who obtained the corresponding solution in the linear approximation of the Einstein theory. Exact solutions of the Einstein-Maxwell equations for such “pencils of light” were found and analyzed by Peres [2] and Bonnor [3–5]. These solutions belong to a general family of pp waves [6,7].

In the impulsive limit (i.e., for an infinitely thin beam, and for the delta-type distribution of the light pulse in time), the simplest of these solutions represents the well-known Aichelburg-Sexl metric [8] which describes the field of a pointlike null particle. Subsequently, more general impulsive waves were found [9–15] (for recent reviews see [16,17]).

The gyraton solutions are generalization of pp waves which belong to the Kundt class for which the source—the beam of radiation—carries not only energy, but also an additional angular momentum. Such spacetimes were first considered by Bonnor in [18], who studied the gravitational field created by a spinning null fluid. In some cases, this may be interpreted as a massless neutrino field [19].

Gyratons on Minkowski background are locally isometric to standard pp waves in the exterior vacuum region, outside the source. The interior region contains a nonexpanding null matter which possesses an intrinsic spin. In general, these solutions are obtained by keeping nondiag-

onal terms g_{ui} in the Brinkmann form [20] of the pp -wave solution, where u is the null coordinate and x^i are orthogonal spatial coordinates. The corresponding energy-momentum tensor thus also contains an extra nondiagonal term $T_{ui} = j_i$. In four dimensions, the terms g_{ui} can be set to zero *locally*, using a suitable gauge transformation. However, they cannot be *globally* removed because the gauge invariant contour integral $\oint g_{ui}(u, x^j) dx^i$ around the position of the gyraton is proportional to the nonzero angular momentum density j_i , which is nonvanishing.

These gyratons were investigated (in the linear approximation) in [21] in higher dimensional flat space; the exact gyraton solutions propagating in an asymptotically flat D -dimensional spacetime were further investigated in [22]. They proved that the Einstein’s equations for gyratons reduce to a set of linear equations in the Euclidean $(D - 2)$ -dimensional space and showed that the gyraton metrics belong to a class of so-called *VSI spacetimes* for which all polynomial scalar invariants, constructed from the curvature and its covariant derivatives, vanish identically [23]. (For the discussion of spacetimes with nonvanishing but nonpolynomial scalar invariants of curvature, see [24].) Subsequently, charged gyratons in Minkowski space in any dimension were presented in [25].

In [26], the exact gyraton solutions in the asymptotically anti-de Sitter spacetime were found. Namely, they obtained Siklos gyratons which generalize the Siklos family of nonexpanding waves [27] (investigated further in [28]) which belong to the class of spacetimes with constant scalar invariants (*CSI spacetimes*) [29–32].

Recently, the large class of gyratons on the direct-product spacetimes was found in [33], where we showed that this class of gyratons has similar properties as the previous gyratonic solutions: the Einstein’s equations reduce to a set of linear equations in transversal 2-space and these spacetimes belong to the CSI class of spacetimes.

Let us also mention that string gyratons in supergravity were recently found in [34]. Supersymmetric gyraton so-

*hedvika.kadlecova@centrum.cz

†Pavel.Krtous@mff.cuni.cz

lutions were also obtained in minimal gauged theory in five dimensions in [35], where the configuration represents a generalization of the Siklos waves with a nonzero angular momentum in anti-de Sitter space.

The gyratons are important in studies of production of mini black holes or in cosmic ray experiments. The theory of high energy particle collisions was developed in [36,37] and was applied to gyraton models in [38].

The main purpose of this paper is to further extend the family of gyratonic solutions. In particular, we present new gyraton solutions of algebraic type II, propagating in the Melvin universe.

The Melvin universe [39,40] is a nonsingular electrovacuum solution with physical properties which are interesting both from a classical and a quantum point of view. The spacetime represents a parallel bundle of magnetic (or electric) flux held together by its own gravitational attraction. The transverse space orthogonal to the direction of the flux has a nontrivial spatial geometry. It was represented in [41] by a suitable embedding diagram which resembles a tall narrow-necked vase. Also in [41,42] it was shown that no motion can get too far from the axis of symmetry. This aspect is analogous to the attractive effect of a negative cosmological constant in the anti-de Sitter universe.

The Melvin universe was considered as an important model in astrophysical processes related to gravitational collapse because of its stability. It was shown in [40,41] that the spacetime is surprisingly stable against small radial perturbations and also against large perturbations which are concentrated in a finite region about the axis of symmetry. The asymmetries are radiated away in gravitational and electromagnetic waves [43,44].

The Melvin universe also appears as a limit in more complicated solutions; in [45] it is obtained as a specific limit of a charged C-metric. The Melvin universe has been generalized to Kaluza-Klein and dilaton theories [46], to nonlinear electrodynamics [47], and has important applications in the study of quantum black hole pair creation in a background electromagnetic field [48–54].

The Melvin universe recently attracted a new interest because it is possible to find gravitational waves in the Melvin universe [43] by an ultrarelativistic boost of the Schwarzschild-Melvin black hole metric [44]. It was shown that these wave solutions are straightforward impulsive limits of a more general class of Kundt spacetimes of type II with an arbitrary profile function, which can be interpreted as gravitational waves propagating on the Melvin spacetime. The gyraton spacetimes investigated in this paper are generalizations of such Kundt waves when their ultrarelativistic source is made of a “spinning matter.”

The paper is organized as follows. In Sec. II we review basic information about the Melvin universe which will be useful in the paper. We derive the ansatz for the gyraton metric by a direct transformation from the Kundt form of

the metric to Melvin’s coordinates. We also review the transverse space geometry of the wave front.

In Sec. III, we derive the field equations and we simplify them introducing potentials. We discuss the structure of the equations and the gauge freedom of the solutions.

Next, in Sec. IV, we solve the Einstein-Maxwell equations in the special case of the ϕ -independent spacetimes, especially with a thin matter source localized on the axis of symmetry.

In Sec. V we concentrate on the interpretation of the gyraton solutions. We discuss the properties of the scalar polynomial invariants and the geometric properties of the principal null congruence. We evaluate the curvature tensor in an appropriate tetrad, discuss the Petrov type, the matter content of the spacetime, and properties of the electromagnetic field.

The main results of the paper are summarized in concluding Sec. VI. Some technical results needed to derive the field equations are left to the Appendix.

II. THE GYRATONS ON THE MELVIN SPACETIME

A. The Melvin universe

In this section we briefly review basic properties of the Melvin spacetime [39,40,44] which will be useful throughout the paper. The Melvin universe describes an axial electromagnetic field concentrating under influence of its self-gravity. The strength of the electromagnetic field is determined by the parameters E and B . In cylindrical coordinates (t, z, ρ, ϕ) , the metric and the Maxwell tensor read

$$ds^2 = \Sigma^2(-dt^2 + dz^2 + d\rho^2) + \Sigma^{-2}\rho^2 d\phi^2, \quad (2.1)$$

$$F = E dz \wedge dt + B \Sigma^{-2} \rho d\rho \wedge d\phi, \quad (2.2)$$

where

$$\Sigma = 1 + \frac{1}{4} \varrho_{\text{EM}} \rho^2. \quad (2.3)$$

The constant ϱ_{EM} is given by the parameters E, B as¹

$$\varrho_{\text{EM}} = \frac{\kappa \varepsilon_o}{2} (E^2 + B^2). \quad (2.4)$$

Introducing double null coordinates,

$$u = \frac{1}{\sqrt{2}}(t - z), \quad v = \frac{1}{\sqrt{2}}(t + z), \quad (2.5)$$

we obtain an alternative expression for metric (2.1) and the Maxwell tensor (2.2):

$$ds^2 = \Sigma^2(-2dudv + d\rho^2) + \Sigma^{-2}\rho^2 d\phi^2, \quad (2.6)$$

¹ $\kappa = 8\pi G$ and ε_o are gravitational and electromagnetic constants. There are two standard choices of geometrical units: the Gaussian with $\kappa = 8\pi$ and $\varepsilon_o = 1/4\pi$, and SI like with $\kappa = \varepsilon_o = 1$.

$$F = E dv \wedge du + B \Sigma^{-2} \rho d\rho \wedge d\phi. \quad (2.7)$$

The electromagnetic field can be rewritten also in the complex self-dual form,²

$$\mathcal{F} = \mathcal{B}(dv \wedge du - i \Sigma^{-2} \rho d\rho \wedge d\phi) \quad (2.8)$$

with the complex constant \mathcal{B} defined as

$$\mathcal{B} = E + iB. \quad (2.9)$$

The metric (2.1) resembles a vacuum solution of the Levi-Civita family [6] for a large value of ρ . For $\mathcal{Q}_{\text{EM}} = 0$ ($E, B = 0$) spacetime reduces to the Minkowski spacetime in cylindrical coordinates. For $E \neq 0, B = 0$ the Maxwell tensor describes an electric field pointing along the z direction, whereas for $E = 0, B \neq 0$ we get a purely magnetic field oriented along the z direction.

The metric admits the four Killing vectors

$$\partial_t, \quad \partial_z, \quad \partial_\phi, \quad z\partial_t + t\partial_z = v\partial_v - u\partial_u, \quad (2.10)$$

which correspond to staticity, cylindrical symmetry, and invariance under a boost transformation. Using the adapted null tetrad $k = \partial_v$, $l = \Sigma^{-2} \partial_u$, and $m = \frac{1}{\sqrt{2}}(\Sigma^{-1} \partial_\rho - i \Sigma \rho^{-1} \partial_\phi)$, the only nonvanishing components of Weyl and Ricci tensors are

$$\Psi_2 = \frac{1}{2} \mathcal{Q}_{\text{EM}}(-1 + \frac{1}{4} \mathcal{Q}_{\text{EM}} \rho^2) \Sigma^{-4}, \quad \Phi_{11} = \frac{1}{2} \mathcal{Q}_{\text{EM}} \Sigma^{-4}. \quad (2.11)$$

This demonstrates that the Melvin universe is a nonvacuum solution of the Petrov type D, except at points satisfying $\rho = 2/\sqrt{\mathcal{Q}_{\text{EM}}}$, where the Weyl tensor vanishes. It is interesting to note that the scalar curvature vanishes, $R = 0$.

To conclude, the Melvin spacetime belongs to the family of nonexpanding, nontwisting type-D electrovacuum solutions investigated by Plebański [55]. As a consequence, it also belongs to the general Kundt class [6,7] which will be important in the following text.

B. The ansatz for the gyratons on Melvin universe

Gyratons are generalized gravitational waves corresponding to null sources with intrinsic rotation. In general, the gyron solutions are obtained by adding nondiagonal terms to the metric of the standard gravitational wave solutions, or in other words, by keeping the nondiagonal terms g_{ui} in the standard Kundt metric [6]. Therefore, we derive the ansatz for the gyron on Melvin spacetime by adding such new terms to the Kundt form of the Melvin metric. It can be explicitly obtained from (2.6) by trans-

formation

$$v = \Sigma^{-2} r, \quad (2.12)$$

which leads to

$$ds^2 = -2dudr + ds_\perp^2 + 2rW_i dudx^i. \quad (2.13)$$

Here we introduced a 2-dimensional metric

$$ds_\perp^2 = \Sigma^2 d\rho^2 + \Sigma^{-2} \rho^2 d\phi^2 \quad (2.14)$$

and r -independent 1-form $W = W_i dx^i$,

$$W = 2 \frac{\Sigma, \rho}{\Sigma} d\rho. \quad (2.15)$$

These tensors can be understood as tensors on space spanned by two coordinates ρ, ϕ . This space can be covered by other suitable spatial coordinates x^i , and we will use the Latin indices i, j, \dots to label the corresponding tensor components.

By an appropriate transformation of coordinates [7,44], the 2-dimensional metric ds_\perp^2 can be transformed into a conformally flat form, which in the standard complex null coordinates $\zeta, \bar{\zeta}$ reads

$$ds_\perp^2 = \frac{2}{\rho^2} d\zeta d\bar{\zeta}. \quad (2.16)$$

Such a transformation brings the metric (2.13) into the Kundt form.

The gyron generalization of (2.13) then reads

$$ds^2 = -2\Sigma^2 H du^2 - 2dudr + ds_\perp^2 + 2(rW_i - \Sigma^2 a_i) dudx^i. \quad (2.17)$$

We have added the term $-2\Sigma^2 H du^2$ which represents a gravitational wave on the Melvin universe [43,44] with an arbitrary profile function H , and the nondiagonal terms $-2\Sigma^2 a_i dudx^i$ characteristic for gyratons. It will be shown in the following that these terms can be generated by specific gyron matter.

Transforming back to the Melvin coordinate v and cylindrical coordinates ρ, ϕ , we obtain the ansatz for the metric describing the gyron on Melvin spacetime,³

$$ds^2 = -2\Sigma^2 H du^2 - 2\Sigma^2 dudv + (\Sigma^2 d\rho^2 + \Sigma^{-2} \rho^2 d\phi^2) + 2\Sigma^2 (a_\rho dud\rho + a_\phi dud\phi). \quad (2.18)$$

The function $H(u, v, \rho, \phi)$ can depend on all coordinates, but we assume that the functions $a_i(u, \rho, \phi)$ are v independent (it actually follows from the Maxwell equations as will be shown below). Let us note that the previously cited works assumed also the function H v independent.

³Here we use notation different from [43,44]; we use $-2H du^2$ instead of $-H du^2$ to match our notation in [33].

²We follow the notation of [6], namely, $\mathcal{F} \equiv F + i \star F$ is a complex self-dual Maxwell tensor, where the 4-dimensional Hodge dual is $\star F_{\mu\nu} = \frac{1}{2} \varepsilon_{\mu\nu\rho\sigma} F^{\rho\sigma}$. The self-dual condition reads $\star \mathcal{F} = -i \mathcal{F}$. The orientation of the 4-dimensional Levi-Civita tensor is fixed by the sign of the component $\varepsilon_{vur\phi} = \rho \Sigma^2$. The energy-momentum tensor of the electromagnetic field is given by $T_{\mu\nu} = \frac{\varepsilon_v}{2} \mathcal{F}_\mu{}^\rho \bar{\mathcal{F}}_{\nu\rho}$.

C. The ansatz for the matter

The metric should satisfy the Einstein equations with a stress-energy tensor generated by the electromagnetic field and the gyratonic source,

$$G_{\mu\nu} = \kappa(T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{gyr}}). \quad (2.19)$$

We assume that the electromagnetic field (2.7) modified by the gyration is given by

$$F = E dv \wedge du + B \Sigma^{-2} \rho d\rho \wedge d\phi + \sigma_j du \wedge dx^j. \quad (2.20)$$

Similar to [33], we have added the term $\sigma_j du \wedge dx^j$.

To evaluate Maxwell equations, it is useful to write down the self-dual form of the Maxwell tensor \mathcal{F} . The Hodge dual of (2.20) reads

$$\begin{aligned} \star F &= B dv \wedge du - E \Sigma^{-2} \rho d\rho \wedge d\phi \\ &+ (*\sigma + Ba - E*a)_j du \wedge dx^j, \end{aligned} \quad (2.21)$$

where the asterisk $*$ means the 2-dimensional Hodge dual defined on the transversal space; see Sec. II D. For the self-dual Maxwell tensor \mathcal{F} we thus obtain

$$\begin{aligned} \mathcal{F} &= \mathcal{B}(dv \wedge du - i \Sigma^{-2} \rho d\rho \wedge d\phi \\ &+ (\mathcal{S} - i*a)_j du \wedge dx^j), \end{aligned} \quad (2.22)$$

where we introduced a complex transverse 1-form \mathcal{S}_j ,

$$\mathcal{B} \mathcal{S}_j = (\sigma + i*\sigma)_j + iB(a + i*a)_j. \quad (2.23)$$

This form is self-dual with respect to the Hodge duality $*$ on the transversal space,

$$*\mathcal{S}_j = -i\mathcal{S}_j, \quad (2.24)$$

and therefore it can be written using a real 1-form s_j :

$$\mathcal{S}_j = (s + i*s)_j. \quad (2.25)$$

The original 1-form σ_i can be expressed in terms of s_j as

$$\sigma_j = E s_j - B*(s - a)_j. \quad (2.26)$$

In the following we use $s_j(v, u, \rho, \phi)$ as a basic variable for the electromagnetic field.

The stress-energy tensor $T_{\mu\nu}^{\text{EM}}$ corresponding to the field (2.22) is given in (A3).

Finally, we must define the gyratonic matter by specifying the structure of its stress-energy tensor. It is obtained from the standard stress-energy tensor of a null fluid by adding terms corresponding to ‘‘internal spatial rotation’’ of the fluid:

$$\kappa T^{\text{gyr}} = j_u du^2 + 2j_\rho du d\rho + 2j_\phi du d\phi. \quad (2.27)$$

We admit a general coordinate dependence of the source functions $j_u(v, u, \rho, \phi)$ and $j_j(v, u, \rho, \phi)$. However, it will be shown below that the field equations enforce a trivial v dependence.

The gyration source is described only on a phenomenological level, by its stress-energy tensor (2.27), which is assumed to be given, and our aim is to determine its influence on the metric and the electromagnetic field. However, we have to consider that the gyration stress-energy tensor is locally conserved. It means that the functions j_u and j_j must satisfy the constraint given by

$$T^{\text{gyr}}{}_{\mu\nu}{}^{;\nu} = 0. \quad (2.28)$$

Of course, if we had considered a specific internal structure of the gyratonic matter, the local energy-momentum conservation would have been a consequence of field equations for the gyration. Without that, we have to require (2.28) explicitly.

To conclude, the fields are characterized by functions Σ , H , a_j , and s_j , which must be determined by the field equations provided the gyration sources j_u and j_j and the constants E and B are given.

D. The geometry of the transverse space

The geometry (2.18) identifies the null geodesic congruence generated by ∂_v , which is parametrized by an affine parameter v , the family of null hypersurfaces $u = \text{const}$, and 2-dimensional *transverse spaces* $u, v = \text{const}$.

The gravitational wave moves along the null direction ∂_v , i.e., it propagates with the speed of light along the z direction, which is the direction of the electromagnetic field. The hypersurface $u = \text{const}$ corresponds to the surface of the constant ‘‘phase,’’ and the transverse spaces $u, v = \text{const}$ are spatial wave fronts of the wave.

Physical quantities do not depend on the affine parameter v , or this dependence is trivial and it will be explicitly found. Specifically, the geometry of the transverse space is v independent.

It turns out to be convenient to restrict various quantities to the transverse space. For example, we can interpret a_i and s_i as components of u -dependent 1-forms on the transverse space. Our goal is to formulate all equations for physical quantities on the transverse spaces. For that we need to review some properties of the transverse geometry. It was studied in detail in [44], and on a general level in [33]; nevertheless it will be useful to mention some of the properties explicitly.

The transverse metric is obtained by restriction of the full metric (2.18) to the transverse space and it is given by the expression (2.14),

$$ds_\top^2 = g_{\top ij} dx^i dx^j = \Sigma^2 d\rho^2 + \Sigma^{-2} \rho^2 d\phi^2. \quad (2.29)$$

The associated Gauss curvature is given by the scalar curvature R_\top ,

$$K = \frac{1}{2} R_\top = \frac{\rho_{\text{EM}}}{\Sigma^4} \left(2 - \frac{1}{4} \rho_{\text{EM}}^2 \right). \quad (2.30)$$

It is obvious that only the electromagnetic field is responsible for the nonflatness of the transversal space—it is

insensitive to the presence of the gyron. For $\varrho_{\text{EM}} = 0$ the curvature vanishes and we get the flat plane.

In general, the curvature is not constant and it is finite everywhere. The Gauss curvature K has maximum on the axis $\rho = 0$ where it is equal to $2\varrho_{\text{EM}}$; it is positive for $0 \leq \rho < 2\sqrt{2}/\sqrt{\varrho_{\text{EM}}}$, and vanishes on the circle at $\rho = 2\sqrt{2}/\sqrt{\varrho_{\text{EM}}}$. For $\rho > 2\sqrt{2}/\sqrt{\varrho_{\text{EM}}}$, it goes to negative values and it has its minimum $K = -\frac{\varrho_{\text{EM}}}{256}$ at $\rho = 2\sqrt{3}/\sqrt{\varrho_{\text{EM}}}$. Then it grows again, and as $\rho \rightarrow +\infty$ the curvature vanishes, $K \rightarrow 0^-$.

The circumference of a circle of constant radius ρ is vanishing when $\rho \rightarrow +\infty$. Therefore, we measure a much shorter circumference for larger ρ —as if we would move “along the stem of the wine glass toward the narrowing end,” [40,41,44].

The 2-dimensional Levi-Civita tensor ϵ_{ij} associated with the metric (2.29) is $\epsilon = \rho d\rho \wedge d\phi$. The covariant derivative will be denoted by a colon, e.g., $a_{i;j}$. We raise and lower the Latin indices using $g_{\top ij}$, which differs from lowering indices using $g_{\alpha\beta}$ thanks to nonvanishing terms g_{ui} . We use a shorthand for a transverse square of the norm of a 1-form a_i as

$$a^2 \equiv a^i a_i = \Sigma^{-2} a_\rho^2 + \rho^{-2} \Sigma^2 a_\phi^2. \quad (2.31)$$

In two dimensions, the Hodge duals of 0-, 1-, and 2-forms φ , a_i , and f_{ij} , respectively, read

$$\begin{aligned} (*\varphi)_{ij} &= \varphi \epsilon_{ij}, & (*a)_i &= a_j \epsilon^j_i, \\ *f &= \frac{1}{2} f_{ij} \epsilon^{ij} = \frac{1}{\rho} f_\rho \phi. \end{aligned} \quad (2.32)$$

For convenience, we also introduce an explicit notation for 2-dimensional divergence and rotation of a transverse 1-form a_i ,

$$\begin{aligned} \text{div} a &\equiv a_i{}^{;i} = \frac{1}{\Sigma^2} a_{\rho,\rho} + \frac{\Sigma^2}{\rho^2} a_{\phi,\phi} + \frac{1}{\rho \Sigma^2} a_\rho - \frac{\Sigma_\rho^2}{\Sigma^4} a_\rho, \\ \text{rota} &\equiv \epsilon^{ij} a_{j,i} = \frac{1}{\rho} (a_{\phi,\rho} - a_{\rho,\phi}). \end{aligned} \quad (2.33)$$

For 2-form f_{ij} we get

$$\text{div} f \equiv f_{ij}{}^{;j} = \frac{1}{\Sigma^2} \left(f_{\phi\rho,\rho} - \frac{1}{\rho} f_{\phi\rho} \right) d\phi + \frac{\Sigma^2}{\rho^2} f_{\rho\phi,\phi} d\rho, \quad (2.34)$$

and $\text{rot} f = 0$. We can generalize the action of divergence and rotation also on a scalar function f as $\text{div} f = 0$ and $\text{rot} f = -*df$. Note that the divergence and rotation are related as $\text{div} a = \text{rot} *a$, and the relation to the transverse exterior derivative is $da = *\text{rot} a$. Clearly, $\text{div} \text{div} a = 0$, $\text{div} \text{rot} a = 0$, and $\text{rot} da = 0$.

The Laplace operator of a function ψ reads

$$\begin{aligned} \Delta \psi &= \psi_{;i}{}^{;i} \\ &= \frac{1}{\Sigma^2} \psi_{,\rho\rho} + \frac{\Sigma^2}{\rho^2} \psi_{,\phi\phi} + \frac{1}{\rho \Sigma^2} \psi_{,\rho} - \frac{\Sigma_\rho^2}{\Sigma^4} \psi_{,\rho}, \end{aligned} \quad (2.35)$$

and for a transverse 1-form η it is defined as $\Delta \eta \equiv d \text{div} \eta - \text{rot} \text{rot} \eta$.

Finally, the transverse space is topologically trivial since it has topology of a plane. We can thus assume that the Poincare lemma ($d\omega = 0 \Rightarrow \omega = d\sigma$) holds, which in terms of rotation and divergence means that $\text{div} \omega = 0$ implies $\omega = \text{rot} \sigma$. However, since the transverse space is noncompact and we do not know *a priori* boundary conditions for various quantities at infinity $\rho \rightarrow \infty$, we have to admit nontrivial harmonics. Therefore, we cannot assume a uniqueness of the Hodge decomposition. Moreover, in some cases it can be physically relevant to consider also topologically nontrivial harmonics which are singular, e.g., at the origin $\rho = 0$. Such solutions would correspond to fields around singular sources localized on the axis. However, we will ignore these cases in a general discussion.

III. THE FIELD EQUATIONS

A. The field equations for matter

Now, we will investigate the equations for matter, i.e., the Maxwell equations for electromagnetic field and the condition (2.28) for the gyron source.

Both Maxwell equations for real Maxwell tensor are equivalent to the cyclic Maxwell equation for the self-dual Maxwell tensor (2.22),

$$\begin{aligned} d\mathcal{F} &= \mathcal{B} [\partial_\nu (s + i*(s-a))_j dv \wedge du \wedge dx^j \\ &\quad - (\text{rots} + i \text{div}(s-a)) du \wedge \epsilon] = 0. \end{aligned} \quad (3.1)$$

From the real part we immediately get that the 1-form s_i is ν independent, $\partial_\nu s_i = 0$, and rotation free,

$$\text{rot} s = 0. \quad (3.2)$$

From the imaginary part it follows that the 1-form a_i is also ν independent (as we have already mentioned above) and it satisfies

$$\text{div}(s-a) = 0. \quad (3.3)$$

Equations (3.2) and (3.3) guarantee the existence and determine the structure of potentials which will be discussed in detail in Sec. III C.

Next, we analyze the condition (2.28) for the gyron source. When translated to the transverse space, it gives

$$-\partial_\nu j_i dx^i + (-\partial_\nu j_u + \text{div}(\Sigma^2 j) + \Sigma^2 a^i \partial_\nu j_i) du = 0. \quad (3.4)$$

The source functions j_i must be thus v independent and j_u has to have the structure

$$j_u = v \operatorname{div}(\Sigma^2 j) + \iota, \quad (3.5)$$

where $\iota(u, x^i)$ is a v -independent function. The gyraton source (2.27) is therefore fully determined by three v -independent functions $\iota(u, x^j)$ and $j_i(u, x^j)$.

Equation (3.5) gives us also an insight into interpretation of the gyratonic terms j_i . They are composed from two contributions: one is related to a kind of ‘‘heat flow’’ which changes energy j_u of the fluid, and the other which is related to intrinsic rotation of the fluid. The source representing heat flow has thus nonvanishing divergence $\operatorname{div}(\Sigma^2 j)$ and we require a vanishing rotational part $\operatorname{rot}(\Sigma^2 j)$. In opposite, the source representing intrinsic rotation has vanishing heat flow, i.e., it satisfies

$$\operatorname{div}(\Sigma^2 j) = 0. \quad (3.6)$$

Such a source can be written in terms of a rotational potential ν as $j = -\Sigma^{-2} \operatorname{rot} \nu$. In components it means

$$j_\rho = -\frac{1}{\rho} \nu_{,\phi}, \quad j_\phi = \frac{\rho}{\Sigma^4} \nu_{,\rho}. \quad (3.7)$$

Physically more relevant is the rotational part of the source, since it can describe the spin of the null fluid, or, in a specific limit, of the polarized beam of light. Terms related to heating flow have bad causal behavior and therefore typically do not satisfy various energy conditions and they are thus rather unphysical. Interpretation of the gyraton source was discussed previously also in [21,22,26,33].

B. The Einstein equations

The Einstein gravitational law (2.19) needs the Einstein tensor and the electromagnetic and gyraton stress-energy tensors. These quantities can be found in the Appendix. We can combine them and inspect various components of the Einstein equations.

The vu component determines the function Σ , namely, it gives the condition

$$-\rho(\Sigma_{,\rho})^2 + 2\Sigma\Sigma_{,\rho} = \varrho_{\text{EM}}\rho. \quad (3.8)$$

It is straightforward to check that it is satisfied again by Σ in the Melvin form (2.3).

The transverse diagonal components $\rho\rho$ and $\phi\phi$ require

$$\partial_v^2 H = 0, \quad (3.9)$$

thus we obtain the explicit v dependence of the metric function H as

$$H = gv + h, \quad (3.10)$$

where we have introduced v -independent functions $g(u, x^j)$ and $h(u, x^j)$.

The remaining nontrivial components of the Einstein equations are those involving the gyraton source (2.27).

The ui components give the equation related to j_i ,

$$j_i = \frac{1}{2} \Sigma^2 f_{ij}{}^{;j} - (\Sigma^2)_{,k} g^{kj} f_{ji} + g_{,i} + \frac{2\varrho_{\text{EM}}}{\Sigma^2} (s_i - a_i), \quad (3.11)$$

where we introduce the exterior derivative f_{ij} of the transverse 1-form a_i and its Hodge dual $b(u, x^j)$,

$$f_{ij} \equiv a_{j,i} - a_{i,j} = (*b)_{ij}, \quad (3.12)$$

$$b \equiv *f = \operatorname{rot} a. \quad (3.13)$$

In terms of b , Eq. (3.11) can be rewritten as

$$\Sigma^2 j = \frac{1}{2} \operatorname{rot}(\Sigma^4 b) + \Sigma^2 dg + 2\varrho_{\text{EM}}(s - a). \quad (3.14)$$

Here and in the following dg represents just the transverse gradient $dg = g_{,i} dx^i$, and for simplicity we skipped the transverse indices.

It is useful to split the equation into divergence and rotation parts by applying div and rot :

$$\operatorname{div}(\Sigma^2 j) = \operatorname{div}(\Sigma^2 dg), \quad (3.15)$$

$$\operatorname{rot}(\Sigma^2 j) = -\frac{1}{2} \Delta(\Sigma^4 b) + \operatorname{rot}(\Sigma^2 dg) - 2\varrho_{\text{EM}} b. \quad (3.16)$$

Here we have used the relations (3.3) and (3.13). The formula (3.15) is the equation for g , (3.16) is the equation for b , and together with (3.13) it determines a_i . In the next section we will return to these equations introducing suitable potentials which allow us to escape the necessity of taking an additional derivative of (3.14) when deriving the equation for a_i .

Finally, from the uu component of the Einstein equation we obtain

$$\begin{aligned} j_u = & \operatorname{div}(\Sigma^2 dg)v + \Sigma^2(\Delta h - (\Sigma^{-2})_{,\rho} h_{,\rho}) + \frac{1}{2} \Sigma^4 b^2 \\ & + 2\Sigma^2 a^i g_{,i} + \partial_u \operatorname{div}(\Sigma^2 a) + g \operatorname{div}(\Sigma^2 a) \\ & - 2\varrho_{\text{EM}}(s - a)^2. \end{aligned} \quad (3.17)$$

When we compare the coefficient in front of v with (3.15) we find that it has structure consistent with (3.5). The nontrivial v -independent part of (3.17) gives the equation for the metric function h ,

$$\begin{aligned} & \Sigma^2(\Delta h - (\Sigma^{-2})_{,\rho} h_{,\rho}) \\ = & \iota - \frac{1}{2} \Sigma^4 b^2 - 2\Sigma^2 a^i g_{,i} - \partial_u \operatorname{div}(\Sigma^2 a) \\ & - g \operatorname{div}(\Sigma^2 a) + 2\varrho_{\text{EM}}(s - a)^2. \end{aligned} \quad (3.18)$$

C. Introducing potentials

In the previous section we have found that the Maxwell and Einstein equations reduce to two potential equations (3.2) and (3.3), and two source equations (3.11) and (3.17).

According to the two-dimensional Hodge decomposition we can express the 1-form a_i using two scalar poten-

tials $\kappa(u, x^j)$ and $\lambda(u, x^j)$,

$$a = d\kappa + \text{rot}\lambda. \quad (3.19)$$

These potentials control the divergence and the rotation of a_i as

$$\text{div } a = \Delta\kappa, \quad \text{rot } a = -\Delta\lambda. \quad (3.20)$$

Comparing with (3.13) we thus obtain the equation for λ in terms of b ,

$$\Delta\lambda = -b. \quad (3.21)$$

The first potential equation (3.2) gives immediately that s_i has a potential $\varphi(u, x^j)$,

$$s = d\varphi. \quad (3.22)$$

Equation (3.3) implies that there exists a potential $\psi(u, x^j)$ satisfying

$$s - a = -\text{rot}\psi. \quad (3.23)$$

In terms of these potentials the 1-form (2.26) from the real Maxwell tensor (2.7) reads

$$\sigma = Ed\varphi + Bd\psi. \quad (3.24)$$

The potential φ and ψ are not, however, independent. Substituting (3.19) and (3.22) into (3.23) we obtain the key relation among the potentials:

$$d(\varphi - \kappa) + \text{rot}(\psi - \lambda) = 0. \quad (3.25)$$

If the Hodge decomposition was unique, the gradient and rotational parts would be vanishing separately, i.e., we would get $\varphi = \kappa$ and $\psi = \lambda$ (up to unimportant constants). The nonuniqueness of the Hodge decomposition is linked to the possible existence of a nontrivial harmonic 1-form $\eta_i(u, x^j)$,

$$\Delta\eta = 0, \quad (3.26)$$

in terms of which the gradient and rotation parts of (3.25) can be expressed as

$$\varphi - \kappa = \text{div}\eta, \quad (3.27)$$

$$\psi - \lambda = -\text{rot}\eta. \quad (3.28)$$

These are equations for electromagnetic potentials φ (or ψ , respectively) in terms of the metric potentials κ and λ . The 1-form η encodes an extra freedom, which allows a nontrivial electromagnetic field not uniquely determined by the metric. (Such contributions would allow one to take into account, for example, an additional electromagnetic charge localized at the origin of the transverse space; cf. the discussion of particular cases in Sec. IV.) However, sufficiently restrictive conditions at the infinity and the smoothness on the whole transverse space for the potentials would eliminate this freedom, so the case $\eta = 0$ is a rather representative choice.

After eliminating the electromagnetic potentials, we need to formulate the equations for κ and λ . We start with the divergence part of (3.14) which can be rewritten using the modified Laplace operator acting on the metric function g :

$$\Sigma^2(\Delta g - (\Sigma^{-2})_{,\rho}g_{,\rho}) = \text{div}(\Sigma^2 j). \quad (3.29)$$

However, for g solving this equation, (3.15) is also the integrability condition for the quantity $\Sigma^2(dg - j)$. It can thus be written in terms of a potential ω ,

$$\Sigma^2(dg - j) = \text{rot } \omega, \quad (3.30)$$

or, more explicitly,

$$d\omega = -\Sigma^2(\text{rot } g + *j). \quad (3.31)$$

For the source (3.7) without intrinsic ‘‘heating,’’ the right-hand side of (3.29) is zero and the function ω has an additive contribution from the rotational potential ν , namely, it has to satisfy

$$d(\omega - \nu) = -\Sigma^2 \text{rot } g. \quad (3.32)$$

The function ω contains information from the source j and from the metric function g relevant for the rotational part of Eq. (3.14). Indeed, substituting the potentials and $\text{rot } \omega$ into (3.14) we obtain

$$\text{rot}(\frac{1}{2}\Sigma^4 b - 2\varrho_{\text{EM}}\psi + \omega) = 0. \quad (3.33)$$

Substituting (3.21) and (3.28), and integrating (absorbing an integration constant to ω), we derive the equation for the potential λ

$$\frac{1}{2}\Sigma^4 \Delta\lambda + 2\varrho_{\text{EM}}\lambda = \omega + 2\varrho_{\text{EM}} \text{rot}\eta. \quad (3.34)$$

Taking into account relations (3.26) and (3.28), we obtain the alternative equation for ψ

$$\frac{1}{2}\Sigma^4 \Delta\psi + 2\varrho_{\text{EM}}\psi = \omega. \quad (3.35)$$

The other metric potential κ remains unrestricted. This nonuniqueness is related to the gauge freedom discussed in detail in Sec. III E. This coordinate freedom allows us to set the potential κ to an arbitrary convenient form, e.g., to eliminate it completely.

D. Discussion of the field equations

We have thus formulated all field equations as equations on the transverse space. They are written in a separated form, i.e., they can be solved one after the other: First, one has to find harmonic 1-form η satisfying Eq. (3.26) and metric function g satisfying (3.15). It allows one to integrate the function ω which together with $\text{rot}\eta$ appears as a source in Eq. (3.34) for the potential λ . Using the gauge freedom one can choose the other potential κ . Equations (3.27) and (3.28) determine electromagnetic potentials and through Eq. (2.26) the electromagnetic field. Finally, the remaining metric function h is determined by

Eq. (3.18), in which the previously computed quantities contribute to the source on the right-hand side.

The most complicated field equations—(3.18), (3.29), and (3.34)—are partial differential equations on the 2-dimensional space, which are solvable, at least in principle. They all contain a modified Laplace operator, Eqs. (3.18) and (3.29) for h and g , the same one, namely,

$$\Sigma^2(\Delta f - (\Sigma^{-2})_{,\rho} f_{,\rho}) \equiv \text{div}(\Sigma^2 \text{d}f). \quad (3.36)$$

Solutions for particular cases (assuming rotational symmetry) will be discussed in Sec. IV.

It is important to observe that, except Eq. (3.18) for h , the field equations are linear. We can thus superpose two solutions simply by adding the fields together. Only in the last step, when computing the source for Eq. (3.18), one has to include total superposition of the fields g , a_i , and σ_i since the expression for the source is nonlinear.

Finally, we have not paid much attention to the u dependence of the studied quantities. All metric functions, matter fields, and sources can depend on the coordinate u and this dependence does not enter the field equations except in one term on the right-hand side of Eq. (3.18). The profile of the gyraton in the u direction can thus be specified arbitrarily. It corresponds to the fact that both matter and gravitational field move with the speed of light and information on different hypersurfaces $u = \text{const}$ evolves rather independently.

Also the dependence of the metric and fields on the coordinate v is very simple and it was found for all quantities explicitly.

E. The gauge transformation

The coordinate transformation $\tilde{v} \rightarrow v = \tilde{v} - \chi(u, x^j)$ accompanied by the following redefinition of the metric functions and matter fields:

$$\begin{aligned} v &= \tilde{v} - \chi, & g &= \tilde{g}, & h &= \tilde{h} + \tilde{g}\chi + \partial_u \chi, \\ a_i &= \tilde{a}_i - \chi_{,i}, & s_i &= \tilde{s}_i - \chi_{,i}, & \sigma_i &= \tilde{\sigma}_i - E\chi_{,i}, \\ \kappa &= \tilde{\kappa} - \chi, & \lambda &= \tilde{\lambda}, & \varphi &= \tilde{\varphi} - \chi, & \psi &= \tilde{\psi}, \\ \omega &= \tilde{\omega}, & \eta &= \tilde{\eta}, & j_i &= \tilde{j}_i, & \iota &= \tilde{\iota} + \chi \text{div} j, \end{aligned} \quad (3.37)$$

leaves the metric, the Maxwell tensor, and the gyraton stress-energy tensor in the same form. Therefore, all the field equations remain the same. This transformation is thus a pure gauge transformation and we can use it to simplify the solution of the equations.

There are two natural choices of gauge: we can eliminate either the metric potential κ or the electromagnetic potential φ . In the first case $a = \text{rot} \lambda$ and $\varphi = \text{div} \eta$. In the latter case $s = 0$, $\sigma = B \text{d}\psi$, $\kappa = -\text{div} \eta$, and $a = \text{rot} \psi$.

Let us mention that in [33] an analogous gauge transformation allowed us to choose also the metric function g .

For the gyraton on the Melvin universe the metric function g decouples from κ and it is gauge independent.

The discussed gauge transformation has a clear geometrical meaning: it corresponds to a shift of the origin of the affine parameter v of the null congruence ∂_v . Note that such a change redefines transverse spaces.

IV. SPECIAL CASES

In this section we will study the special solutions of the field equations. Namely, we restrict to the axially symmetric situation, i.e., to the case when the geometry and the fields are invariant under action generated by the rotational vector ∂_ϕ . Further, we concentrate on the gyraton generated by a thin beam of matter concentrated at the origin of the transverse space which means on the axis of symmetry.

Thank to linearity mentioned at the end of Sec. III D, we can discuss various special cases separately. However, the geometry of the spacetime with gyraton is not merely a superposition of the individual contribution since the nonlinear coupling in the metric function h .

We do not discuss in detail the u dependence of the fields. It does not enter the field equations, except in the source term of Eq. (3.18) for h . The u dependence of the gyraton sources and corresponding dependence of other fields can thus be chosen arbitrarily.

The symmetry assumption enforces that quantities a_i , g , h , σ_i , j_i , and ι are ϕ independent. It can induce a slightly weaker condition on the potentials: typically, they have a linear dependence on ϕ .

The thin beam approximation requires that gyratonic matter is concentrated at the origin o of the transverse space given by $\rho = 0$, i.e., j_i and ι should be distributions with the support at the origin. However, we relax this condition slightly in the case of gyratonic heat flow discussed in Sec. IV B.

In all discussed cases we use a natural gauge

$$\kappa = 0, \quad \text{i.e.,} \quad a = \text{rot} \lambda. \quad (4.1)$$

Together with the symmetry assumptions it implies $\lambda_{,\phi} = \text{const}$.

A. Pure gravitational gyraton

We start with the simplest vacuum case: we set $j_i = 0$ and $\iota = 0$ and we assume no pure electromagnetic contribution, i.e., $\eta = 0$. The equation for g has only a trivial solution $g = g_o = \text{const}$, the function ω is also a trivial constant and we obtain Eq. (3.34) with vanishing right-hand side. Taking into account that $\lambda_{,\phi} = \text{const}$ we obtain that λ must be ϕ independent (a ϕ -linear term in λ would require an analogous term in the source) and we obtain an ordinary differential equation

$$\frac{1}{\rho} \left(\frac{\rho}{\Sigma^2} \lambda_{,\rho} \right)_{,\rho} + \frac{4q_{\text{EM}}}{\Sigma^4} \lambda = 0. \quad (4.2)$$

Substituting (2.3) for Σ , it is possible to obtain two independent solutions, one regular at the origin,

$$\lambda = -\gamma \Sigma^{-1} \left(1 - \frac{3}{4} \varrho_{\text{EM}} \rho^2\right), \quad (4.3)$$

and the other behaving as $\log \rho$ near the origin, which corresponds to a delta source at the origin and it will be discussed in Sec. IV D.

The metric 1-form $a = \text{rot } \lambda$ has thus components

$$a_\rho = 0, \quad a_\phi = -\frac{\rho}{\Sigma^2} \lambda_{,\rho} = -\gamma \frac{2\varrho_{\text{EM}}}{\Sigma^4} \rho^2, \quad (4.4)$$

and its ‘‘strength’’ $b = \text{rot } a$ is then

$$b = -\gamma \frac{4\varrho_{\text{EM}}}{\Sigma^5} \left(1 - \frac{3}{4} \varrho_{\text{EM}} \rho^2\right). \quad (4.5)$$

These can be plugged into (3.18) which turns out to be

$$\frac{1}{\rho} (\rho h_{,\rho})_{,\rho} = -\gamma^2 \frac{8\varrho_{\text{EM}}^2}{\Sigma^6} \left(1 - \frac{1}{4} \varrho_{\text{EM}} \rho^2\right) \left(1 - \frac{9}{4} \varrho_{\text{EM}} \rho^2\right). \quad (4.6)$$

It can be integrated explicitly; however, the result is rather long, so we skip it.

B. Nonspinning beam with heat flow

As we discussed in Sec. III A, the gyraton source can have two contributions: one, which changes gyratonic energy density j_u and the other corresponding to intrinsic rotation. Let us investigate the case when the gyratonic energy is concentrated at the origin (a thin beam) but there is axially symmetric energy flow in the ∂_ρ direction which accumulates energy at the beam (a heating process). Namely, we assume that $\partial_\nu j_u = \text{div}(\Sigma^2 j)$ is nonzero only at the origin, so elsewhere the transverse flow j_i must satisfy Eq. (3.6). Such j_i has the form

$$j_\rho = \frac{\alpha}{2\pi} \frac{1}{\rho}, \quad j_\phi = 0, \quad (4.7)$$

and the increasing energy of the gyraton is given by

$$j_u = \alpha \nu \delta_o. \quad (4.8)$$

Here, δ_o stands for the transverse delta function localized at the origin o , normalized to the standard metric volume element on the transverse space.

Since the heating is localized only at the origin, the gyraton source (4.7) can be locally written using the source potential ν , cf. (3.7), as

$$\nu = -\frac{\alpha}{2\pi} \phi. \quad (4.9)$$

Note however, that the potential cannot be defined globally and it is not well behaved at the origin.

We again assume no pure electromagnetic contribution, i.e., $\eta = 0$. The requirement of the axial symmetry enforces that the difference $\Sigma^{-2} \text{rot } \omega$ between $\text{d}g$ and j , cf.

(3.30), must be zero, i.e., $\omega = 0$. The ϕ -independent metric function g must thus satisfy $g_{,\rho} = j_\rho$ which gives

$$g = \frac{\alpha}{2\pi} \log \rho + g_o. \quad (4.10)$$

The equation for λ has again the form (4.2) and in this case we choose the trivial solution $\lambda = 0$. With the gauge $\kappa = 0$ we thus obtain the only nontrivial field to be the metric function g , otherwise $a_i = 0$ and $\sigma_i = 0$. The source for Eq. (3.18) is also trivial, given only by ι , and we will study it in the next case.

C. Nonspinning light beam

A particular example of the gyraton source is standard null fluid. The thin nonspinning beam localized at the origin is described by the source

$$\iota = \varepsilon \delta_o, \quad j_i = 0. \quad (4.11)$$

We can set all the fields except h to be zero: $g = 0$, $a_i = 0$, and $\sigma_i = 0$. The equation for h outside the origin is $\rho^{-1} (\rho h_{,\rho})_{,\rho} = 0$ which (with proper fixing of the source constant) gives

$$h = \frac{\varepsilon}{2\pi} \log \rho. \quad (4.12)$$

D. Thin gyraton—spinning light beam

Finally we proceed with the most characteristic representant of the gyratonic matter. It is a simple null beam of energy localized at the origin with no heating, which, however, contains an intrinsic energy rotation. Since we have a pointlike source at the transverse space, we can speak about inner spin instead of a global rotational energy flow.

The gyraton source has a form⁴

$$\iota = \varepsilon \delta_o, \quad (4.13)$$

$$j = -\Sigma^{-2} \text{rot } \nu, \quad \text{with } \nu = \beta \delta_o. \quad (4.14)$$

The symmetry assumptions together with the no-heating requirement implies $g = \text{const}$, which we choose to be zero in this case. Equation (3.30) also implies that $\omega = \nu$. Ignoring again the electromagnetic contribution, $\eta = 0$, we obtain the equation for λ

$$\frac{1}{\rho} \left(\frac{\rho}{\Sigma^2} \lambda_{,\rho} \right)_{,\rho} + \frac{4\varrho_{\text{EM}}}{\Sigma^4} \lambda = \Sigma^{-4} \nu. \quad (4.15)$$

The solution of the homogeneous equation with a singular behavior $\sim \log \rho$ corresponding to the delta function (4.14) at the origin reads

⁴The exact structure of the singular source j_i at the origin can be read out from the singular solution (4.16) of Eq. (3.34) for λ below.

$$\lambda = -\frac{\beta}{2\pi}\Sigma^{-1}\left(1 + \frac{1}{2}\varrho_{\text{EM}}\rho^2 - \frac{3}{64}\varrho_{\text{EM}}^2\rho^4 - \frac{1}{768}\varrho_{\text{EM}}^3\rho^4 + \frac{1}{2}\left(1 - \frac{3}{4}\varrho_{\text{EM}}\rho^2\right)\log\left(\frac{1}{4}\varrho_{\text{EM}}\rho^2\right)\right). \quad (4.16)$$

The multiplicative constant in the argument of the logarithm can be chosen arbitrary since it generates only an additional homogeneous contribution of the form (4.3).

The metric 1-form a_i and its rotation b are

$$a_\rho = 0, \\ a_\phi = -\frac{\beta}{2\pi}\Sigma^{-4}\left(1 - \frac{3}{8}\varrho_{\text{EM}}^2\rho^4 - \frac{1}{32}\varrho_{\text{EM}}^3\rho^6 - \frac{1}{768}\varrho_{\text{EM}}^4\rho^8 - \varrho_{\text{EM}}\rho^2\log\left(\frac{1}{4}\varrho_{\text{EM}}\rho^2\right)\right), \quad (4.17)$$

and

$$b = 2\frac{\beta}{2\pi}\varrho_{\text{EM}}\Sigma^{-5}\left(\left(1 - \frac{3}{4}\varrho_{\text{EM}}\rho^2\right)\log\left(\frac{1}{4}\varrho_{\text{EM}}\rho^2\right) + 2 + \varrho_{\text{EM}}\rho^2 - \frac{3}{32}\varrho_{\text{EM}}^2\rho^4 - \frac{1}{384}\varrho_{\text{EM}}^3\rho^6\right). \quad (4.18)$$

The source for Eq. (3.18) becomes cumbersome and lengthy, but treatable, in principle.

E. Electromagnetic wave

In the previous examples we have ignored the possibility of a nontrivial electromagnetic field. Namely, we have assumed that the electromagnetic field is given by the metric potentials via relations $\varphi = \kappa$ and $\psi = \lambda$. However, we have already observed that Eq. (3.25) admits also other solutions which we have parametrized using a harmonic 1-form η . To include such solutions we should classify all 1-form harmonics on the transverse space. But if we restrict to the axially symmetric fields we can solve the potential equation (3.25) directly, without referring to η explicitly.

Let us study a pure electromagnetic contribution to the matter, i.e., we assume $\iota = 0$, $j_i = 0$ here. We can thus take a trivial vanishing solution for g which implies $\omega = 0$.

The symmetry assumptions tell us that 1-forms s , σ , and a are ϕ independent, which implies that all potentials including $\varphi - \kappa$ and $\psi - \lambda$ can be at most linear in ϕ :

$$\varphi - \kappa = \hat{\varphi}(\rho) + \varphi_\phi\phi, \quad \psi - \lambda = \hat{\psi}(\rho) + \psi_\phi\phi, \quad (4.19)$$

φ_ϕ , ψ_ϕ being constants and $\hat{\varphi}$ and $\hat{\psi}$ functions of ρ only. Substituting into (3.25) we get

$$\left(\hat{\varphi}_{,\rho} - \frac{\Sigma^2}{\rho}\psi_\phi\right)d\rho + \left(\varphi_\phi + \frac{\rho}{\Sigma^2}\hat{\psi}_{,\rho}\right)d\phi = 0, \quad (4.20)$$

which implies

$$\hat{\varphi}_{,\rho} = \frac{\Sigma^2}{\rho}\psi_\phi, \quad \hat{\psi}_{,\rho} = -\frac{\Sigma^2}{\rho}\varphi_\phi. \quad (4.21)$$

Taking into account (2.3), it can be easily integrated and substituting back to (4.19) we obtain

$$\varphi = \kappa + \psi_\phi(\log\rho + \frac{1}{8}\varrho_{\text{EM}}\rho^2) + \varphi_\phi\phi, \\ \psi = \lambda - \varphi_\phi(\log\rho + \frac{1}{8}\varrho_{\text{EM}}\rho^2) + \psi_\phi\phi. \quad (4.22)$$

At this moment it is easier to solve Eq. (3.35) for ψ with vanishing right-hand side instead of Eq. (3.34) for λ . It has solutions in the form (4.3) and (4.16). The metric potential λ is then given by the second of the equations (4.19). The metric potential κ is vanishing thanks to our gauge.

After choosing the solution for ψ , we can thus compute all quantities a , b , s , and σ and substitute them to Eq. (3.18) for h .

Let us mention that solution (4.21) is singular at the origin. A careful distributional calculation would show that equations for the potentials (3.22) and (3.23) may not be satisfied at the origin. Tracing this singular term back to Maxwell equations, it could lead to nonvanishing electric charges localized at the origin. However, since we have not written down the Maxwell equations with sources, we do not discuss these terms in more detail.

V. PROPERTIES OF THE GYRATON SPACETIMES

A. Gravitational field

In this section we discuss some of the geometrical properties of the gyratonic solutions.

One of the important characteristics of spacetimes are the scalar polynomial invariants which are constructed only from the curvature and its covariant derivatives. It was shown that gyratons in the Minkowski spacetime [22] have all the scalar polynomial invariants vanishing (VSI spacetimes) [23], the gyratons in the anti-de Sitter [26] and direct-product spacetimes [33] have all invariants constant (CSI spacetimes) [29].

In these cases, the invariants are independent of all metric functions which characterize the gyraton, and have the same values as the corresponding invariants of the background spacetime. We observe that similar property is valid also for the gyraton on Melvin spacetime, however, in this case the invariants are generally *nonconstant*, namely, they depend on the coordinate ρ . This property is a consequence of the general theorem holding for the relevant subclass of the Kundt solution; see Theorem II.7 in [56].

The metric (2.18) admits the null vector $k = \partial_\nu$. It is a Killing vector for $g = 0$ and $\text{div}(\Sigma^2 j) = 0$, i.e., for the heating part in the gyratonic source. The covariant derivative of k is given by

$$k_{\alpha;\beta} = -\Sigma^{-2}(\partial_\nu H)k_\alpha k_\beta + \Sigma^{-2}k_{[\alpha}\nabla_{\beta]}\Sigma^2. \quad (5.1)$$

We observe that the congruence is not even recurrent [6] as

in the case of a gyration on direct-product spacetimes [33]. For $\partial_v H = 0$ we recover the formula in Garfinkle and Melvin [43]. In general, the nonrecurrency of the congruence is related to the nonvanishing spin coefficient $\tau_{\text{NP}} = -\frac{1}{\sqrt{2}}\frac{1}{\Sigma^2}\Sigma_{,\rho}$; cf. [23].

The null character of k and the condition (5.1) imply that the null congruence with tangent vector k is geodesic, expansion free, shear free, and twist free, and the spacetime thus belongs to the Kundt class.

Next we calculate components of the curvature tensors with respect to the following adapted null tetrad $\{k, l, m, \bar{m}\}$ [6]

$$\begin{aligned} k &= \partial_v, & l &= \frac{1}{\Sigma^2}(\partial_u - H\partial_v), \\ m &= \frac{1}{\sqrt{2}\Sigma}\left(a_m\partial_v + \partial_\rho - i\frac{\Sigma^2}{\rho}\partial_\phi\right). \end{aligned} \quad (5.2)$$

Here, we have introduced the projection of a transverse 1-form a on the vector m

$$a_m = m^i a_i = a_\rho - i\frac{\Sigma^2}{\rho}a_\phi = (a + i * a)_\rho, \quad (5.3)$$

and we will use an analogous notation also for components of the transverse gradient of a real function f

$$f_{,m} = m^i f_{,i} = f_{,\rho} - i\frac{\Sigma^2}{\rho}f_{,\phi}, \quad f_{,\bar{m}} = \overline{f_{,m}}. \quad (5.4)$$

The dual tetrad of 1-forms $\{\Theta^{(k)}, \Theta^{(l)}, \Theta^{(m)}, \Theta^{(\bar{m})}\}$ has a simple form

$$\begin{aligned} \Theta^{(k)} &= dv + Hdu - a, & \Theta^{(l)} &= \Sigma^2 du, \\ \Theta^{(m)} &= \frac{\Sigma}{\sqrt{2}}(d\rho - i * d\rho), & \Theta^{(\bar{m})} &= \frac{\Sigma}{\sqrt{2}}(d\rho + i * d\rho). \end{aligned} \quad (5.5)$$

With respect to this tetrad, we have found that the nonvanishing curvature components are given by four new components and by those which are the same for the Melvin universe (2.6). The nonvanishing Ricci scalars are

$$\begin{aligned} \Phi_{12} &= \frac{1}{4\sqrt{2}}\frac{1}{\Sigma^3}(\Sigma^2 ib_{,m} + 2ib(\Sigma^2)_{,\rho} + 2g_{,m}), \\ \Phi_{22} &= \frac{1}{2}\frac{1}{\Sigma^2}\left(\Delta H - (\Sigma^{-2})_{,\rho}H_{,\rho} + \frac{1}{2}\Sigma^2 b^2 + 2a^i g_{,i} \right. \\ &\quad \left. + \Sigma^{-2}(g + \partial_u)\text{div}(\Sigma^2 a)\right), \end{aligned} \quad (5.6)$$

and the nonvanishing Weyl scalars read

$$\begin{aligned} \Psi_3 &= \frac{1}{4\sqrt{2}}\frac{1}{\Sigma^3}(\Sigma^2 ib_{,\bar{m}} + ib(\Sigma^2)_{,\rho} + 2g_{,\bar{m}}), \\ \Psi_4 &= \frac{1}{2}\frac{1}{\Sigma^5}\left[2\Sigma a_{\bar{m}}g_{,\bar{m}} + \Sigma\left(H_{,\rho\rho} - \frac{\Sigma^4}{\rho^2}H_{,\phi\phi} + 2i\frac{\Sigma^2}{\rho}H_{,\phi\rho}\right) \right. \\ &\quad \left. + \Sigma(g + \partial_u)\left(a_{\rho,\rho} - \frac{\Sigma^4}{\rho^2}a_{\phi,\phi} + i\frac{\Sigma^2}{\rho}(a_{\rho,\phi} + a_{\phi,\rho})\right) \right. \\ &\quad \left. + \left(2\Sigma_{,\rho} - \frac{\Sigma}{\rho}\right)\left(H_{,\rho} + \partial_u a_\rho + g a_\rho \right. \right. \\ &\quad \left. \left. + 2i\frac{\Sigma^2}{\rho}(H_{,\phi} + \partial_u a_\phi + g a_\phi)\right)\right]. \end{aligned} \quad (5.7)$$

In particular, there exists a relation between Ψ_3 and Φ_{12} :

$$\bar{\Phi}_{12} + \Psi_3 = \frac{1}{4\sqrt{2}}\frac{1}{\Sigma^3}[-i(\Sigma^2)_{,\rho}b + 4g_{,\bar{m}}]. \quad (5.8)$$

Therefore, the metric (2.18) describes the transversal gravitational wave (Ψ_4 term) in the k direction with a longitudinal wave component (Ψ_3 term). The gravitational wave is accompanied by an aligned pure radiation field (Φ_{22} term) with non-null component (Φ_{12} term) propagating in the Melvin universe. In fact, the scalars Ψ_3 and Φ_{12} are generated by the gyration functions a_i and the function g . In general the spacetime (2.18) is of Petrov type II.

Let us now investigate the subcases of our solutions. When we set the gyration functions $a_i = 0$, the Ricci scalars become

$$\begin{aligned} \Phi_{12} &= \frac{1}{2\sqrt{2}}\frac{1}{\Sigma^3}g_{,m}, \\ \Phi_{22} &= \frac{1}{2}\frac{1}{\Sigma^4\rho^2}[\rho(\rho H_{,\rho})_{,\rho} + \Sigma^4 H_{,\phi\phi}] \end{aligned} \quad (5.9)$$

and the Weyl scalars then read

$$\begin{aligned} \Psi_3 &= \frac{1}{2\sqrt{2}}\frac{1}{\Sigma^3}g_{,\bar{m}}, \\ \Psi_4 &= \frac{1}{2}\frac{1}{\Sigma^5\rho}[\Sigma\rho H_{,\rho\rho} + (2\rho\Sigma_{,\rho} - \Sigma)H_{,\rho}] \\ &\quad - \frac{1}{2}\frac{1}{\rho^2}H_{,\phi\phi} + i\frac{1}{2}\frac{1}{\Sigma^3\rho^2}[2\partial_\rho(\Sigma g H) - 3\Sigma H]_{,\phi}. \end{aligned} \quad (5.10)$$

We again obtain the spacetime with similar characteristics as for the full gyration metric (2.18). However, the scalars Ψ_3 and Φ_{12} , which now depend only on function g , are now related by an even simpler relation:

$$\bar{\Phi}_{12} = \Psi_3. \quad (5.11)$$

If we assume additionally H to be v independent (i.e., $g = 0$) we obtain the only nonvanishing scalars Φ_{22} and Ψ_4 in the same form as in (5.9) and (5.10). This case and its subcases were thoroughly discussed in [44].⁵

⁵The terms Φ_{22} and Ψ_4 have a little different form which is caused by the slightly different choice of null tetrad in our paper.

Finally, let us mention that for $\varrho_{\text{EM}} = 0$ the background reduces to Minkowski spacetime and we thus recover the gyraton moving on the Minkowski background as an important subcase of our solutions.

B. Electromagnetic field

The gyraton propagates in a non-null electromagnetic field (2.22), the influence of which on the geometry is characterized by its density ϱ_{EM} (2.4). The electromagnetic field is modified by the gyraton through the $\sigma_i du \wedge dx^i$ terms. The electromagnetic field can be rewritten in terms of potentials using (3.24) as

$$F = E (dv \wedge du + du \wedge d\varphi) + B (\Sigma^{-2} \rho d\rho \wedge d\phi + du \wedge d\psi). \quad (5.12)$$

It describes a superposition of electric and magnetic fields, both pointing along the z direction, which are modified by the gravitation field of the gyraton. The additional term does not have a simple structure of electric or magnetic field, however both are of the form $du \wedge df$ with f being the proper potential.

The electromagnetic field projected on the null tetrad (5.2) is characterized by three scalars Φ_i ,

$$\Phi_0 = 0, \quad \Phi_1 = \frac{\bar{B}}{2\Sigma^2}, \quad \Phi_2 = \frac{\bar{B}}{\sqrt{2}\Sigma^3} [a_{\bar{m}} - \bar{S}_p]. \quad (5.13)$$

It follows that the non-null electromagnetic field is aligned with the principal null direction k of the gravitation field, but this vector is not a double degenerate vector of the field.

VI. CONCLUSION

We have derived and analyzed new gyraton solutions moving with the speed of light on electrovacuum Melvin background spacetime in four dimensions. This solution extends the gyraton solutions previously known on the Nariai, anti-Nariai, and Plebański-Hacyan universes of type D, and on conformally flat Bertotti-Robinson and Minkowski space.

The gyraton solutions describe a gravitational field created by a stress-energy tensor of a spinning (circularly polarized) high-frequency beam of electromagnetic radiation, neutrino, or any other massless fields. The gyratons generalize standard gravitational pp waves or Kundt waves by admitting a nonzero angular momentum of the source. The interpretation is that the null matter in the interior of the gyratonic source possesses an intrinsic spin (or nonzero angular momentum). This leads to other nontrivial components of the Einstein equations, namely, G_{ui} in addition to the pure radiation uu component which appears for pp waves or Kundt waves.

We have shown that it is possible to define the gyraton by adding the gyratonic terms a_i to the gravitational wave

on the Melvin spacetime in cylindrical coordinates in a similar way as we have defined them in the general Kundt class. We were able to find an ansatz for the gyraton metric on the Melvin spacetime by direct transformation from the Kundt class of metrics (Sec. II B).

We have further demonstrated that the Einstein-Maxwell equations reduce to the set of linear equations on the 2-dimensional transverse spacetime which has a nontrivial geometry given by the transverse metric. These equations can be solved exactly for any distribution of the matter sources. In general, the problem has been thus reduced to a construction of scalar Green functions for certain differential operators on the transverse space.

We have solved and analyzed the field equations for particular examples with the axial symmetry. In these cases the equations reduce to ordinary differential equations.

We have analyzed geometric properties of the principal null congruence and we have found that it is not recurrent contrary to the case of gyratons on direct-product spacetimes. We have explicitly calculated the curvature tensor and determined that the gyratons on Melvin spacetime are of Petrov type II and belong to the Kundt family of shear-free and twist-free nonexpanding spacetimes. The gyratonic term a_i generates the nontrivial Ricci Φ_{12} and Weyl Ψ_3 scalars, in addition to the gravitational waves investigated in [44]. We found also a very simple relation (5.8) between these components. By studying particular subclasses we have shown that our solutions are generalizations of those from [44].

The scalar polynomial invariants of the metric (2.18) are in general nonconstant (although, some of them are zero)—they depend on the coordinate ρ . The invariants are not affected by the presence of the gyratons, they are the same as for the Melvin background. The same property was proved for gyratons on backgrounds belonging to VSI or CSI families of spacetime.

It would be interesting to investigate a generalization of our ansatz for more complicated spacetimes which could allow, e.g., an inclusion of a cosmological constant.

ACKNOWLEDGMENTS

We wish to thank to Tomáš Pecháček and Otakar Svítek for helpful discussions and to Marcello Ortaggio for his paper about gravitational waves in the Melvin universe which motivated our work. H.K. was supported by Grants No. GAČR-202/09/H033, No. GAUK 12209, and Project No. SVV 261301 of the Charles University in Prague. P.K. was supported by Grant No. GAČR 202/09/0772, and both authors thank the Project No. LC06014 of the Center of Theoretical Astrophysics.

APPENDIX: THE EINSTEIN EQUATIONS

Here we present quantities needed for evaluation of the Einstein equations.

The inverse to the metric (2.18) is

$$\begin{aligned} g^{\mu\nu}\partial_\mu\partial_\nu &= \frac{1}{\Sigma^2}\partial_\rho\partial_\rho + \frac{\Sigma^2}{\rho^2}\partial_\phi\partial_\phi - \frac{2}{\Sigma^2}\partial_u\partial_\nu \\ &+ 2\left(a_\rho\frac{1}{\Sigma^2}\partial_\rho + a_\phi\frac{\Sigma^2}{\rho^2}\partial_\phi\right)\partial_\nu \\ &+ \left(2\frac{H}{\Sigma^2} + a^2\right)\partial_\nu\partial_\nu. \end{aligned} \quad (\text{A1})$$

The stress-energy tensor T^{EM} of the electromagnetic field (2.22) can be defined as

$$T_{\mu\nu}^{\text{EM}} = \frac{\varepsilon_o}{2}\mathcal{F}_{\mu\rho}\bar{\mathcal{F}}_{\nu\rho}, \quad (\text{A2})$$

where $\mathcal{F} \equiv F + i\star F$ is the complex self-dual Maxwell tensor. The 4-dimensional Hodge dual is defined by $\star F_{\mu\nu} = \frac{1}{2}\varepsilon_{\mu\nu\rho\sigma}F^{\rho\sigma}$ and the Maxwell tensor $\mathcal{F}_{\mu\nu}$ satisfies the self-duality condition $\star\mathcal{F} = -i\mathcal{F}$.

The nonvanishing components of the stress-energy tensor (A2) are

$$\begin{aligned} \kappa T_{uv}^{\text{EM}} &= \frac{\varrho_{\text{EM}}}{\Sigma^2}, & \kappa T_{uu}^{\text{EM}} &= 2\varrho_{\text{EM}}\left(\frac{H}{\Sigma^2} + (s-a)^2\right), \\ \kappa T_{u\rho}^{\text{EM}} &= \frac{\varrho_{\text{EM}}}{\Sigma^2}(a_\rho - 2s_\rho), & \kappa T_{u\phi}^{\text{EM}} &= \frac{\varrho_{\text{EM}}}{\Sigma^2}(a_\phi - 2s_\phi), \\ \kappa T_{\rho\rho}^{\text{EM}} &= \frac{\varrho_{\text{EM}}}{\Sigma^2} = \frac{\varrho_{\text{EM}}}{\Sigma^4}g_{\rho\rho}, & \kappa T_{\phi\phi}^{\text{EM}} &= \frac{\varrho_{\text{EM}}\rho^2}{\Sigma^6} = \frac{\varrho_{\text{EM}}}{\Sigma^4}g_{\phi\phi}, \end{aligned} \quad (\text{A3})$$

where the density ϱ_{EM} was defined in (2.4).

The nonvanishing components of the stress-energy tensor (2.27) of the gyratonic matter are

$$\kappa T_{uu}^{\text{gyr}} = j_u = v \operatorname{div}(\Sigma^2 j) + \iota, \quad \kappa T_{ui}^{\text{gyr}} = j_i. \quad (\text{A4})$$

The Einstein tensor for the metric (2.18) reads

$$\begin{aligned} G_{uv} &= \frac{1}{\Sigma^2\rho}(-\rho(\Sigma, \rho)^2 + 2\Sigma(\Sigma, \rho)), \\ G_{uu} &= \frac{1}{2}\Sigma^4 b^2 + \Sigma^2\left(\Delta H + \frac{(\Sigma^2)_{,\rho}}{\Sigma^4}H, \rho\right) \\ &+ \Sigma^2(\partial_\nu^2 H)a^2 + 2\Sigma^2 a^i \partial_\nu H, i + (\partial_\nu H + \partial_u)\operatorname{div}(\Sigma^2 a) \\ &+ 2HG_{uv}, \\ G_{u\rho} &= \frac{1}{2}\frac{\Sigma^4}{\rho}b_{,\phi} - a_\rho(G_{uv} - \partial_\nu^2 H) + \partial_\nu H, \rho, \\ G_{u\phi} &= -\frac{1}{2}\rho\left(b_{,\rho} + \frac{4\Sigma_{,\rho}}{\Sigma}b\right) - a_\phi(G_{uv} - \partial_\nu^2 H) + \partial_\nu H, \phi, \\ G_{\rho\rho} &= G_{uv} + \partial_\nu^2 H, \quad G_{\phi\phi} = \frac{\rho^2}{\Sigma^4}(G_{uv} + \partial_\nu^2 H). \end{aligned} \quad (\text{A5})$$

Here we have used only the metric (2.18), without any usage of the field equations.

-
- [1] R. C. Tolman, *Relativity, Thermodynamics, and Cosmology* (Clarendon Press, Oxford, 1934).
- [2] A. Peres, *Phys. Rev.* **118**, 1105 (1960).
- [3] W. B. Bonnor, *Commun. Math. Phys.* **13**, 163 (1969).
- [4] W. B. Bonnor, *Int. J. Theor. Phys.* **2**, 373 (1969).
- [5] W. B. Bonnor, *Int. J. Theor. Phys.* **3**, 57 (1970).
- [6] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers, and E. Herlt, *Exact Solutions of Einstein's Field Equations* (Cambridge University Press, Cambridge, England, 2003).
- [7] J. B. Griffiths and J. Podolský, *Exact Space-Times in Einstein's General Relativity* (Cambridge University Press, Cambridge, England, 2009).
- [8] P. C. Aichelburg and R. U. Sexl, *Gen. Relativ. Gravit.* **2**, 303 (1971).
- [9] V. Ferrari and P. Pendenza, *Gen. Relativ. Gravit.* **22**, 1105 (1990).
- [10] C. O. Loustó and N. Sánchez, *Nucl. Phys.* **B383**, 377 (1992).
- [11] M. Hotta and M. Tanaka, *Classical Quantum Gravity* **10**, 307 (1993).
- [12] H. Balasin and H. Nachbagauer, *Classical Quantum Gravity* **12**, 707 (1995).
- [13] H. Balasin and H. Nachbagauer, *Classical Quantum Gravity* **13**, 731 (1996).
- [14] J. Podolský and J. B. Griffiths, *Phys. Rev. D* **56**, 4756 (1997).
- [15] J. Podolský and J. B. Griffiths, *Phys. Rev. D* **58**, 124024 (1998).
- [16] J. Podolský, in *Gravitation: Following the Prague Inspiration*, edited by O. Semerák, J. Podolský, and M. Žofka (World Scientific, Singapore, 2002), pp. 205–246.
- [17] C. Barrabès and P. A. Hogan, *Singular Null Hypersurfaces in General Relativity* (World Scientific, Singapore, 2003).
- [18] W. B. Bonnor, *Int. J. Theor. Phys.* **3**, 257 (1970).
- [19] J. B. Griffiths, *Int. J. Theor. Phys.* **5**, 141 (1972).
- [20] H. W. Brinkmann, *Math. Ann.* **94**, 119 (1925).
- [21] V. P. Frolov and D. V. Fursaev, *Phys. Rev. D* **71**, 104034 (2005).
- [22] V. P. Frolov, W. Israel, and A. Zelnikov, *Phys. Rev. D* **72**, 084031 (2005).
- [23] V. Pravda, A. Pravdova, A. Coley, and R. Milson, *Classical Quantum Gravity* **19**, 6213 (2002).
- [24] D. N. Page, *Classical Quantum Gravity* **26**, 055016 (2009).

- [25] V. P. Frolov and A. Zelnikov, *Classical Quantum Gravity* **23**, 2119 (2006).
- [26] V. P. Frolov and A. Zelnikov, *Phys. Rev. D* **72**, 104005 (2005).
- [27] S. T. C. Siklos, *Galaxies, Axisymmetric Systems and Relativity*, edited by M. A. H. MacCallum (Cambridge University Press, Cambridge, England, 1985).
- [28] J. Podolský, *Classical Quantum Gravity* **15**, 719 (1998).
- [29] A. A. Coley, S. Hervik, and N. Pelavas, *Classical Quantum Gravity* **23**, 3053 (2006).
- [30] A. A. Coley, G. W. Gibbons, S. Hervik, and C. N. Pope, *Classical Quantum Gravity* **25**, 145017 (2008).
- [31] A. A. Coley, S. Hervik, and N. Pelavas, *Classical Quantum Gravity* **25**, 025008 (2008).
- [32] A. A. Coley, S. Hervik, and N. Pelavas, *Classical Quantum Gravity* **26**, 025013 (2009).
- [33] H. Kadlecová, A. Zelnikov, P. Krtouš, and J. Podolský, *Phys. Rev. D* **80**, 024004 (2009).
- [34] V. P. Frolov and F.-L. Lin, *Phys. Rev. D* **73**, 104028 (2006).
- [35] M. M. Caldarelli, D. Klemm, and E. Zorzan, *Classical Quantum Gravity* **24**, 1341 (2007).
- [36] H. Yoshino and V. S. Rychkov, *Phys. Rev. D* **71**, 104028 (2005).
- [37] H. Yoshino and R. B. Mann, *Phys. Rev. D* **74**, 044003 (2006).
- [38] H. Yoshino, A. Zelnikov, and V. Frolov, *Phys. Rev. D* **75**, 124005 (2007).
- [39] W. B. Bonnor, *Proc. Phys. Soc. London Sect. A* **67**, 225 (1954).
- [40] M. A. Melvin, *Phys. Rev.* **139**, B225 (1965).
- [41] K. S. Thorne, *Phys. Rev.* **139**, B244 (1965).
- [42] Melvin and Wallingford, *J. Math. Phys. (N.Y.)* **7**, 333 (1966).
- [43] D. Garfinkle and M. A. Melvin, *Phys. Rev. D* **45**, 1188 (1992).
- [44] M. Ortaggio, *Phys. Rev. D* **69**, 064034 (2004).
- [45] L. Havrdová and P. Krtouš, *Gen. Relativ. Gravit.* **39**, 291 (2007).
- [46] G. W. Gibbons and K. Maeda, *Nucl. Phys.* **B298**, 741 (1988).
- [47] G. W. Gibbons and C. A. R. Herdeiro, *Classical Quantum Gravity* **18**, 1677 (2001).
- [48] G. W. Gibbons, in *Fields and Geometry: Proceedings of the XXII Winter School of Theoretical Physics*, edited by A. Jadczyk (World Scientific, Singapore, 1986).
- [49] D. Garfinkle and A. Strominger, *Phys. Lett. B* **256**, 146 (1991).
- [50] D. Garfinkle, S. B. Giddings, and A. Strominger, *Phys. Rev. D* **49**, 958 (1994).
- [51] F. Dowker, J. P. Gauntlett, D. A. Kastor, and J. Traschen, *Phys. Rev. D* **49**, 2909 (1994).
- [52] F. Dowker, J. P. Gauntlett, S. B. Giddings, and G. T. Horowitz, *Phys. Rev. D* **50**, 2662 (1994).
- [53] S. W. Hawking and S. F. Ross, *Phys. Rev. D* **52**, 5865 (1995).
- [54] R. Emparan, *Phys. Rev. Lett.* **75**, 3386 (1995).
- [55] J. F. Plebanski, *J. Math. Phys.* **20**, 1946 (1979).
- [56] A. A. Coley, S. Hervik, and N. Pelavas, *Classical Quantum Gravity* **27**, 102001 (2010).