

Ultrarelativistic boost of spinning black rings

Marcello Ortaggio,* Jiří Podolský and Pavel Krtouš

*Institute of Theoretical Physics, Faculty of Mathematics and Physics
Charles University in Prague
V Holešovičkách 2, 180 00 Prague 8, Czech Republic
E-mail: marcello.ortaggio@comune.re.it, Jiri.Podolsky@mff.cuni.cz,
Pavel.Krtous@mff.cuni.cz*

ABSTRACT: We study the $D = 5$ Emparan-Reall spinning black ring under an ultrarelativistic boost along an arbitrary direction. We analytically determine the resulting shock pp -wave, in particular for boosts along axes orthogonal and parallel to the plane of rotation. The solution becomes physically more interesting and simpler if one enforces equilibrium between the forces on the ring. We also comment on the ultrarelativistic limit of recently found supersymmetric black rings with two independent angular momenta. Essential distinct features with respect to the boosted Myers-Perry black holes are pointed out.

KEYWORDS: Black Holes, Classical Theories of Gravity.

*Also at INFN, Rome, Italy.

Contents

1. Introduction	1
2. The black ring solution	2
3. General boost	3
4. Orthogonal boost: $\alpha = 0$	6
5. Parallel boost: $\alpha = \pi/2$	8
6. Boost of the supersymmetric black ring	11
A. Results for the boosted $D = 5$ Myers-Perry black hole	12
A.1 Orthogonal boost	12
A.2 Parallel boost	13

1. Introduction

Shock pp -wave geometries describe the spacetime surrounding very fast moving objects, and are thus relevant to the study of planckian scattering [1]. They are also of interest in string theory, since strings may be exactly solved in such backgrounds [2, 3]. The prototype of shock wave solutions is the Aichelburg-Sexl spacetime, which represents the gravitational field of a massless point particle. It was originally obtained by boosting the Schwarzschild black hole to the speed of light, while rescaling the mass to zero in an appropriate way [4]. According to recent extra-dimension scenarios, the fundamental Planck scale of (higher dimensional) gravity could be as low as a few TeV. This has stimulated renewed interest in the study of gravitational effects in high energy collisions, especially in view of the possible observation of microscopic black holes at near future colliders [5–8] (see, e.g., [9] for a recent review and for further references). It has been shown that closed trapped surfaces do indeed form in the ultrarelativistic collision of Aichelburg-Sexl point particles [10, 11] and of finite-size beams [12], which can more accurately model string-size effects. Nevertheless, it is desirable to understand how other effects could influence high energy scattering. A first step in this direction is to investigate more general shock wave solutions of higher dimensional gravity, which can naturally be obtained by applying the boosting technique of [4] to black hole spacetimes. This has been done in any $D \geq 4$ for static black holes with

electric charge [13] or immersed in an external magnetic field [14, 15]. The ultrarelativistic limit of the Myers-Perry rotating black holes [16] has been studied in [17] (for the case of one non-vanishing spin). However, a striking feature of General Relativity in $D > 4$ is the non-uniqueness of the spherical black holes of [16]. In five-dimensional vacuum gravity, there exist also asymptotically flat rotating black rings with an event horizon of topology $S^1 \times S^2$ [18]. In the present contribution, we aim at studying the gravitational field generated by such rings in the Aichelburg-Sexl limit. As we will see in detail, this results in shock waves generated by extended lightlike sources (with a characteristic length-scale) which are remnants of the ring singularity of the original spacetime [18]. Our recent results on boosted non-rotating black rings [19] will be recovered as a special subcase. In general, the presence of spin is important because it allows black rings to be in equilibrium [18] without introducing “unphysical” membranes via conical singularities [20]. This will be reflected also in the shock geometry resulting from the boost. From a supergravity and string theory point of view, it is remarkable that supersymmetric black rings have been also constructed [21–24]. We will conclude this article with a brief comment on the boost of such solutions. In the Appendix, we compare our results with those obtained for the ultrarelativistic limit of Myers-Perry black holes [17] in $D = 5$.

2. The black ring solution

In this section we briefly summarize the basic properties of the black ring, referring to [18, 25] for details. In the coordinates of [25],¹ the line element reads

$$\begin{aligned}
 ds^2 = & -\frac{F(y)}{F(x)} \left(dt + C(\nu, \lambda) L \frac{1+y}{F(y)} d\psi \right)^2 + \\
 & + \frac{L^2}{(x-y)^2} F(x) \left[-\frac{G(y)}{F(y)} d\psi^2 - \frac{dy^2}{G(y)} + \frac{dx^2}{G(x)} + \frac{G(x)}{F(x)} d\phi^2 \right], \quad (2.1)
 \end{aligned}$$

where

$$F(\zeta) = \frac{1 + \lambda\zeta}{1 - \lambda}, \quad G(\zeta) = (1 - \zeta^2) \frac{1 + \nu\zeta}{1 - \nu}, \quad C(\nu, \lambda) = \sqrt{\frac{\lambda(\lambda - \nu)(1 + \lambda)}{(1 - \nu)(1 - \lambda)^3}}. \quad (2.2)$$

The dimensionless parameters λ and ν satisfy $0 \leq \nu \leq \lambda < 1$, and for $\lambda = 0 = \nu$ the spacetime (2.1) is flat. The constant $L > 0$ represents a length related to the radius of the “central circle” of the ring. For a physical interpretation of the spacetime (2.1) we take $y \in (-\infty, -1]$, $x \in [-1, +1]$ (see a discussion in [26] for other possible choices) and ψ and ϕ as periodic angular coordinates (see below). Surfaces of constant y have topology $S^1 \times S^2$. The coordinate ψ runs along the S^1 factor, whereas (x, ϕ) parametrize S^2 (see [20, 25, 22] for illustrative pictures). Within the above range, y parametrizes “distances” from the ring circle. At $y \rightarrow -\infty$ the spacetime has a inner spacelike curvature singularity, $y = -1/\nu$

¹Up to simple constant rescalings of $F(\zeta)$, $G(\zeta)$, $C(\lambda, \nu)$, ψ and ϕ , cf. eqs. (2.2) and (2.4) with the corresponding ones in [25]. In addition, multiply our L^2 by $(1 - \nu)/(1 - \lambda)$ to obtain the parameter used in [25].

is a horizon and $y = -1/\lambda$ an ergosurface, both with topology $S^1 \times S^2$. The black ring solution (2.1) is asymptotically flat near spatial infinity $x, y \rightarrow -1$, where it tends to Minkowski spacetime in the form

$$ds_0^2 = -dt^2 + \frac{L^2}{(x-y)^2} \left[(y^2 - 1)d\psi^2 + \frac{dy^2}{y^2 - 1} + \frac{dx^2}{1 - x^2} + (1 - x^2)d\phi^2 \right]. \quad (2.3)$$

To avoid conical singularities at the axes $x = -1$ and $y = -1$, the angular coordinates must have the standard periodicity

$$\Delta\phi = 2\pi = \Delta\psi. \quad (2.4)$$

Centrifugal repulsion and gravitational self-attraction of the ring are in balance if conical singularities are absent also at $x = +1$, which requires

$$\lambda = \frac{2\nu}{1 + \nu^2}. \quad (2.5)$$

When this equilibrium condition holds, the metric (2.1) is a vacuum solution (of $D = 5$ General Relativity) everywhere. With different choices (e.g., in the static limit $\nu = \lambda$ [20]), the conical singularity at $x = +1$ describes a disk-shaped membrane inside the ring.

The mass, angular momentum and angular velocity (at the horizon) of the black ring are

$$M = \frac{3\pi L^2}{4} \frac{\lambda}{1 - \lambda}, \quad J = \frac{\pi L^3}{2} \sqrt{\frac{\lambda(\lambda - \nu)(1 + \lambda)}{(1 - \nu)(1 - \lambda)^3}}, \quad \Omega = \frac{1}{L} \sqrt{\frac{(\lambda - \nu)(1 - \lambda)}{\lambda(1 + \lambda)(1 - \nu)}}. \quad (2.6)$$

The algebraic type of the Weyl tensor of the ring spacetime is I_i [26].

3. General boost

For our purposes, it is convenient to decompose the line element (2.1) as

$$ds^2 = ds_0^2 + \Delta, \quad (3.1)$$

in which ds_0^2 is Minkowski spacetime (2.3) and

$$\begin{aligned} \Delta = & \lambda \frac{x-y}{1+\lambda x} dt^2 - 2(1-\lambda)C(\lambda, \nu)L \frac{1+y}{1+\lambda x} dt d\psi \\ & + \frac{\lambda - \nu}{1 - \nu} \frac{L^2}{1 + \lambda y} \left[-\lambda \frac{1 + \lambda}{1 - \lambda} \frac{(1 + y)^2}{1 + \lambda x} + \frac{y^2 - 1}{x - y} \right] d\psi^2 + \\ & + \frac{L^2}{(x - y)^2} \left[\nu \frac{x + 1}{1 - \nu} (y^2 - 1) d\psi^2 + \frac{\lambda(1 - \nu)(x - y) + (\lambda - \nu)(1 + y)}{(1 - \lambda)(1 + \nu y)} \frac{dy^2}{y^2 - 1} + \right. \\ & \left. + \frac{\lambda - \nu}{1 - \lambda} \frac{dx^2}{(1 - x)(1 + \nu x)} + \nu \frac{x + 1}{1 - \nu} (1 - x^2) d\phi^2 \right]. \end{aligned} \quad (3.2)$$

The above splitting is such that near infinity ($x, y \rightarrow -1$) one has $ds^2 \rightarrow ds_0^2$, while Δ becomes “negligible” (in the sense of the “background” metric ds_0^2). This enables us to define a notion of Lorentz boost using the symmetries of the asymptotic minkowskian

background ds_0^2 . Cartesian coordinates will visualize it most naturally. These can be introduced in two steps. First, we replace the coordinates (y, x) with new coordinates (ξ, η) via the substitution

$$y = -\frac{\xi^2 + \eta^2 + L^2}{\Sigma}, \quad x = -\frac{\xi^2 + \eta^2 - L^2}{\Sigma}, \quad (3.3)$$

where

$$\Sigma = \sqrt{(\eta^2 + \xi^2 - L^2)^2 + 4L^2\eta^2}. \quad (3.4)$$

The flat term ds_0^2 in eq. (3.1) now takes the form $ds_0^2 = -dt^2 + d\eta^2 + \eta^2 d\phi^2 + d\xi^2 + \xi^2 d\psi^2$. Then, cartesian coordinates adapted to the Killing vectors ∂_ϕ and ∂_ψ are given by

$$x_1 = \eta \cos \phi, \quad x_2 = \eta \sin \phi, \quad y_1 = \xi \cos \psi, \quad y_2 = \xi \sin \psi, \quad (3.5)$$

so that $\eta = \sqrt{x_1^2 + x_2^2}$, $\xi = \sqrt{y_1^2 + y_2^2}$, and $ds_0^2 = -dt^2 + dx_1^2 + dx_2^2 + dy_1^2 + dy_2^2$. This enables us to study a boost along a general direction. Since the original spacetime (2.1) is symmetric under (separate) rotations in the (x_1, x_2) and (y_1, y_2) planes, such a direction can be specified by a single parameter α , namely introducing rotated axes z_1 and z_2

$$x_1 = z_1 \cos \alpha - z_2 \sin \alpha, \quad y_1 = z_1 \sin \alpha + z_2 \cos \alpha. \quad (3.6)$$

Defining now suitable double null coordinates (u', v') by

$$t = \frac{-u' + v'}{\sqrt{2}}, \quad z_1 = \frac{u' + v'}{\sqrt{2}}, \quad (3.7)$$

a Lorentz boost along z_1 takes the simple form

$$u' = \epsilon^{-1}u, \quad v' = \epsilon v. \quad (3.8)$$

The parameter $\epsilon > 0$ is related to the standard Lorentz factor via $\gamma = (\epsilon + \epsilon^{-1})/2$. We are interested in “ultrarelativistic” boosts to the speed of light, i.e. in taking the limit $\epsilon \rightarrow 0$ in the transformation (3.8). While $\epsilon \rightarrow 0$, we will rescale the mass as $M = \gamma^{-1}p_M \approx 2\epsilon p_M$ [4], which physically means that the total energy remains finite in the limit ($p_M > 0$ is a constant). Moreover, during the ultrarelativistic limit we wish to keep the angular velocity Ω finite (a similar condition was imposed in [17]), and to allow for the possibility of black rings in equilibrium (when the condition (2.5) holds). From eq. (2.6), these requirements imply the rescalings²

$$\lambda = \epsilon p_\lambda, \quad \nu = \epsilon p_\nu, \quad (3.9)$$

where $p_\lambda = 8p_M/(3\pi L^2)$ and p_ν is another positive constant such that $p_\lambda \geq p_\nu$. In terms of these parameters, for $\epsilon \rightarrow 0$ the equilibrium condition (2.5) becomes

$$p_\lambda = 2p_\nu. \quad (3.10)$$

²This appears to be physically the most interesting and simple choice. See Footnote 3 for a subtler, slightly more general comment.

Values $p_\nu \leq p_\lambda < 2p_\nu$ correspond to black rings (2.1) which are “underspinning” before the boost (and therefore balanced by a membrane of negative energy density), values $p_\lambda > 2p_\nu$ to “overspinning” black rings (with a membrane of positive energy density). Notice, however, that under the limit $\epsilon \rightarrow 0$ the angular momentum J will tend to zero (as $\sim \epsilon$).

We can now evaluate how the black ring metric (2.1) (that is, eq. (3.1) with eqs. (2.3) and (3.2)) transforms under the boost (3.8). We have first to substitute eq. (3.3) into eqs. (2.3) and (3.2). Then, we apply the sequence of substitutions (3.5)–(3.7) into the thus obtained expressions for ds_0^2 and for Δ . Finally, we perform the boost (3.8) with the rescalings (3.9), which make $\Delta = \Delta_\epsilon$ dependent on ϵ . The ds_0^2 is invariant under the boost and at the end it reads

$$ds_0^2 = 2dudv + dx_2^2 + dy_2^2 + dz_2^2. \tag{3.11}$$

The next step is to take the ultrarelativistic limit $ds^2 = ds_0^2 + \lim_{\epsilon \rightarrow 0} \Delta_\epsilon$. This is delicate because the expansion of Δ_ϵ in ϵ has a different structure in different regions of the spacetime (even away from the singularity $y = -\infty$). In particular, a peculiar behaviour is obtained for $u = 0$, because Δ_ϵ depends on u through the combination

$$z_\epsilon = \frac{1}{\sqrt{2}}(\epsilon^{-1}u + \epsilon v). \tag{3.12}$$

In order to have control over the exact distributional structure of the limit, it is convenient to isolate such dependence on $\epsilon^{-1}u$ by performing first an expansion of Δ_ϵ with z_ϵ unexpanded. This leads to an expression

$$\Delta_\epsilon = \frac{1}{\epsilon}h(z_\epsilon)du^2 + [k_1(z_\epsilon)dx_2 + k_2(z_\epsilon)dy_2 + k_3(z_\epsilon)dz_2 + k_4(z_\epsilon)du] du + \dots, \tag{3.13}$$

where the dots denote terms proportional to higher powers of ϵ , which are negligible in the limit. We have emphasized here the dependence of the functions h and k_i ($i = 1, \dots, 4$) on z_ϵ (and thus on ϵ), because this is essential in our limit, but they depend also on x_2 , y_2 and z_2 . The quantities k_i are rather involved, but it suffices to observe here that $\lim_{\epsilon \rightarrow 0} k_i(z_\epsilon) = 0$. We can thus also drop all the terms of order ϵ^0 in (3.13).³ For h , after all the steps described above, we obtain explicitly

$$\begin{aligned} h(z_\epsilon) = & p_\lambda \frac{L^2}{\Sigma} + p_\nu \frac{L^2}{\Sigma^3} \left[(\xi^2 - \eta^2 - L^2) \frac{y_1}{\xi} \sin \alpha + 2\xi x_1 \cos \alpha \right]^2 + \\ & + \frac{1}{2}(2p_\nu - p_\lambda) \left(1 - \frac{\xi^2 + \eta^2 - L^2}{\Sigma} \right) \left(\frac{y_2^2}{\xi^2} \sin^2 \alpha + \frac{x_2^2}{\eta^2} \cos^2 \alpha \right) + \\ & + \sqrt{p_\lambda(p_\lambda - p_\nu)} \frac{Ly_2 \sin \alpha}{\xi^2} \left(-1 + \frac{\xi^2 + \eta^2 + L^2}{\Sigma} \right) + (p_\lambda - p_\nu) \frac{L^2 y_2^2}{\xi^2 \Sigma} \sin^2 \alpha + \\ & + \frac{1}{2}(p_\lambda - p_\nu) \left(1 - \frac{\xi^2 + \eta^2 - L^2}{\Sigma} \right). \end{aligned} \tag{3.14}$$

³A remark on the “triviality” of the ϵ^0 terms is in order, since they could be non-vanishing for certain more general scalings of the original metric parameters. While with higher order (in ϵ) corrections in eq. (3.9) $\lim_{\epsilon \rightarrow 0} k_i(z_\epsilon) = 0$ would still hold, we could introduce a non-vanishing contribution by allowing an ϵ -dependence in the ring “radius” via $L_\epsilon = L + c_1\epsilon + c_2\epsilon^2 + \dots$. The convergence of the integral (3.17) would then require $c_1 = 0$, but the quantity $c_2\epsilon^2$ would affect the limit of (3.13) via $\lim_{\epsilon \rightarrow 0} k_4(z_\epsilon) = c_2$. The resulting term $c_2 du^2$ is, however, obviously removable with a coordinate transformation.

Recall that the dependence of h on ϵ is contained in x_1 and y_1 via eqs. (3.6)–(3.8), in η and ξ via eq. (3.5) and in Σ via eq. (3.4). In taking the limit $\epsilon \rightarrow 0$ of eq. (3.13), we apply the distributional identity

$$\lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} f(z_\epsilon) = \sqrt{2} \delta(u) \int_{-\infty}^{+\infty} f(z) dz. \quad (3.15)$$

The final metric is thus (cf. eqs. (3.11) and (3.13))

$$ds^2 = 2dudv + dx_2^2 + dy_2^2 + dz_2^2 + H(x_2, y_2, z_2) \delta(u) du^2, \quad (3.16)$$

with a profile function given by

$$H(x_2, y_2, z_2) = \sqrt{2} \int_{-\infty}^{+\infty} h(z) dz. \quad (3.17)$$

A black ring boosted to the speed of light in a general direction z_1 is thus described by the metric (3.16) with eq. (3.17). This is evidently a $D = 5$ impulsive pp -wave with wave vector ∂_v . Such a spacetime is flat everywhere except on the null hyperplane $u = 0$, which represents the impulsive wave front. Note that the equilibrium condition (3.10) has not yet been enforced in the above expression for h (in particular, in the static limit $p_\nu = p_\lambda$ we recover the result of [19]). In order to write the solutions in a completely explicit form, it only remains to perform the integration in eq. (3.17), with h given by eq. (3.14) with eqs. (3.4)–(3.8) and (3.12). For any α , this integral is always convergent and can in principle be expressed using elliptic integrals (because Σ is a square root of a fourth order polynomial in z , see [19] for related comments). Therefore, no singular coordinate transformation of the type of [4] has to be performed. In the following, we will explicitly calculate the integral, and study the corresponding solution in the case of two different boosts of the black ring along the privileged axes x_1 ($\alpha = 0$) and y_1 ($\alpha = \pi/2$), which are respectively “orthogonal” and “parallel” to the 2-plane (y_1, y_2) (i.e., (ξ, ψ)) in which the ring rotates.

4. Orthogonal boost: $\alpha = 0$

For the orthogonal boost $\alpha = 0$, from eq. (3.6) one has $z_1 = x_1$ and $z_2 = y_1$, so that the general pp -wave (3.16) reduces to

$$ds^2 = 2dudv + dx_2^2 + dy_1^2 + dy_2^2 + H_\perp(x_2, y_1, y_2) \delta(u) du^2. \quad (4.1)$$

Also, it is now convenient to rewrite h in eq. (3.14) as

$$h_\perp(z_\epsilon) = [3p_\lambda L^2 - (p_\lambda - p_\nu) \xi^2 - p_\nu (x_2^2 + L^2)] \frac{1}{2\Sigma} + p_\nu \frac{4L^2 \xi^2 z_\epsilon^2}{\Sigma^3} + \frac{1}{2} (2p_\nu - p_\lambda) \left[\frac{x_2^2 (L^2 - \xi^2)}{(z_\epsilon^2 + x_2^2) \Sigma} + \frac{x_2^2}{z_\epsilon^2 + x_2^2} \right] + \frac{1}{2} (p_\lambda - p_\nu) \left(1 - \frac{z_\epsilon^2}{\Sigma} \right), \quad (4.2)$$

and Σ (from eq. (3.4)) as

$$\Sigma = \sqrt{[z_\epsilon^2 + x_2^2 + (\xi + L)^2] [z_\epsilon^2 + x_2^2 + (\xi - L)^2]}. \quad (4.3)$$

Hereafter, it is understood that $\xi = \sqrt{y_1^2 + y_2^2}$. In the orthogonal boost there is no contribution to h_{\perp} from the off-diagonal term $g_{t\psi}$ in the metric (2.1). Performing the integration (3.17) with h given by eqs. (4.2) and (4.3), we find

$$\begin{aligned}
 H_{\perp}(x_2, y_1, y_2) = & \sqrt{2} \frac{3p_{\lambda}L^2 + (2p_{\nu} - p_{\lambda})\xi^2}{\sqrt{(\xi + L)^2 + x_2^2}} K(k) + \sqrt{2}(2p_{\nu} - p_{\lambda}) \times \\
 & \times \left[-\sqrt{(\xi + L)^2 + x_2^2} E(k) + \frac{\xi - L}{\xi + L} \frac{x_2^2}{\sqrt{(\xi + L)^2 + x_2^2}} \Pi(\rho, k) + \right. \\
 & \left. + \pi|x_2|\Theta(L - \xi) \right], \tag{4.4}
 \end{aligned}$$

where

$$k = \sqrt{\frac{4\xi L}{(\xi + L)^2 + x_2^2}}, \quad \rho = \frac{4\xi L}{(\xi + L)^2}, \tag{4.5}$$

and $\Theta(L - \xi)$ denotes the step function. In the above calculation, we have used the standard elliptic integrals and their properties summarized in the Appendix of [19], and the additional integral (Σ given by eq. (4.3) with z_{ϵ} replaced by z)

$$\int_0^{\infty} \left(1 - \frac{z^2}{\Sigma}\right) dz = \sqrt{(\xi + L)^2 + x_2^2} E(k). \tag{4.6}$$

In order to gain physical insight, it is useful to visualize the behaviour of the gravitational field at a large spatial distance within the wave front $u = 0$. Defining the coordinates (r, θ)

$$x_2 = r \cos \theta, \quad \xi = r \sin \theta, \tag{4.7}$$

an expansion for small values of the dimensionless parameter L/r (using the identities summarized in [19]) leads to

$$\begin{aligned}
 H_{\perp} = & \frac{\pi}{\sqrt{2}} p_{\lambda} L \left[3 \frac{L}{r} - \left(\frac{5}{8} + \frac{p_{\nu}}{4p_{\lambda}} \right) (3 \cos^2 \theta - 1) \frac{L^3}{r^3} + \right. \\
 & \left. + \left(\frac{7}{64} + \frac{p_{\nu}}{16p_{\lambda}} \right) (35 \cos^4 \theta - 30 \cos^2 \theta + 3) \frac{L^5}{r^5} + O\left(\frac{L^7}{r^7}\right) \right]. \tag{4.8}
 \end{aligned}$$

We recognize the standard form of multipole terms. The monopole is essentially an Aichelburg-Sexl term. The dipole and the octupole are missing, due to the geometry of the source. The quadrupole and 16-pole reflect the shape of the singularity and depend on the spin of the original black ring, but they persist even in the static limit $p_{\nu} = p_{\lambda}$ [19] (when, in fact, they reach their maximal strength).

It is remarkable that for the physically more interesting case of *black rings in equilibrium*, i.e. those satisfying $p_{\lambda} = 2p_{\nu}$ (see eq. (3.10)), the profile function simplifies significantly to

$$H_{\perp}^e(x_2, y_1, y_2) = \frac{3\sqrt{2} p_{\lambda} L^2}{\sqrt{(\xi + L)^2 + x_2^2}} K(k). \tag{4.9}$$

Interestingly, this is just the newtonian potential generated by a *uniform ring of radius L* and linear density $\mu = 3\sqrt{2}p_{\lambda}L/4$ located at $x_2 = 0$ in the flat three-dimensional space (x_2, y_1, y_2) . Since for a general pp -wave (4.1) the only component of the Ricci tensor is




Figure 1: The profile function H_{\perp} , given by eq. (4.4), in the case of underspinning ($p_{\lambda} < 2p_{\nu}$, left), overspinning ($p_{\lambda} > 2p_{\nu}$, right) and balanced ($p_{\lambda} = 2p_{\nu}$, bottom) black rings (cf. eq. (3.10)) boosted along an orthogonal direction x_1 . It is represented over the plane (x_2, ξ) (cf. eq. (3.5)), and the Killing coordinate ψ is suppressed. In the equilibrium case, H_{\perp} reduces to H_{\perp}^e of eq. (4.9) and the disk membrane at $x_2 = 0$, $\xi < L$ disappears (no jump of $\partial H_{\perp}^e / \partial x_2$ occurs at $x_2 = 0$). In all cases, there is a ring singularity at $x_2 = 0$, $\xi = L$, as indicated by the thick points in the pictures.

$R_{uu} = -\frac{1}{2}\delta(u)\Delta H_{\perp}$, Δ denoting the Laplace operator over the transverse space (x_2, y_1, y_2) , it follows that the profile function (4.9) represents a spacetime which is vacuum everywhere except on the circle $u = 0 = x_2$, $\xi = L$ (so that $k = 1$ in eq. (4.5)). This lies on the wave front and corresponds to a singular ring-shaped source moving with the speed of light. It is obviously a remnant of the curvature singularity ($y = -\infty$) of the original stationary black ring (2.1). For the non-equilibrium solution (4.4), the discontinuous term proportional to $\Theta(L - \xi)$ is responsible for a disk membrane supporting the ring [19]. We have plotted typical profile functions H_{\perp} and H_{\perp}^e in figure 1.

5. Parallel boost: $\alpha = \pi/2$

For the parallel boost $\alpha = \pi/2$, from eq. (3.6) one has $z_1 = y_1$ and $z_2 = -x_1$, and the general pp -wave (3.16) reduces to

$$ds^2 = 2dudv + dx_1^2 + dx_2^2 + dy_2^2 + H_{\parallel}(x_1, x_2, y_2)\delta(u)du^2. \tag{5.1}$$




Figure 2: Plot of the profile function $H_{||}^e$, given by eq. (5.10), for balanced black rings ($p_\lambda = 2p_\nu$) boosted along the direction y_1 in the plane of rotation. It is depicted over the plane (y_2, η) (cf. eq. (3.5)), and the Killing coordinate ϕ is suppressed. The profile function $H_{||}^e$ diverges at the rod singularity $\eta = 0$, $|y_2| \leq L$, as indicated by the thick line. The two smaller pictures represent the symmetric and antisymmetric part (with respect to the origin of the y_2 -axis) of $H_{||}^e$, respectively. The case of unbalanced black rings, eq. (5.5), does not produce qualitative changes, since the disk membrane Lorentz-contracts to the singular rod region.

The function h can be reexpressed as

$$\begin{aligned}
 h_{||}(z_\epsilon) = & \left[(3p_\lambda + p_\nu)L^2 - p_\nu y_2^2 + 2\sqrt{p_\lambda(p_\lambda - p_\nu)}Ly_2 - (p_\lambda - p_\nu)\eta^2 \right] \frac{1}{2\Sigma} - p_\nu \frac{4L^2\eta^2 z_\epsilon^2}{\Sigma^3} + \\
 & + \frac{1}{2} \left[(2p_\nu - p_\lambda)y_2^2 - 2\sqrt{p_\lambda(p_\lambda - p_\nu)}Ly_2 \right] \left[-\frac{L^2 + \eta^2}{(z_\epsilon^2 + y_2^2)\Sigma} + \frac{1}{z_\epsilon^2 + y_2^2} \right] + \\
 & + \frac{1}{2}(p_\lambda - p_\nu) \left(1 - \frac{z_\epsilon^2}{\Sigma} \right), \tag{5.2}
 \end{aligned}$$

and

$$\Sigma = \sqrt{z_\epsilon^4 + 2(y_2^2 + \eta^2 - L^2)z_\epsilon^2 + a^4}, \tag{5.3}$$

with

$$a = [(\eta^2 + y_2^2 - L^2)^2 + 4\eta^2 L^2]^{1/4}. \tag{5.4}$$

It is understood that $\eta = \sqrt{x_1^2 + x_2^2}$. Performing the integration (3.17) with h given by eqs. (5.2)–(5.4), one obtains

$$\begin{aligned}
 H_{||}(x_1, x_2, y_2) = & \left[2(2p_\lambda - p_\nu)L^2 + (2p_\nu - p_\lambda)a^2 \left(1 + \frac{L^2 + \eta^2}{a^2 - y_2^2} \right) + \right. \\
 & \left. + 2\sqrt{p_\lambda(p_\lambda - p_\nu)}Ly_2 \left(1 - \frac{L^2 + \eta^2}{a^2 - y_2^2} \right) \right] \frac{\sqrt{2}}{a} K(k) - 2\sqrt{2}(2p_\nu - p_\lambda)aE(k) + \\
 & + \frac{\sqrt{2}}{2} \left[(2p_\nu - p_\lambda)y_2 - 2\sqrt{p_\lambda(p_\lambda - p_\nu)}L \right] \times \\
 & \times \left[-\frac{\eta^2 + L^2}{ay_2} \frac{a^2 + y_2^2}{a^2 - y_2^2} \Pi(\rho, k) + \pi \operatorname{sgn}(y_2) \right], \tag{5.5}
 \end{aligned}$$

where

$$k = \frac{(a^2 - \eta^2 - y_2^2 + L^2)^{1/2}}{\sqrt{2}a}, \quad \rho = -\frac{(a^2 - y_2^2)^2}{4a^2 y_2^2}. \quad (5.6)$$

Again, we refer to the [19, appendix], the only additional integral used here being (Σ given by eq. (5.3) with z_ϵ replaced by z)

$$\int_0^\infty \left(1 - \frac{z^2}{\Sigma}\right) dz = 2aE(k) - aK(k). \quad (5.7)$$

With the coordinates

$$y_2 = r \cos \theta, \quad \eta = r \sin \theta, \quad (5.8)$$

the behaviour at large spatial distances is given by

$$\begin{aligned} H_{||} = \frac{\pi}{\sqrt{2}} p_\lambda L \left[3\frac{L}{r} + 2\sqrt{\frac{p_\lambda - p_\nu}{p_\lambda}} \cos \theta \frac{L^2}{r^2} + \left(\frac{7}{8} - \frac{p_\nu}{4p_\lambda}\right) (3 \cos^2 \theta - 1) \frac{L^3}{r^3} + \right. \\ \left. + \frac{3}{4} \sqrt{\frac{p_\lambda - p_\nu}{p_\lambda}} (5 \cos^3 \theta - 3 \cos \theta) \frac{L^4}{r^4} + \right. \\ \left. + \left(\frac{11}{64} - \frac{p_\nu}{16p_\lambda}\right) (35 \cos^4 \theta - 30 \cos^2 \theta + 3) \frac{L^5}{r^5} + O\left(\frac{L^6}{r^6}\right) \right]. \quad (5.9) \end{aligned}$$

Notice that now there appear also a dipole and an octupole term, as a remnant of the angular momentum of the black ring.

We are especially interested in *black rings in equilibrium* (3.10), for which one is left with

$$\begin{aligned} H_{||}^e(x_1, x_2, y_2) = p_\lambda L \left[\frac{3\sqrt{2}L}{a} + \frac{2y_2}{a} \left(1 - \frac{L^2 + \eta^2}{a^2 - y_2^2}\right) \right] K(k) + \\ + p_\lambda L \left[\frac{\eta^2 + L^2}{ay_2} \frac{a^2 + y_2^2}{a^2 - y_2^2} \Pi(\rho, k) - \pi \operatorname{sgn}(y_2) \right]. \quad (5.10) \end{aligned}$$

This function is singular at the points satisfying $u = 0 = \eta$ and $|y_2| \leq L$ ($k = 1$ in eq. (5.6)), i.e. on a *rod of length* $2L$ contained within the wave front. This is a remnant of the curvature singularity of the original stationary black ring (2.1), which has (infinitely) Lorentz-contracted because of the ultrarelativistic boost in the plane of the ring. For the same reason, and because the original ring was rotating, the rod-source corresponding to eq. (5.10) is not uniform. The profile (5.10) corresponds to a vacuum spacetime everywhere except on the rod. Notice also that the apparent divergences of $H_{||}^e$ at $y_2^2 = a^2$ and $y_2 = 0$ is only a fictitious effect: the singular behaviour of the coefficient of Π in eq. (5.10) is exactly compensated from that of K in the first case and from the $\operatorname{sgn}(y_2)$ function in the second case (recall also the form of ρ in eq. (5.6)). Finally, it is interesting to observe that the antisymmetric part (in the coordinate y_2) of $H_{||}$ and $H_{||}^e$ comes entirely from the off-diagonal term $g_{t\psi}$ in the metric (2.1), which was responsible for rotation before the boost (and produces the terms even in $\cos \theta$ in the expansion (5.9)). The profile function $H_{||}^e$ is plotted in figure 2.

6. Boost of the supersymmetric black ring

To conclude, we demonstrate that the above method can also be employed to calculate the gravitational field generated by other black rings in the ultrarelativistic limit. The first supersymmetric black ring (solution of $D = 5$ minimal supergravity) was presented in [21] (and subsequently generalized in [22–24]). The line element reads

$$ds^2 = -f^2(dt + \omega_\psi d\psi + \omega_\phi d\phi)^2 + f^{-1}(ds_0^2 + dt^2), \quad (6.1)$$

with ds_0^2 as in eq. (2.3) and

$$f^{-1} = 1 + \frac{Q - q^2}{2L^2}(x - y) - \frac{q^2}{4L^2}(x^2 - y^2), \quad (6.2)$$

$$\omega_\psi = \frac{3}{2}q(1 + y) + \frac{q}{8L^2}(1 - y^2)[3Q - q^2(3 + x + y)], \quad (6.3)$$

$$\omega_\phi = -\frac{q}{8L^2}(1 - x^2)[3Q - q^2(3 + x + y)]. \quad (6.4)$$

The $S^1 \times S^2$ horizon is localized at $y \rightarrow -\infty$, and asymptotic infinity at $x, y \rightarrow -1$. The Maxwell field $F = dA$ is determined by

$$A = \frac{\sqrt{3}}{2}f(dt + \omega_\psi d\psi + \omega_\phi d\phi) - \frac{\sqrt{3}}{4}q[(1 + x)d\phi + (1 + y)d\psi]. \quad (6.5)$$

The net electric charge and the local dipole magnetic charge are proportional to the positive parameters Q and q , respectively, which (for a physical interpretation) are assumed to satisfy $Q \geq q^2$ and $L < (Q - q^2)/(2q)$ [21]. The mass and angular momenta of the ring are

$$M = \frac{3\pi}{4}Q, \quad J_\psi = \frac{\pi}{8}q(6L^2 + 3Q - q^2), \quad J_\phi = \frac{\pi}{8}q(3Q - q^2). \quad (6.6)$$

In the limit $q = 0$ the black ring becomes a static charged naked singularity, solution of the pure Einstein-Maxwell theory. In order to boost the line element (6.1), we can follow a procedure almost identical to the one used for the vacuum ring. The standard mass rescaling of [4] together with the inequality $L < (Q - q^2)/(2q)$ suggests that during the boost we rescale the charges as

$$Q = \epsilon p_Q, \quad q = \epsilon p_q \quad (p_Q > 2Lp_q). \quad (6.7)$$

Omitting straightforward intermediate steps, in the case of a boost orthogonal to the plane (ξ, ψ) we obtain a shock pp -wave (4.1) with

$$H_\perp^s(x_2, y_1, y_2) = \frac{3\sqrt{2}p_Q}{\sqrt{(\xi + L)^2 + x_2^2}} K(k), \quad (6.8)$$

and k given by eq. (4.5). For a parallel boost, we obtain the metric (5.1) with

$$\begin{aligned} H_\parallel^s(x_1, x_2, y_2) = & 3\sqrt{2} \left[p_Q \frac{1}{a} + p_q \frac{y_2}{a} \left(1 - \frac{L^2 + \eta^2}{a^2 - y_2^2} \right) \right] K(k) + \\ & + \frac{3\sqrt{2}p_q}{2} \left[\frac{\eta^2 + L^2}{ay_2} \frac{a^2 + y_2^2}{a^2 - y_2^2} \Pi(\rho, k) - \pi \operatorname{sgn}(y_2) \right], \end{aligned} \quad (6.9)$$

where k and ρ as in eq. (5.6). To obtain the field of a boosted naked singularity ($q = 0$) just set $p_q = 0$ in eq. (6.9). Notice that the dipole charge q has an effect only in the case of a parallel boost, since p_q does not appear in H_{\perp}^s [which is in fact equivalent to the expression (4.9) for balanced vacuum rings]. This is related to the “asymmetry” between the angular momenta J_{ψ} and J_{ϕ} in eq. (6.6). In both boosts, one also finds that $F = dA$ tends to zero together with its associated energy-momentum tensor (so that the “peculiar configuration” of [13] does not arise here). In fact, both H_{\perp}^s and H_{\parallel}^s correspond to vacuum pp -waves. In principle, rescalings different from eq. (6.7) can be considered if one drops the requirement $L < (Q - q^2)/(2q)$. The detailed investigation of this and other possibilities is left for possible future work.

Acknowledgments

M.O. is supported by a post-doctoral fellowship from Istituto Nazionale di Fisica Nucleare (bando n.10068/03).

A. Results for the boosted $D = 5$ Myers-Perry black hole

Ref. [17] analyzed the ultrarelativistic boost of D -dimensional Myers-Perry black holes [16] with a single non-vanishing angular momentum. As in the present work, the calculation was performed in the case of two particular boosts orthogonal and parallel to the plane of rotation, and for $D = 5$ it resulted in impulsive pp -waves of the type (4.1) and (5.1), respectively. It is thus interesting to compare the results of [17] to ours. First of all, the angular momentum of black holes in $D = 5$ must obey a Kerr-like bound $a^2 < \mu$ [16]. Since, in the Aichelburg-Sexl limit, ref. [17] sent the mass parameter μ to zero while keeping the spin parameter a fixed, for $D = 5$ the final metrics refer to boosted naked singularities rather than black holes [17]. On the other hand, there is no upper limit on the spin of black rings [18], so that in our limit the rings do remain “black” until the final pp -wave is obtained (the same applies to the solutions of [17] in $D \geq 6$, when also black holes can be ultra-spinning). In the rest of this appendix we shall present the profile functions of [17] (for the case $D = 5$) using an explicit form adapted to our notation,⁴ and we shall compare them with our functions (4.4) and (5.5).

A.1 Orthogonal boost

For an orthogonal boost, the result of [17] can be rearranged as

$$\begin{aligned} \tilde{H}_{\perp}(x_2, y_1, y_2) = \frac{8\sqrt{2}p_M}{3\pi} & \left[\frac{2\sqrt{2}}{(\xi^2 + x_2^2 + L^2 + b^2)^{1/2}} K(k_1) + \frac{\sqrt{2}}{L^2} (\xi^2 + x_2^2 + L^2 + b^2)^{1/2} \times \right. \\ & \left. \times E(k_1) - \frac{2\sqrt{2}}{L^2} \frac{b^2}{(\xi^2 + x_2^2 + L^2 + b^2)^{1/2}} \Pi(\rho_1, k_1) \right], \end{aligned} \quad (\text{A.1})$$

⁴In particular, the quantity L will replace the original spin parameter a .

where

$$\begin{aligned}
 k_1 &= \left(\frac{\xi^2 + x_2^2 + L^2 - b^2}{\xi^2 + x_2^2 + L^2 + b^2} \right)^{1/2}, & \rho_1 &= -\frac{(\xi^2 + x_2^2 - L^2 - b^2)^2}{4L^2x_2^2}, \\
 b &= [(\xi^2 + x_2^2 - L^2)^2 + 4x_2^2L^2]^{1/4}.
 \end{aligned}
 \tag{A.2}$$

The above elliptic functions are singular for $k_1 = 1$, that is on a circle of radius L given by $x_2 = 0$, $\xi = L$. This was already remarked in [17] and it resembles our results of section 4. Other physical properties are more “hidden” in the expression (A.1). First of all, for $x_2 \rightarrow 0$ one has $\rho_1 \rightarrow 0$ if $\xi > L$, whereas ρ_1 diverges if $\xi < L$. This implies (with [19, identity (A5)]) that, when $\xi < L$ and x_2 is small, H_\perp contains a non-smooth term proportional to $|x_2|$, namely there is an additional membrane at $x_2 = 0$ and $\xi < L$ (i.e. within the ring singularity discussed above). The presence of such a disk-shaped source is related to the structure of the singularities of the Myers-Perry solutions [16], and it should be contrasted with the simpler profile function (4.9) for balanced black rings, which has only a “uniform” circle as a source. From a complementary viewpoint, we can compare an expansion of the profile function (A.1) at large spatial distances with the analogous result (4.8) for the black ring. From eq. (A.1) we obtain

$$\tilde{H}_\perp = \frac{1}{\sqrt{2}} \frac{8p_M}{3L} \left[3\frac{L}{r} - \frac{5}{8}(3\cos^2\theta - 1)\frac{L^3}{r^3} + \frac{7}{64}(35\cos^4\theta - 30\cos^2\theta + 3)\frac{L^5}{r^5} + O\left(\frac{L^7}{r^7}\right) \right].
 \tag{A.3}$$

The monopole term coincides with the one in the corresponding expression (4.8) for the black ring, which we should expect since we are boosting objects with the same mass (which scales as $M = \gamma^{-1}p_M$). However, eqs. (A.3) and (4.8) in general differ already in the quadrupole term, in particular for the physically most interesting case of balanced rings $p_\lambda = 2p_\nu$. They coincide only in the limiting case $p_\nu = 0$, corresponding to $\nu = 0$, when the black ring in fact reduces to a naked singularity isometric to that of Myers and Perry (see the discussion above about the Kerr bound). In addition, in the limit of vanishing rotation $L = 0$ of eq. (A.3) only the Aichelburg-Sexl monopole survives, which corresponds to the ultrarelativistic boost of the $D = 5$ Schwarzschild-Tangherlini black hole.⁵

A.2 Parallel boost

For a parallel boost, the profile function of [17] is

$$\begin{aligned}
 \tilde{H}_\parallel(x_1, x_2, y_2) &= \frac{8\sqrt{2}p_M}{3\pi} \left[\frac{4}{a} \left(1 + \frac{\eta^2 + y_2^2 + L^2 + a^2}{2Ly_2} \right) K(k) + \frac{2a}{L^2} E(k) - \right. \\
 &\quad \left. - \frac{2L + y_2}{a} \frac{\eta^2 + y_2^2 + L^2 + a^2}{L^2y_2} \Pi(\rho_1, k) \right],
 \end{aligned}
 \tag{A.4}$$

with a as in eq. (5.4), k as in eq. (5.6) and

$$\rho_1 = -\frac{\eta^2 + y_2^2 + L^2 - a^2}{2a^2}.
 \tag{A.5}$$

⁵Recall that, instead, balanced black rings can not be static, while unbalanced static rings correspond to setting $p_\nu = p_\lambda$ in eq. (4.8) [19] (and *not* $L = 0$).

Similarly as in section 5, the elliptic integrals are singular at $k = 1$, i.e. on a rod of length $2L$ located at $\eta = 0$, $|y_2| \leq L$ [17]. At large spatial distances, the expression (A.4) behaves as

$$\begin{aligned} \tilde{H}_{\parallel} = \frac{1}{\sqrt{2}} \frac{8p_M}{3L} \left[3\frac{L}{r} + 2\cos\theta \frac{L^2}{r^2} + \frac{7}{8}(3\cos^2\theta - 1)\frac{L^3}{r^3} + \frac{3}{4}(5\cos^3\theta - 3\cos\theta)\frac{L^4}{r^4} + \right. \\ \left. + \frac{11}{64}(35\cos^4\theta - 30\cos^2\theta + 3)\frac{L^5}{r^5} + O\left(\frac{L^6}{r^6}\right) \right]. \end{aligned} \quad (\text{A.6})$$

The discussion is similar as the one above for \tilde{H}_{\perp} . Again, the monopole term coincides with the one in the corresponding expression (5.9) for the black ring. Higher multipoles in general differ, in particular for balanced rings. Boosted black holes reduce to the Aichelburg-Sexl monopole in the static limit $L = 0$.

References

- [1] G. 't Hooft, *Graviton dominance in ultrahigh-energy scattering*, *Phys. Lett.* **B 198** (1987) 61.
- [2] D. Amati and C. Klimčík, *Strings in a shock wave background and generation of curved geometry from flat space string theory*, *Phys. Lett.* **B 210** (1988) 92.
- [3] H.J. de Vega and N. Sánchez, *Quantum string scattering in the Aichelburg-Sexl geometry*, *Nucl. Phys.* **B 317** (1989) 706.
- [4] P.C. Aichelburg and R.U. Sexl, *On the gravitational field of a massless particle*, *Gen. Rel. Grav.* **2** (1971) 303.
- [5] T. Banks and W. Fischler, *A model for high energy scattering in quantum gravity*, [hep-th/9906038](#).
- [6] R. Emparan, G.T. Horowitz and R.C. Myers, *Black holes radiate mainly on the brane*, *Phys. Rev. Lett.* **85** (2000) 499 [[hep-th/0003118](#)].
- [7] S.B. Giddings and S.D. Thomas, *High energy colliders as black hole factories: the end of short distance physics*, *Phys. Rev.* **D 65** (2002) 056010 [[hep-ph/0106219](#)].
- [8] S. Dimopoulos and G. Landsberg, *Black holes at the LHC*, *Phys. Rev. Lett.* **87** (2001) 161602 [[hep-ph/0106295](#)].
- [9] V. Cardoso, E. Berti and M. Cavagliá, *What we (don't) know about black hole formation in high-energy collisions*, *Class. and Quant. Grav.* **22** (2005) L61 [[hep-ph/0505125](#)].
- [10] D.M. Eardley and S.B. Giddings, *Classical black hole production in high-energy collisions*, *Phys. Rev.* **D 66** (2002) 044011 [[gr-qc/0201034](#)].
- [11] H. Yoshino and Y. Nambu, *High-energy head-on collisions of particles and hoop conjecture*, *Phys. Rev.* **D 66** (2002) 065004 [[gr-qc/0204060](#)].
- [12] E. Kohlrath and G. Veneziano, *Black holes from high-energy beam-beam collisions*, *JHEP* **06** (2002) 057 [[gr-qc/0203093](#)].
- [13] C.O. Loustó and N. Sánchez, *The curved shock wave space-time of ultrarelativistic charged particles and their scattering*, *Int. J. Mod. Phys.* **A 5** (1990) 915.
- [14] M. Ortaggio, *Ultrarelativistic black hole in an external electromagnetic field and gravitational waves in the Melvin universe*, *Phys. Rev.* **D 69** (2004) 064034 [[gr-qc/0311088](#)].

- [15] M. Ortaggio, *Higher dimensional black holes in external magnetic fields*, *JHEP* **05** (2005) 048 [[gr-qc/0410048](#)].
- [16] R.C. Myers and M.J. Perry, *Black holes in higher dimensional space-times*, *Ann. Phys. (NY)* **172** (1986) 304.
- [17] H. Yoshino, *Lightlike limit of the boosted Kerr black holes in higher-dimensional spacetimes*, *Phys. Rev. D* **71** (2005) 044032 [[gr-qc/0412071](#)].
- [18] R. Emparan and H.S. Reall, *A rotating black ring in five dimensions*, *Phys. Rev. Lett.* **88** (2002) 101101 [[hep-th/0110260](#)].
- [19] M. Ortaggio, P. Krtouš and J. Podolský, *Ultrarelativistic boost of the black ring*, *Phys. Rev. D* **71** (2005) 124031 [[gr-qc/0503026](#)].
- [20] R. Emparan and H.S. Reall, *Generalized Weyl solutions*, *Phys. Rev. D* **65** (2002) 084025 [[hep-th/0110258](#)].
- [21] H. Elvang, R. Emparan, D. Mateos and H.S. Reall, *A supersymmetric black ring*, *Phys. Rev. Lett.* **93** (2004) 211302 [[hep-th/0407065](#)].
- [22] H. Elvang, R. Emparan, D. Mateos and H.S. Reall, *Supersymmetric black rings and three-charge supertubes*, *Phys. Rev. D* **71** (2005) 024033 [[hep-th/0408120](#)].
- [23] I. Bena and N.P. Warner, *One ring to rule them all . . . and in the darkness bind them?*, [hep-th/0408106](#).
- [24] J.P. Gauntlett and J.B. Gutowski, *General concentric black rings*, *Phys. Rev. D* **71** (2005) 045002 [[hep-th/0408122](#)].
- [25] R. Emparan, *Rotating circular strings and infinite non-uniqueness of black rings*, *JHEP* **03** (2004) 064 [[hep-th/0402149](#)].
- [26] V. Pravda and A. Pravdová, *WANDs of the black ring*, *Gen. Rel. Grav.* **37** (2005) 1277.